















supplementary foods was minimum means 25% only, whereas 75% mothers have no knowledge about them. Thus the growth & development may be declined if they do not taking sufficient breast milk. The factors responsible for stopping the breastfeeding are 5% due to lack of sufficient milk secretion & 70% had stopped breast feeding due to absence of sucking reflex. In terms of infant formula milk, 10% mothers think that it is similar to breast, 72% mothers were not agreed, whereas 18% were confused either it is similar or not. Some nursing devices are available in the market which may be beneficial for cleft lip & cleft palate babies but here no respondents had any awareness regarding nursing devices.

## References

- [1] Amstalden-M L, Magna, L. A., & Gil-da-Silva-Lopes. L. (2007). Neonatal care of infants with cleft lip and/or palate: Feeding orientation and evolution of weight gain in a no specialized Brazilian hospital. *Cleft Palate-Craniofacial Journal*, 44, 329-334
- [2] Bushra f, Ahmad N, Hussain T, Manan F, Effect of infant feeding practices on birth interval and morbidity among children in North West Province of Pakistan. *Journal of Islamic academy of science* 1995; 8:69-72.
- [3] Chezem, J., C. Friesen, and J. Boettcher, Breastfeeding knowledge, breastfeeding confidence, and infant feeding plans: effects on actual feeding practices. *JOGNN: Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 2003. 32(1): p. 40-47.
- [4] Clarren, S., Anderson, B., & Wolf, L. (1987). Feeding infants with cleft lip and cleft palate, *cleft palate journal*, 24,244-249.
- [5] Clarren, S., Anderson, B., & Wolf, L. (1987). Feeding infants with cleft lip, cleft palate, *cleft palate journal*, and 24,244-249.
- [6] *Craniofacial Journal*, 40, 84-87. Dalben, G. D., Costa, B., Ribeiro-Gomide, M., & Teixeira das Neves, L. (2003). Breast-feeding and sugar intake in babies with cleft lip and palate.
- [7] Danner SC. Breast feeding the infants with a cleft defect. *NAACOGS Clin Issu Perinat Womens Health*, Nurs1992; 3: 634-9.
- [8] Dennis, C., et al., The effect of peer support on breast-feeding duration among primiparous women: a randomized controlled trial. *Canadian Medical Association Journal*, 2002. 166(1): p. 21-28.
- [9] Donath, S. and L.H. Amir, Rates of breastfeeding in Australia by state and socioeconomic status: evidence from the 1995 National Health Survey. *Breastfeeding Review*, 2000. 8(3): p. 23-7.
- [10] Kulkarni RN, Ajenaya s, Gujar R. Breast feeding practices in an urban community of Kalamboli, Navi Mumbai. *Indian journal of community medicine* 2004;
- [11] Lande, B., et al., *Infant feeding practices and associated factors in the first six months of life: the Norwegian infant nutrition survey*. *Acta Paediatrica*, 2003. 92(2): p. 152-61.
- [12] Mizuno, K. (2002). Feeding behaviors of infants with cleft lip and palate. *Acta Paediatr*, 91, 1227-1232. Reid, J. (2004). A review of feeding intervention for infants with cleft palate. *Cleft Palate-Craniofacial Journal*, 31, 268-278.
- [13] National Health and Medical Research Council, Dietary Guidelines for Children and Adolescents in Australia incorporating the Infant Feeding Guideline for Health Workers. 2003, Canberra: Commonwealth of Australia.
- [14] Pincombe, J., et al., Baby Friendly Hospital Initiative (BFHI) Practices and breastfeeding duration in first time mothers in an Adelaide teaching maternity hospital, Australia. in press, 2006.