





control and overall survival. In the future we hope it could be possible. Also we emphasize the careful patient's selection and their treatment in specialized centers.

## 5. Conclusions

Our study demonstrated that also in developing countries reirradiation is a possible option of treatment for patients with in field recurrence. It improves quality of life with acceptable side effects. However we should take into account that we treated a small number of patients from which clear conclusions cannot be drawn. Prospective randomized studies should be planned in the future to decide the optimal fractionation and total reirradiation dose to the tumor with less side effects in normal tissues. Installation of new and more advanced radiotherapy machines in our department will give patients the opportunity for a curative treatment with a better local control and overall survival.

## Reference

- [1] Lavey RS, Taylor J, Tward JD et al, The extent, time course, and fraction size dependence of mouse spinal cord recovery from radiation injury. *Int J Radiat Oncol Biol Phys.* 1994;30:609–617
- [2] Ang KK, Jiang GL, Feng Y, et al. Extent and kinetics of recovery of occult spinal cord injury. *Int J Radiat Oncol Biol Phys.* 2001;50:1013–1020.
- [3] Lee AW, Foo W, Law SC, Reirradiation for recurrent nasopharyngeal carcinoma: factors affecting the therapeutic ratio and ways for improvement. *Int J Radiat Oncol Biol Phys.* 1997; 38:43–52
- [4] Kruse JJ, Stewart FA Gene expression arrays as a tool to unravel mechanisms of normal tissue radiation injury
- [5] and prediction of response. *World J Gastroenterol* 2007 13:2669–2674
- [6] Alsner J, Andreassen CN, Overgaard J. Genetic markers for prediction of normal tissue toxicity after radiotherapy. *Semin Radiat Oncol.* 2008 18:126–135
- [7] Valdagni R, Rancati T, Fiorino C. Predictive models of toxicity with external radiotherapy for prostate cancer: clinical issues. *Cancer.* 2009 115:3141–3149
- [9] Joseph KJ, Al-Mandhari Z, Pervez N et al Reirradiation after radical radiation therapy: a survey of patterns of practice among Canadian Radiation Oncologists. *Int J Radiat Oncol Biol Phys* 2008; 72,1523–1529
- [10] Dresen RC, Gosens MJ, Martijn H, et al. Radical resection after IORT-containing multimodality treatment is the most important determinant for outcome in patients treated for locally recurrent rectal cancer. *Ann Surg Oncol* 2008;15:1937–47.
- [11] Bouchard P, Efron J. Management of recurrent rectal cancer. *Ann Surg Oncol* 2010;17:1343–56.
- [12] Das P, Delclos M, Skibber J, Rodriguez-Bigas M et al. Hyperfractionated accelerated radiotherapy for rectal cancer in patients with prior pelvic irradiation. *Int J Radiat Oncol Biol Phys* 2010; 77; 60-65
- [13] Baujat B, Bourhis J, Blanchard P et al. Hyperfractionated or accelerated radiotherapy for head and neck cancer. *Cochrane Database Syst Rev.* 2010; 8;(12)
- [14] Van den Hout WB, Van der Linden YM, Steenland E, et al. Single- versus multiple-fraction radiotherapy in patients with painful bone metastases: Cost-utility analysis based on a randomized trial. *J Natl Cancer Inst* 2003;95:222-229
- [15] Wong CS, Van Dyk J, Milosevic M, Laperriere NJ, Radiation myelopathy following single courses of radiotherapy and retreatment. *Int J Radiat Oncol Biol Phys* 1994;30:575–81
- [16] Maranzano E, Bellavita R, Floridi P, Radiation-induced myelopathy in long-term metastatic spinal cord compression patients after hypofractionated radiotherapy: a clinical and magnetic resonance imaging analysis. *Radiother Oncol* 2001;60:281–8.
- [17] Nieder C, Grosu AL, Andratschke NH, Molls M, Update of human spinal cord reirradiation tolerance based on additional data from 38 patients. *Int J Radiat Oncol Biol Phys* 2006; 66, 5,1446–1449