

Unfortunately, in Bulgaria there is still no team approach in the work among anaesthesiologists-reanimators and psychologists; this poses several questions for the active role of psychologist in pre- and post-operative preparation and care for these children.

References

- [1] Isaksen, J., P. Holth, An operant approach to teaching joint Attention skills to children with autism, *Behavioral Interventions*, 2009, 24: 215–236.
- [2] www.doctorbg.com
- [3] Levterova, D. Actual problems of special education, Pd., 2002, p. 106-108/.
- [4] Levterova, D. Actual problems of special education, Pd., 2002, p. 104-105/.
- [5] Morales, R. Autism fear. New unique approach, S., 2009, p. 21-23/
- [6] Levterova, D, p 105; E. McGuire A., R. Michalko, *Minds Between Us: Autism, mindblindness and the uncertainty of communication*, Educational Philosophy and Theory, Vol. 43, No. 2, 201, p. 162-177.
- [7] McGuire E. A., Michalko R., *Minds Between Us: Autism, mindblindness and the uncertainty of communication*, *Educational Philosophy and Theory*, Vol. 43, No. 2, 201, p. 162-177
- [8] Luis L. Special Diets for Special Kids. 2010.
- [9] Dillenburger, K., M. Keenan, A. Doherty, T. Byrne, St. Gallagher, Living with children diagnosed with autistic spectrum disorder: parental and professional views, *British Journal of Special Education*, 2010, Vol. 37, Issue 1, p. 13-23.
- [10] Rainey L, van der Walt JH. The anaesthetic management of autistic children. *Anaesth Intensive Care*. 1998; 26: 682-686.
- [11] Proczkowska-Björklund M, Svedin CG. Child related background factors affecting compliance with induction of anaesthesia. *Paediatr Anaesth*. 2004; 14: 225-234.
- [12] Judith Alison Short. Anaesthesia for children with special needs, including autistic spectrum disorder. *Continuing Education in Anaesthesia, Critical Care & Pain Advance Access published January 29, 2013*.
- [13] www.pecs.org.uk
- [14] Karam VY, Barakat H. Perioperative management of the child with behavioral disorders.
- [15] Bozkurt P. Premedication of the pediatric patient - anesthesia for the uncooperative child. *Curr Opin Anaesthesiol*. 2007; 20: 211-215.
- [16] Ghai B, Grandhe RP, Kumar A, Chari P. Comparative evaluation of midazolam and ketamine with midazolam alone as oral premedication. *Paediatr Anaesth*. 2005; 15: 554-559.
- [17] Filatov SM, Baer GA, Rorarius MG, Oikkonen M. Efficacy and safety of premedication with oral ketamine for day-case adenoidectomy compared with rectal diazepam/diclofenac and EMLA. *Acta Anaesthesiol Scand*. 2000; 44: 118-124.
- [18] Christiansen E, Chambers N. Induction of anesthesia in a combative child; management and issues. *Paediatr Anaesth*. 2005; 15: 421-425.
- [19] Darlong V, Shende D, Subramanian MS, Sunder R, Naik A. Oral ketamine or midazolam or low dose combination for premedication in children. *Anaesth Intensive Care*. 2004; 32:246-9.
- [20] Shailesh Shah, Sonia Shah, Jesus Apuya, Senthil Gopalakrishnan, Timothy Martin. Combination of oral ketamine and midazolam as a premedication for a severely autistic and combative patient. *Journal of Anesthesia* 2009; 23:126-128
- [21] Thompson DG, Tielsch-Goddard A. Improving Management of Patients With Autism Spectrum Disorder Having Scheduled Surgery: Optimizing Practice. 2013.
- [22] Greenberg B. MD, Schoonover J. NP, Jedlicki K. NP, Graf A. NP, Overbey E. MD, et al (2010). Intranasal Combination Dexmedetomidine and Midazolam for Pediatric Procedural Sedation: A New and Palatable Approach. Paper Presented at SPA meeting in San Antonio.
- [23] Zub D1, Berkenbosch JW, Tobias JD. Preliminary experience with oral dexmedetomidine for procedural and anesthetic premedication. *Paediatr Anaesth*. 2005; 15: 932-938.
- [24] Alyssa Davi. Anesthesia and Sedation Risks in Children Labeled with Autistic Spectrum Disorder. P. 1-2.