

References

- [1] Ashraf, M., Gohil, R.N. 1986. Chromosome number reports LXXXXI. Taxon 35: 408.
- [2] Baquar, S.R., Abid Askari, S.H. 1970. Chromosome numbers in some flowering plants of west Pakistan. Genet. Iber. 22: 1-11.
- [3] Chauhan, N.S. 2003. Important medicinal and aromatic plants of Himachal Pradesh. Indian Forester 129: 979-998.
- [4] Haroun, S.A., Al Shehri, A.M., Al Wadie, H.M. 2004. Cytomixis in the microsporogenesis of *Vicia faba* L. (Fabaceae). Cytologia 69: 7-11.
- [5] Kumar, P., Singhal, V.K., Kaur, D. 2012. Impaired male meiosis due to irregular synapsis coupled with cytomixis in a new diploid cytotype of *Dianthus angulatus* (Caryophyllaceae) from Indian cold deserts. Folia Geobot. 47: 59-68. doi: 10.1007/s12224-011-9107-8
- [6] Mehra, P.N., Dhawan, H. 1971. In IOPB chromosome number reports XXXIV Taxon 20: 785-797.
- [7] Mabberley, D.J. 1997. The plant book, 2nd ed. Cambridge University Press, Cambridge, UK.
- [8] Rana, P.K., Kumar, P., Singhal V.K. 2014 Cytomixis and associated abnormalities during male meiosis in *Lindlofia longiflora* var. *falconeri* (Boraginaceae) Cytologia 79(4): 535-540.
- [9] Rawat, D.S., Kharwal, A.D. 2011. Traditional health practices by 'kinners'- a tribe in alpine and sub-alpine Himalayas of Kinnaur (Himachal Pradesh), India. Life sciences Leaflets 22: 1048-1055.
- [10] Rundel, R.W. 1989. Ecological success in relation to plant form and function in the woody legumes. In C. H. Stirton and J. L. Zarucchi [eds.], Advances in legume biology, Monographs in Systematic Botany from the Missouri Botanical Garden 29: 377-398.
- [11] Sareen, T.S., Singh, P.D. 1976. Cytological studies in some north Indian Leguminosae. Proc. Indian Sci. Congr. Assoc. 93: 122-123.
- [12] Singhal, V.K., Kaur, D. 2011. Cytomixis inducing meiotic irregularities and pollen malformation in *Clematis graveolens* Lindley from the cold deserts of Kinnaur districts of Himachal Pradesh, India. Cytologia 76: 319-327.

