





- strains isolated from human faeces. *Anaerobe* 18(1):96-102.
- [8] **Funmilola A, Ayenia B, Borja Sáncheza, Bolanle A, Adeniyib, Clara G. de los Reyes-Gavilána, Abelardo M, Patricia Ruas-Madiedo** (2011). Evaluation of the functional potential of *Weissella* and *Lactobacillus* isolates obtained from Nigerian traditional fermented foods and cow's intestine. *International Journal of Food Microbiology* 147(2):97–104.
- [9] **Gerald W. T** (1999). Identification of Lactobacilli and Bifidobacteria. *Current Issues in Molecular Biology* 1(1): 53-64.
- [10] **Chin M. L, Chin C. S., Norhani A. and Yin W. H** (2008). Estimation of 16S rRNA gene copy number in several probiotic *Lactobacillus* strains isolated from the gastrointestinal tract of chicken. *FEMS Microbiology Letters* 287(1):136–141.
- [11] **Clarridge J. E. III** (2004). Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. *Clinical Microbiology Reviews* 17(4):840–862.
- [12] *Bergey's manual of systematic bacteriology; Volume 3: The Firmicutes: (464-513); Springer-Verlag, New York. 2009.*
- [13] **Collins C, Grange J, Lyne P, Falkinham J III** (2004). *Collins and Lyne's Microbiological methods. 8<sup>th</sup> Edition; CRC Press.*
- [14] **Harrigan W. F** (2013). *Laboratory methods in food Microbiology. 3<sup>rd</sup> edition; Academic Press; First Indian Print.*
- [15] **Saitou N, Nei M** (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4(4):406-25.
- [16] **Joseph F** (1985). Confidence limits on phylogenies: An approach using the Bootstrap. *Evolution* 39(4):783-791.
- [17] **Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S** (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* (10):2731-9.
- [18] **Gillian E. G, Christine H, Andrew W. B, Beuerman D, and Gregor R** (2002). Persistence of *Lactobacillus fermentum* RC-14 and *Lactobacillus rhamnosus* GR-1 but not *L rhamnosus* GG in the human vagina as demonstrated by randomly amplified polymorphic DNA. *Clinical and Diagnostic Laboratory Immunology* 9(1): 92–96.
- [19] **Pan, Dao D, Xia Q. Z, and Tian Y** (2011). Characterization of *Lactobacillus fermentum* SM-7 isolated from koumiss, a potential probiotic bacterium with cholesterol lowering effects. *Journal of the Science of Food and Agriculture* 91(3): 512-518.
- [20] **John-Hwan, Park, Yeonhee L, Enpyo M, Seun H S, and Min-Won B** (2005). Safety assessment of *Lactobacillus fermentum* PL 9005, a potential probiotic lactic acid bacterium, in mice. *Journal of Microbiology and Biotechnology* 15(3): 603-608.
- [21] **Zeng, Xian Q, Dao D. P, and Pei D. Z** (2010). Functional characteristics of *Lactobacillus fermentum* F1. *Current Microbiology* 62(1): 27-31.
- [22] **Marika M and Mihkel Z** (2009). *Lactobacillus fermentum* ME-3 – an antimicrobial and antioxidative probiotic. *Microbial Ecology in Health and Disease* 21(1): 1-27.
- [23] **Yeonhee L** (2005). Characterization of *Weissella kimchii* PL9023 as a potential probiotic for women *FEMS Microbiology Letters* 250(1):157–162.