











## 6. Acknowledgments

This modest work geared to support human category, which suffer from chronic diseases, I wish to commend the patients who participated and understand the value of work. The authors wish to thank all the thalassemic patients who participated in this study; clinical laboratory of at "AL-Emam AL-Sajisd hospital " in Najaf city, (Iraq) for execution the laboratory analyses.

## References

- [1] (A. J. Kadhum Antioxidant and lipidperoxidation in chronic renal failure patients Coll.of Sci./Univ.of Al-Qadisiya AL-Qadisiya Journal of Vet. Med. Sci. Vol. /7 No./ 2 2008)
- [2] (David S. Baldwin, MD: A Legacy in Nephrology Paul L. Kimmel, Joel Neugarten, and Jerome Lowenstein JASN March 2015 26: 531-535; published ahead of print August 22, 2014, doi:10.1681/ASN.2014030305)
- [3] (Dr Donald Richardson, Consultant Nephrologist, York District Hospital NIH Publication No. 14-4619 May 2014)
- [4] (Michael Hsieh, David A Power bnormal Renal Function and Electrolyte Disturbances in Older People J Pharm Pract Res 2009; 39: 230-4)
- [5] (Pedro Iglesias1 and Juan J. Díez2Insulin therapy in renal disease Department of Endocrinology, Hospital General, Segovia, Spain.Department of Endocrinology, Hospital Ramo'n y Cajal, Madrid, Spain) 0.1111/j.1463-1326.2007.00802.x.
- [6] Artiss JD, Vinogradov S, Zak B. Spectrophotometric study of several sensitive reagents for serum iron.Clin Biochem. 1981;14:311-315. [PubMed]
- [7] Biesenbach G, Raml A, Schmekal B, Eichbauer-Sturm G. Decreased insulin requirement in relation to GFR in nephropathic type 1 and insulin-treated type 2 diabetic patients. Diabet Med 2003; 20: 642-645
- [8] Centers for Disease Control and Prevention (CDC). Incidence of end-stage renal disease among persons with diabetes – United States, 1990-2002. MMWR Morb Mortal Wkly Rep 2005; 54: 1097-1100.
- [9] Centers for Disease Control and Prevention (CDC).Incidence of end-stage renal disease among persons with diabetes – United States, 1990-2002. MMWR
- [10] Coward RJ, Welsh GI, Yang J et al. The human glomerular podocyte is a novel target for insulin action. Diabetes 2005; 54: 3095-3102.
- [11] Freeman V, Arneson W. Hemoglobin production, disorders, and testing. In: Anderson W, Brickell J, editors. Clinical Chemistry: A Laboratory Prospective. Philadelphia: FA Daves Co.; 2007. p. 185.
- [12] Hostetter TH, Rennke HG, Brenner BM. The case for intrarenal hypertension in the initiation and progression of diabetic and other glomerulopathies. Am J Med 1982; 72: 375-380.
- [13] Iron Panel of the International Committee for Standardization in Hematology The measurement of total and unsaturated iron-binding capacity in serum. Br J Haematol. 1978;38:281-290. [PubMed]
- [14] Maeda M, Yabuki A, Suzuki S, Matsumoto M, Taniguchi K, Nishinakagawa H. Renal lesions in spontaneous insulin-dependent diabetes mellitus in the nonobese diabetic mouse: acute phase of diabetes. Vet Pathol 2003; 40: 187-19
- [15] Maeda M, Yabuki A, Suzuki S, Matsumoto M, Taniguchi K, Nishinakagawa H. Renal lesions in spontaneous insulin-dependent diabetes mellitus in the nonobese diabetic mouse: acute phase of diabetes. Vet Pathol 2003; 40: 187-195.]
- [16] Morb Mortal Wkly Rep 2005; 54: 1097-1100
- [17] Morgan EH. Transferrin. In: Haerberli A, editor. Human Protein Data, Part B. Weinheim: Wiley-VCH; 1992. p. 291.
- [18] Morioka T, Emoto M, Tabata T et al. Glycemic control is a predictor of survival for diabetic patients on hemodialysis. Diabetes Care 2001; 24: 909-913.]
- [19] Rabkin R, Ryan MP, Duckworth WC. The renal metabolism of insulin. Diabetologia 1984; 27: 351-357.
- [20] Rayssiguier Y, Mbega JD, Durlach V, et al. Magnesium and blood pressure. I. Animal studies. Book of abstracts V1th International Mg Symposium abstracts. Indore, India, Dec 2-6, 1991.
- [21] Scholey JW, Meyer TW. Control of glomerular hypertension by insulin administration in diabetic rats.J Clin Invest 1989; 83: 1384-1389].
- [22] Wu MS, Yu CC, Wu CH, Haung JY, Leu ML, Huang CC. Pre-dialysis glycemic control is an independent predictor of mortality in type II diabetic patients on continuous ambulatory peritoneal dialysis. Perit Dial Int 1999; 19 (Suppl. 2) : S179-S183.]