









- [25] Simpson DA, Furness CA, Hodkinson TR, Muasya AM, Chase MW (2003) Phylogenetic relationships in Cyperaceae subfamily Mapanioideae inferred from pollen and plastid DNA sequence data. *Amer J Bot* 90:1071–1086
- [26] Simpson DA, Muasya AM, Alves M, Bruhl JJ, Dhooge S, Chase MW, Furness CA, Ghamkhar K, Goetghebeur P, Hodkinson TR, Marchant AD, Nieuborg R, Reznicek AA, Roalson EH, Smets E, Starr JR, Thomas WW, Wilson KL, Zhang X (2008) Phylogeny of Cyperaceae based on DNA sequence data—a new rbcL analysis. In: *MonocotsIII/Grasses IV*. Claremont, CA: Aliso 23: 72–83
- [27] Smith-White S (1959) Pollen development patterns in the Epacridaceae. *Proc Linn Soc NSW* 84:8–35
- [28] Strandhede SO (1973) Pollen development in *Eleocharis Palustris* Group (Cyperaceae). 2. Cytokinesis and microspore degeneration. *Bot Not* 126:255–265
- [29] Sultan, S., Perveen, A & Qaiser, M. 1994. A Palynological study of monocots from Karachi (excluding Gramineae ) – *Pakistan Journal of Botany*. 26 (1) : 21-34.
- [30] Van Wichelen J, Camelbeke K, Chaerle P, Goetghebeur P, Huysmans S (1999) Comparison of different treatments for LM and SEM studies and systematic value of pollen grains in Cyperaceae. *Grana* 38:50–58
- [31] Wodehouse RP (1935) Pollen grains. Their structure identification and significance in science and medicine. Hafner, New York