







- [10] Fernandez I (1987) Contribution al conocimiento palinologico de Cyperaceae.
- [11] Acta Bot Malacitana 12:173–182
- [12] Furness CA, Rudall PJ (1999) Microsporogenesis in monocotyledons. *Ann Bot* 84:475–499.
- [13] Goetghebeur, P. (1986) *Genera Cyperacearum*. Een bijdrage tot de Kennis van de morfologie, systematick en fylogense van de Cyperaceae genera – DPhil. Thesis. State University of Ghent, Belgium.
- [14] Goetghebeur P (1998) Cyperaceae. In: Kubitzki K (ed) *The families and genera of vascular plants. Flowering plants-monocotyledons*, vol 4. Springer, Berlin, pp 141–190
- [15] Govaerts R, Simpson DA, Goetghebeur P, Wilson K, Egorova T, Bruhl JJ (2007) World checklist of Cyperaceae. The Board of Trustees of the Royal Botanic Gardens, Kew. Available at <http://www.kew.org/wcsp/monocots/>, Accessed on 1 October 2007
- [16] Haines RW, Lye KA (1983) *The sedges and rushes of East Africa*. African Natural History Society, Nairobi
- [17] Koyama T. 1961 Classification of family Cyperaceae 1. *J. Fac. Sci. Univ. Tokyo* 8 (3). 37-148.
- [18] Kunjalwar S.G.(2008 ) Scanning electron microscopy as aid to the taxonomy of sedges: sporoderm and spermoderm pattern in some Cyperaceae of Nagpur District, Ph.D. Thesis Rashtrasant Tukadoji Maharaj Nagpur Univrsity Nagpur India
- [19] Kuprianova, L.A. 1948 Pollen morphology and the phylogeny of the monocotyledons. *Comm. Komorrow. Indst. Acad. Sci.* 1:7.
- [20] Muasya AM, Simpson DA, Verboom GA, Goetghebeur P, Naczi RFC, Chase MW, Smets E (2008) Phylogeny of cyperaceae based on DNA sequence data: current progress and future prospects.
- [21] Nair P. K. K. (1970) *Pollen morphology of Angiosperms*. Vikas Publication, Delhi.
- [22] Padhye M.D. (1966-67) The pollen grains of *Kyllinga Rottb.* *Palynological Bull.* vol. II & III. pp. 101 – 103.
- [23] Padhye MD, Makde KH (1980) Pollen morphology of Cyperaceae. *J Palynol* 16:71–81
- [24] Selling OH (1947) Studies in the Hawaiian pollen statistics, Part II. The pollens of the Hawaiian phanerograms. *Bulletin of the Bishop Museum.* Honolulu 38:1–360
- [25] Shah CK (1962) Pollen development in some members of the Cyperaceae. *Plant embryology—a symposium.* CSIR, New Delhi, pp 81–93
- [26] Sharma, M. 1967 Pollen morphology of Indian Monocotyledons *J. Palynol* Special Volume.
- [27] Simpson DA, Furness CA, Hodkinson TR, Muasya AM, Chase MW (2003) Phylogenetic relationships in Cyperaceae subfamily Mapanioideae inferred from pollen and plastid DNA sequence data. *Amer J Bot* 90:1071–1086
- [28] Simpson DA, Muasya AM, Alves M, Bruhl JJ, Dhooge S, Chase MW, Furness CA, Ghamkhar K, Goetghebeur P, Hodkinson TR, Marchant AD, Nieuborg R, Reznicek AA, Roalson EH, Smets E, Starr JR, Thomas WW, Wilson KL, Zhang X (2008) Phylogeny of Cyperaceae based on DNA sequence data—a new rbcL analysis. In: *MonocotsIII/Grasses IV*. Claremont, CA: Aliso 23: 72–83
- [29] Smith-White S (1959) Pollen development patterns in the Epacridaceae. *Proc Linn Soc NSW* 84:8–35
- [30] Strandhede SO (1973) Pollen development in *Eleocharis Palustris* Group (Cyperaceae). 2. Cytokinesis and microspore degeneration. *Bot Not* 126:255–265
- [31] Van Wichelen J, Camelbeke K, Chaerle P, Goetghebeur P, Huysmans S (1999) Comparison of different treatments for LM and SEM studies and systematic value of pollen grains in Cyperaceae. *Grana* 38:50–58
- [32] Wodehouse RP (1935) *Pollen grains. Their structure identification and significance in science and medicine.* Hafner, New York