







- [2] Franchini E, Yuri Takito M, Yuzo Nakamura F, Ayumi Matsushigue K, Peduti Dal'Molin Kiss MA, "Effects of Recovery type after a Judo combat on Blood Lactate Removal and on Performance in an intermittent task," *Journal of Sports Medicine and Physical Fitness*, 43(4): 424-31, 2003.
- [3] Jeffreys, I, "A multidimensional approach to enhancing recovery," *Strength and Conditioning Journal*. 27(5): 78-85, 2005.
- [4] Kinugasa T., and Kilding AE, "A Comparison of Post-Match Recovery Strategies in Youth Soccer players", *Journal of Strength and Conditioning Research*, 23 (5), 1402-1407, 2009.
- [5] Signorile FJ, Ingalls C, Tremblay LM, "The Effects of Active and Passive Recovery on Short-Term, High Intensity power output," *Canadian Journal of Applied Physiology*, 18(1), 31-42, 1993.
- [6] Spierer DK, Goldsmith R, Baran DA, Hryniewicz K, Katz SD, "Effect of Active versus Passive Recovery on Work performed during serial supramaximal exercise tests", *International Journal of Sports Medicine*, 25(2), 109-14, 2004.
- [7] Tessitore A, Meeusen R, Cortis C, Capranica L, "Effects of different Recovery interventions on Anaerobic Performances following preseason soccer training," *The Journal of Strength & Conditioning Research*, 21(3): 745-50, 2007.
- [8] Weltman A, Stamford BA, Fulco C, "Recovery from Maximal Effort Exercise: Lactate disappearance and Subsequent Performance", *Journal of Applied Physiology: Respiratory Environmental and Exercise Physiology*, 47(4), 677-82, 1979.
- [9] Verma, J.P, *Statistical Methods for Sports and Physical Education*, Tata Mcgraw Hill Education Private Limited, New Delhi, 2011.