

Standardization of Explant Bud Break in *Melia dubia* - Australian Teak Using Tissue Culture Techniques

R. Aishwarya¹, V. G. Saravana Kumar²

¹Student, Department of Biotechnology, St. Michaels College of Engineering and Technology, Kalayarkoil

²Associate Professor, Department of Biotechnology, St. Michaels College of Engineering and Technology, Kalayarkoil

Corresponding Author: R. Aishwarya

Abstract: *Meliadubia* is the fast growing multipurpose tree species, Australian Teak wood variety, also called as a Mahaneem or Forest neem, which has a great demand in Wood industry. The harvesting period is 6-7 years unlike the commercial Burma teak wood variety *Tectonagrandis* which takes almost a generation (60-70 years). Moreover, the Australian teak wood is cheaper when compared to Burma Teak wood. Conventional growth of *Meliadubia* through seeds is not much successful where the germination rate is very poor of about 14-34% only. So, *Invitro* propagation of the *Meliadubia* is very essential for large scale propagation of the species which also possess therapeutic properties. Also, *Meliadubia* can generate substantial power, the yellow mature stem, with more fibre and lignin content yielded around 200 dry metric tonnes per hectare. This study is aimed at the Standardization of Initiation stage of *Meliadubia* explants through *Invitro* propagation. Comparison of best combination of growth regulator was identified. The developed hardened sapling through *Invitro* propagation can be harvested in 5 years which then will be useful for various purposes. Through this technique, the mass propagation is simple and effective, propagated plant will be like mother who is identified the best and disease free.

Keywords: *Meliadubia*, *Invitro* propagation, Initiation, Mass propagation

1. Introduction

Meliadubia is a good energy source crop which allows more diversity than bamboo species. It is commonly found in the hills. It provides a rapid growth. The extract from different parts of the *Meliadubia* plant possess antiviral, antibacterial, antifungal, antidiabetic, antineoplastic, antihelmintic and antileprosy properties. Conventionally, *M. dubia* is propagated through seeds, which have very poor (14%-34.3%) germination rates because of its hard stony seed coat, which makes it difficult to germinate without any treatment (Anand et al., 2012). The combined juice of papaya leaf, malaivembu or hill neem and common neem has been a daily dose for dengue patients for getting rid of the disease (Anand B et al., 2012).

Demand for *Meliadubia*

It is a fast growing multipurpose tree species, which has high demand in plywood industry, termite and fungal resistant timber and has potential to use in biomass power plants (power generation) (Batcher M.S). *Meliadubia* through conventional propagation using seeds has limited viability period. Fruits mature on tree, but do not drop and cling to the branches and loose viability on the tree itself. Farmers use wilding and raise bund planting of *M. dubia*. Therefore, production of planting material of *M. dubia* has become a problem (Apichart Kaosaard, et al., 1998).

Need for *Invitro* Propagation

Meliadubia is one of the fast growing tree and give good returns. It can be used in collection of biomass or plywood industries. After 3 years, 30-40 tons of biomass/acre can be harvested up to 10 years. Up to 5 years, ground nut, chilli, turmeric, blackgram as inter crops can be cultivated.

Meliadubia has nutritive value in leaves which when fed by sheep, goats possess more nutritive value. In addition, the extract from different parts of the *M. dubia* plant are known to have antiviral, antibacterial, antifungal, antidiabetic, antineoplastic, antihelmintic and antileprosy properties. For this reason *M. dubia* trees growing naturally have been indiscriminately logged and resulted in significant decline in its population. Therefore, it is imperative to use plant tissue culture method for large scale production of clonal planting material of the species from superior genotypes for quick rejuvenation (Baskaran P, Jayabalan N, 2008).

2. Materials and Methods

Preparation of Explants

The living parts of the plant are referred as explants which are responsible for the development of a whole plant. Healthy mature 3 year old *Meliadubia* plant was procured. The nodes and young leaves as explants were collected from the plants. The collected explants were brought to the production laboratory and washed thoroughly in running tap water for 10 min in order to eliminate the muddy/unwanted particles from the explants. Excision of nodal explants of about 1 – 1.5 cm was done with the help of secateurs whereas leaf discs were prepared (Bhimi Ram et al., 2012).

Explants Sterilization

The Nodal explants were soaked in antifungal and antibacterial solution containing carbendazim (0.1%) and streptomycin (0.1%) for 15 minutes. The sterilization is followed by the treatment of detergent, Polysorbate-20 for 20 minutes. The explants were washed with sterile water

Volume 5 Issue 6, June 2016

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

three times to ensure the complete wash of detergent. The sterilization is further carried out inside laminar air flow chamber; explants were treated with Ethanol (70%) for 30 sec and Mercuric chloride (0.1%) spanned for 8, 9, 10, 11, 12 min. The explants were removed from the mercuric chloride solution and washed with sterile water 3 times to eliminate the toxic effects of Mercuric chloride (Bhimi Ram et al).

The tender leaves as explants were soaked in antifungal and antibacterial solution containing carbendazim (0.1%) and streptocycline (0.1%) for 15 minutes. The sterilization is followed by the treatment of detergent, Polysorbate-20 for 15-20 minutes (Bonner J. and Devirian P.S, 1999). The explants were washed with sterile water three times to ensure the complete wash of detergent. The sterilization is further carried out inside laminar air flow chamber; explants were treated with Ethanol (70%) for 30 seconds and Mercuric chloride (0.1%) spanned for 3, 5, 7, 10min. The explants were removed from the mercuric chloride solution and washed with sterile water 3 times to eliminate the toxic effects of Mercuric chloride (Biswas K et al., 2002).

The Mortality rate was calculated by

$$\% \text{ Mortality} = \frac{\text{Explants contaminated} \times 100}{\text{Total no of Explants}}$$

3. Explants Initiation

Initiation stage begins with the emergence of shoots using nodes whereas occurrence of callus induction will be therefore leaves. In leaves, callus regeneration was observed in 60 days which is referred as Indirect organogenesis whereas in nodes, Direct organogenesis was observed which gave raise to shoot formation in 15 days.

The surface sterilized explants were inoculated in following MS basal media treatments + Sucrose 3% with various growth regulator concentrations (Bonner J. and Addicott F, 1997).

Nodes Leaves

Table 1: Initiation Media for the Nodes and Leaves

MEDIA	6BAP+NAA COMBINATION (mg/l)	MEDIA	6BAP TRIALS (mg/l)
N1	0.5+0.1	L1	0.1
N2	1+0.1	L2	0.2
N3	1.5+0.1	L3	0.3
N4	2+0.1	L4	0.4
N5	0.5+0.2	L5	0.5
N6	1+0.2	L6	0.6
N7	1.5+0.2	L7	0.7
N8	2+0.2	L8	0.8
		L9	0.9
		L10	1

The explants were placed in upright position on each treatment. The inoculated jars were incubated.

Culture Conditions


The Nodal explants were subjected to light for 10-12 h in the growth room after 2 days of darkness. Initiated leaves were placed under light intensity for 10-12 h in the growth room. Photoperiod provided by cool white fluorescent lamps of 1500-3000 lux, temperature of about $25 \pm 2^\circ\text{C}$ and humidity of 35 - 40%. The observation for the callus induction and shoot formation was observed often.

4. Results and Discussion


Nodal Explant Sterilization:

Table 2: Surface Sterilization with Nodes

Treatment	No Of Explants Taken	Non-Responsive Explants	Response (%)	Mean Rate Of Shoot Regeneration (%)
0.1% HgCl ₂ – 8 min	25	23	8	89
0.1% HgCl ₂ – 9 min		20	20	81
0.1% HgCl ₂ – 10 min		1	96	77
0.1% HgCl ₂ – 11 min		15	40	62
0.1% HgCl ₂ – 12 min		13	48	58



(i)



(ii)

Graph 1: (i) Graph showing Shoot generation response by nodal explants (ii) Graph showing Rate of Mortality with shoot regeneration


Majority of the explants survived after treatment when they were exposed to mercuric chloride for about 10 min. Treatment less than 10 min were found with more mortality of the explants whereas treatment for more than 10 min found to have mortality but comparatively less.

Rate of shoot generation was found as 77% whereas 89% of shoot regeneration was noted in 8 min treatment with mortality rate of 92%. At 10 min, the rate of mortality was found to be lesser and rate of shoot regeneration was moderate when compared to other treatments which showed almost equal mortality rate and shoot regeneration.


Leaves Sterilization

Table 3: Surface Sterilization with Leaves

Treatment	No Of Explants Taken	Non-Responsive Explants	Response (%)	Mean Rate Of Callus Induction (%)
0.1% HgCl ₂ – 3 min	25	24	4	95
0.1% HgCl ₂ – 5 min		2	92	86
0.1% HgCl ₂ – 7 min		19	24	80
0.1% HgCl ₂ – 9 min		16	36	69
0.1% HgCl ₂ – 10 min		15	40	62



(i)



(ii)

Graph 2: (i) Graph showing callus induction response by Leaves explants (ii) Graph showing Rate of Mortality with callus induction


Majority of the leaves explants survived after treatment when they were exposed to mercuric chloride for about 5

min. Other treatments were found with high mortality of the explants. Rate of callus induction was found as 86% when treated with 5 min, whereas 95% of callus induction was noted in 3 min treatment with mortality rate of 96%. At 5 min, the rate of mortality was found to be lesser and rate of callus induction was moderate when compared to other treatments which showed almost equal mortality rate and callus induction.


Initiation in Nodes

Table 4: Initiation Response in Nodes

Trial Media	No Of Survived Explants Taken	No Of Explants Raised Shoots	Mean Height Of Shoots (Cm)	% Initiation Response
N1	20	6.3	0.77	31.5
N2		11.6	1.2	58
N3		16.4	1.6	82.5
N4		18.5	2.3	93
N5		12	1.24	60
N6		10	1.08	50
N7		8.7	0.93	43.5
N8		5.3	0.56	26.5




(i)



(ii)

Graph 3: (i) Graph showing Initiation response in Nodes (ii) Graph showing the Mean height of the shoots in Nodes
 Figure 1: (a) Initiation of Nodes – Day 1 (b) Initiation of Nodes – Day 14 (c) Initiation of Nodes – Day 28





Initiation of nodal response was found as 93% which trailed with the combination of 6BAP+NAA = 2 mg/l + 0.1mg/l with mean height of 2.3 cm. other media trials with 6BAP and NAA of about 1, 3, 4, 5 mg/l and 0.2 mg/l were found with lesser shoot regeneration and less shoot height which was further transferred to multiplication stage.

Initiation in Leaves

Table 5: Initiation Response in Leaves

Trial Media	No Of Survived Explants Taken	No Of Explants Raised Shoots	Mean Height Of Shoots (cm)	Mean Initiation Response (%)
L1	20	6	0.4	30.3
L2		8	0.91	42
L3		9	1.12	56
L4		9	1.32	65.3
L5		11	1.67	72
L6		10	1.64	76
L7		13	1.76	79.8
L8		13	1.83	80
L9		14	1.89	85.5
L10		17	2.16	90.8



Graph 4: (i) Graph showing Mean Height of shoots from Callus (ii) Graph showing the Initiation response from Callus




Figure 2: (a) Initiation of Nodes – Day 14 (b) Initiation of Nodes – Day 28

Initiation of shoot response using leaves as explants was found as 90.8% which trailed with the combination of 6BAP= 1 mg/l with mean shoot height of 2.16 cm from the undifferentiated mass. Other media trials with 6BAP of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 mg/l were found with lesser callus induction and less shoot height which was further transferred to multiplication stage.

5. Discussion

In the present study, *Meliadubia* was selected as target plants which possess medicinal values as it is as equal as *Tectonagrandis* which can be used in plywood industry. The explants such as nodes and leaves which undergone various sterilization trials. Majority of the explants survived after treatment when they were exposed to mercuric chloride for about 10 min. Rate of shoot generation was found as 77% whereas 89% of shoot regeneration was noted in 8 min treatment with mortality rate of 92%. At 10 min, the rate of mortality was found to be lesser and rate of shoot regeneration was moderate when compared to other treatments which showed almost equal mortality rate and shoot regeneration. Majority of the leaves explants survived after treatment when they were exposed to mercuric chloride for about 5 min. Rate of callus induction was found as 86% when treated with 5 min, whereas 95% of callus induction was noted in 3 min treatment with mortality rate of 96%. At 5 min, the rate of mortality was found to be lesser and rate of callus induction was moderate when compared to other treatments which showed almost equal mortality rate and callus induction. Initiation of nodal response was found as 93% which trailed with the combination of 6BAP+NAA = 2 mg/l + 0.1mg/l with mean height of 2.3 cm. Response of callus induction in leaves was found as 90.8% which trailed with the combination of 6BAP + NAA= 1 mg/l + 0.1mg/l with mean shoot height of 2.16 cm from the undifferentiated mass.

6. Conclusion

Direct organogenesis was achieved for nodes and indirect organogenesis was done through leaves where undifferentiated mass called callus was formed and led to the formation of shoots. Mortality rate was lesser in Nodal explants whereas response was higher in leaves. Nodes performed best when compared to leaves which were developed by direct organogenesis.

Acknowledgement

We would like to thank Dr. V. PALANI, MD, Genewin Biotech for giving an opportunity to work in their lab and Mrs. MANASA SATHEESH, CTO, Genewin Biotech, Hosur for the technical guidance and continuous support throughout the work.

References

- [1] Anand B., Devagiri G.M., Maruti G., Vasudev H.S., Khaple A.K., "Effects of Pre-sowing Seed Treatments on Germination and Seedling Growth

- Performance of *Meliadubia* CAV: An Important Multipurpose Tree", International Journal of Life Sciences, 1, 59-63, 2012.
- [2] ApichartKaosaard, Verapong Suangtho and Erik D. Kjax, "Genetic Improvement Of Teak (*Tectona Grandis*) In Thailand", Forest Genetic Resources No. 26, 456-463, 1998.
- [3] Baskaran P, Jayabalan N, "Effect of growth regulators on rapid micropropagation and psoralen production in *Psoralea corylifolia*", Acta Physiological Plant Sciences, 30, 345-351, 2008.
- [4] Batcher M.S., "Element Stewardship Abstract for *Melia azedarach*", The Nature Conservancy, Arlington, USA, 7, 678-682.
- [5] Bhimi Ram, Trilok Singh Rathore, Gillela Ravi Shankar Reddy, "In vitro propagation of *Meliadubia* Cav. from seedling explants", Biotechnology Bioinformatics and Bioengineering, 2(1), 610-616, 2012.
- [6] Bhimi Ram, Trilok Singh Rathore and Beena Derajira Bopanna, "An efficient protocol for micropropagation and genetic stability analysis of *Meliadubia* Cav. - an important multipurpose tree", International Journal of Current Microbiology and Applied Sciences, 3(7), 533-544.
- [7] Biswas K., Chattopadhyay, Ishita, Banerjee R. K., Bandyopadhyay, Uday., "Biological activities and medicinal properties of neem (*Azadirachta indica*)", Current science, 82, 1336-1345, 2002.
- [8] Bonner J. and Addicott F., "Cultivation In Vitro Of Excised Pea Roots", Botany, 99, 144-170, 10, 1997.
- [9] Bonner J. and Devirian P.S., "Growth Factor Requirements Of Four Species Of Isolated Roots", American Journal of Botanical Sciences, 26, 661-665, 1999