

Third Molar Impaction and its Clinical Correlation- A Systematic Review

M. Ashfaq Ahmed¹, Dr. Thenmozhi²

¹I Year BDS, Saveetha Dental College and Hospitals, Saveetha University, P.H.Road, Chennai-600077, India

²Head of the Department, Department of Anatomy, Saveetha Dental College and Hospitals, Saveetha University, P.H.Road, Chennai-600077, India

Abstract: Aim: To determine the intensity of the impacted third molar and its clinical correlation. Objective: This study was done to check the impacted third molar and to study its clinical correlation. Background: A wisdom tooth or third molar is one of the three molars per quadrant of the human dentition. Wisdom teeth generally erupt between the ages of 17 and 25. Wisdom teeth commonly affect other teeth as they develop becoming impacted or "coming in sideways." They are often extracted when this occurs. Impacted wisdom teeth (or impacted third molars) are wisdom teeth which do not fully erupt into the mouth because of blockage from other teeth. Impacted wisdom teeth are classified by their direction of impaction, their depth compared to the biting surface of adjacent teeth and the amount of the tooth's crown that extends through gum tissue or bone. A lack of room to allow the teeth to erupt results in a risk of periodontal disease and dental cavities that increases with age. Only a small minority of adults age 65 years or older maintain the teeth without caries or periodontal disease and 13% maintain unimpacted wisdom teeth without caries or periodontal disease. Reason: This study may help people in understanding the significance of the third molar impaction.

Keywords: Impacted third molar, extraction, anaesthesia, bleeding, gauze, inferior alveolar nerve, lingual nerve, complications

1. Introduction

Third molar also known as wisdom teeth are the last teeth to erupt in the oral cavity. This generally occurs in the ages between 17 to 25, a time that has been called as the age of wisdom^[1].

Anthropologists say that the rough diet of early humans resulted in excessive wear of their teeth. Normal drifting of the teeth to compensate for this wear ensured that space was available for most of the wisdom teeth to erupt by adolescence^[2]. The modern diet which is much softer, and the orthodontic tooth straightening procedures produce a fuller dental arch^[3], which quite commonly doesn't allow room for the third molar to erupt thereby setting the stage for the problems such as impaction when the final four molars enter the mouth^[4,5].

Impacted tooth is defined as the tooth that is prevented from entering into the oral cavity due to malposition, lack of space or other impediments within the expected time^[6]. Impacted third molar teeth are classified by their direction of impaction, their depth compared to the biting surface of adjacent teeth and the amount of the tooth's crown that extends through gum tissue or bone^[7]. Impaction of teeth occurs in 73% of young adults^[8]. The average age of third molar eruption in males is 3 to 6 months ahead of females and it is found that females are more prone to third molar impaction than males^[9]. Third molar eruption and positional changes after eruption can be related with the nature of diet, race, genetic background and intense use of masticatory apparatus^[10]. A lack of room to allow the teeth to erupt results in a risk of periodontal disease and dental cavities that increases with age^[11]. Only a small minority of adults age 65 years or older maintain the teeth without caries or periodontal disease and 13% maintain unimpacted wisdom teeth without caries or periodontal disease^[12].

Impaction of third molar is a common condition related to the difficulty in degree of extraction and complications including iatrogenic trigeminal nerve injury^[13,14].

2. Clinical Anatomy of Third Molar

Third molar is situated at the distal end of the body of the mandible where it is in connection with the ramus of the mandible. In other ways It is found in anterior to the pterygomandibular raphe^[15]. There is a region of weakness and fracture can occur if excess force is applied during impaction of the wisdom tooth elevation. The roots of third molar are in close proximity to mandibular canal. They can penetrate into the mandibular canal or can be deflected. These canals with roots can evoke inferior alveolar nerve damage^[16]. The buccal alveolar bone is thicker than the lingual. The lingual nerve lies close to the third molar, therefore there is a high risk of lingual nerve damage while performing lingual split technique or elevating third molar flap medially to the distoangular recess.^[17]

3. Surgical Extraction of the Third Molar Tooth

Before the removal of the third molar there are certain conditions that are necessary to be checked such as position of the tooth and root development. Third molar removal may require a more involved surgical procedure^[18]. Most of the wisdom tooth are removed with the help of the oral and maxillofacial surgeon. The surgery initially begins with the administration of local anaesthesia, intravenous sedation or general anaesthesia^[19].

- **Local anaesthesia:** It is administered with one or more injections near the site of extraction.
- **Sedation anaesthesia:** It is given through an intravenous(IV) line in the arm. It suppresses the consciousness during the procedure.

Volume 5 Issue 7, July 2016

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

- **General anaesthesia:** Medication is inhaled through the nose or given an IV line in the arm. Then medication, breathing, temperature, fluids, and blood pressure are monitored^[20].

Procedure

- Incision is made in the gum tissue to expose the tooth and the bone^[21].
- Removal of the bone that blocks access to the root.
- Tooth is divided into sections.
- Tooth is removed.
- The site of the removed tooth is cleaned of any debris or bone.
- Wound closure by stitching the wound to promote healing.
- A gauze is placed over the extraction site to control bleeding^[22].

Post Operative Healing

Immediately following the removal of tooth, bleeding commonly occurs. Pressure is applied by biting on the gauze swab for at least 30 minutes and it should be left undisturbed. A thrombus forms in the socket^[23]. Antibiotics are prescribed to reduce the risk of certain post extraction complications^[24].

- **Smoking:** The patient should stop smoking as it reduces the ability for the blood to clot and also retards healing.
- **Pain:** some discomfort is normal after surgery. To minimize pain some medications such as nuprin, Tylenol, advil or some non aspirin pain reliever should be taken every 3 to 4 hours along with the diet.
- **Swelling:** Icebags should be applied over the operated area to minimize swelling. It should be applied every 15 minutes^[25].
- **Brushing:** It should not be done for the first 8 hours after surgery. After this it can be done gently.
- **Rinsing:** Rinsing should be avoided for 24 hours after extraction. Rinsing can disturb the formation of healing blood clot. This could cause rinsing and the risk of dry socket. After 24 hours it can be done with salt water solutions.
- **Diet:** soft foods are to be taken for first two days. Plenty of water must be taken^[26].

Instruments Used In Extraction

- **Lower molar forceps:** The lower third molar has bifurcated root with variable root pattern. Therefore the forceps will need a twin beak on either side.
- **Upper molar forceps:** upper molar has three roots (two buccal and one palatal). The forceps are twin beaked on the buccal and have a side blade palatally. Because of asymmetrical beaks, the instrument is not reversible and separate beaks are needed on right and left side.
- **Bayonet forceps:** It has blades offset substantially in distal direction. The shape resembles bayonet rifle attachment.
- **Cowhorns forceps:** It penetrates between the roots of molars to split the roots especially if the roots are diverged.
- **Elevator:** They differ from forceps in the manner in which the force is applied. The elevator applies forces

between the tooth and the bone. This is an instrument especially used to remove impacted teeth. They exert less directional force so tooth is less likely to be fractured.

- **Scalpel:** It is used to make incisions on the gums where a flap can be raised to gain access to tooth.
- **Chisels and mallet:** They are useful for distoangular third molars and upper third molar extraction.
- **Surgical burs:** Used as a straight hand piece, burs are used to remove bone surrounding the tooth to be apart.
- **Bone curette:** It is probably the most valuable instrument and are used to remove the granulation tissue from the tooth socket^[27].

4. Complications

- **Dry socket (alveolar osteitis):** is a painful phenomenon that mostly occurs after the removal of lower third molar teeth. It typically occurs when the blood clot with the extraction site is disrupted. The empty socket causes ache or throbbing pain in the gingival and can be intense^[28]. The pain is the only symptom seen which radiates up and down the head and neck. There are evidences that the use chlorohexidine can prevent the extent of dry socket but the risk factors increases with smoking after extraction^[29].
- **Nerve injury:** Injury to sections of nerve is another complication of wisdom tooth removal. This can cause pain, a tingling sensation and numbness in the tongue, lips, teeth and gums^[30]. A nerve injury can interfere with our daily activities, making things such as eating and drinking difficult and painful^[31]. There are two types of nerve injury:
 - 1) **Inferior alveolar nerve injury:** This nerve enters the mandible at the mandibular foramen and exits the mandible through the mental foramen^[32]. It supplies the sensation of the lower teeth. Injury is caused while lifting the tooth upwards during extraction. It also causes loss of sensation to all the teeth^[33].
 - 2) **Lingual nerve injury:** This nerve is a branch of mandibular nerve which in turn is a branch of trigeminal nerve. It gives off a sensory branch to the tongue by the chorda tympani nerve^[34]. Therefore damage to this nerve causes loss of pain and numbness in the anterior two-thirds of the tongue^[35]. It is commonly accepted to be 2% temporary and 0.2% permanent^[36].
- **Sinus:** The maxillary sinus lies above the roots of molars. The bony floor of the sinus divides the tooth socket from the sinus. Therefore extracting upper third molar can result in sinus exposure^[37].

5. Conclusion

The information provided above about the third molar impaction and its extraction and complication is promising to be a helpful tool for impacted tooth assessment as well as for planning surgical operation.

References

- [1] Agarwal KN, Gupta R, Faridi MM, Kalra N. Permanent dentition in Delhi boys of age 5-14 years. Indian Pediatr. 2004 Oct;41(10):1031-5.

- [2] Khan NB, Chohan AN, AlMoghrabi B, AlDeyab S, Zahid T, AlMoutairi M. Eruption Time of Permanent First Molars and Incisors Among a Sample of Saudi Male Schoolchildren. *Saudi Dent J.* 2006 Jan-Apr;18(1):18-24.
- [3] Elsey MJ, Rock WP. Influence of orthodontic treatment on development of third molars. *Br J Oral Maxillofac Surg.* 2000 Aug;38(4):350-3.
- [4] Odusanya SA, Abayomi IO. Third molar eruption among rural Nigerians. *Oral Surg Oral Med Oral Pathol.* 1991 Feb;71(2):151-4.
- [5] Dodson TB, Susarta SM (Apr 2010). "Impacted wisdom teeth (systematic review)". *Clin Evid (Online)* 2010(1302).
- [6] Peterson LJ. Principles of Management of Impacted Teeth. In: Peterson LJ, Ellis E III, Hupp JR, Tucker MR, editors. *Contemporary Oral and Maxillofacial Surgery*, 3rd ed. St. Louis: Mosby; 1998. p. 215-48.
- [7] Bishara SE, Andreasen G. Third molars: a review. *Am J Orthod.* 1983 Feb;83(2):131-7.
- [8] Richardson M. The development of third molar impaction. *Br J Orthod.* 1975 Oct;2(4):231-4.
- [9] Odusanya SA. Third molar impaction among Nigerian youths. *Odontostomatol Trop.* 1984 Jun;7(2):79-83.
- [10] Textbook of Oral and Maxillofacial Surgery. Kurger GO, editor. 6 th ed. Mosby: New Delhi: Jaypee Brothers; 1990.
- [11] Nanda, R. S., "Agenesis of the Third Molar in Man," *American Journal of Orthodontics*, Vol. 40, No. 9, 1954, pp. 698-706.
- [12] Akadiri OA, Obiechina AE. Assessment of difficulty in third molar surgery - a systematic review. *J Oral Maxillofac Surg* 2009 Apr;67(4):771-4.
- [13] Kay LW, Killey HC. The extraction of teeth. The impacted third molar. *Dent Update.* 1976 Mar-Apr;3(2):49-52.
- [14] Blondeau F, Daniel NG. Extraction of impacted mandibular third molars: postoperative complications and their risk factors. *J Can Dent Assoc.* 2007 May;73(4):325.
- [15] Xu GZ, Yang C, Fan XD, Yu CQ, Cai XY, Wang Y, He D. Anatomic relationship between impacted third mandibular molar and the mandibular canal as the risk factor of inferior alveolar nerve injury. *Br J Oral Maxillofac Surg.* 2013 Feb 11.
- [16] Westesson PL, Carlsson LE. Anatomy of mandibular third molars. A comparison between radiographic appearance and clinical observations. *Oral Surg Oral Med Oral Pathol* 1980;49:90-4.
- [17] Nelson ST, Ash MM. Wheeler's dental anatomy, physiology and occlusion. 9 th ed. St. Louis: Saunders Elsevier; 2010. p. 184-8.
- [18] Removal of third molars. Sponsored by the National Institute of Dental Research, November 28-30, 1979. *Natl Inst Health Consens Dev Conf Summ.* 1979;2:65-8s.
- [19] Davis, M.J., Vogel, L.D. Local anesthetic safety in pediatric patients. *N Y State Dent J.* 1996;62:32-35.
- [20] Archer WH. *Oral Surgery: A Step-By-Step Atlas of Operative Techniques*, 4th ed. Philadelphia: W.B. Saunders Company; 1966. p. 507-10.
- [21] Koerner KR. The removal of impacted third molars. Principles and procedures. *Dent Clin North Am.* 1994 Apr;38(2):255-78.
- [22] Garcia AG, Sampedro FG, Rey JG, Vila PG, Martin MS. Pell-Gregory classification is unreliable as a predictor of difficulty in extracting impacted lower third molars. *Br J Oral Maxillofac Surg.* 2000 Dec;38(6):585-587.
- [23] Muhonen A, Venta I, Ylipaavalniemi P. Factors predisposing to postoperative complications related to wisdom tooth surgery among university students. *J Am Coll Health* 1997;46:39 - 42.
- [24] Kay LW, Killey HC. The extraction of teeth. The impacted third molar. *Dent Update.* 1976 Mar-Apr;3(2):49-52.
- [25] Yuasa H, Sugiura M. Clinical postoperative findings after removal of impacted mandibular third molars: prediction of postoperative facial swelling and pain based on preoperative variables. *Br J Oral Maxillofac Surg.* 2004 Jun;42(3):209-14.
- [26] Susarla SM, Blaeser BF, Magalnick D. Third molar surgery and associated complications. *Oral Maxillofac Surg Clin North Am.* 2003 May;15(2):177-86.
- [27] Pell GJ, Gregory BT. Impacted mandibular third molars: classification and modified techniques for removal. *Dent Digest* 1933; 39: 330-338.
- [28] Lilly GE, Osborn DB, Rael EM, Samuel HS, Jones JC. Alveolar Osteitis associated with mandibular third molar extraction. *J Am Dent Assoc* 1974;88:802-6.
- [29] Bouloux GF, Steed MB, Perciaccante VJ. Complications of third molar surgery. *Oral Maxillofac Surg Clin North Am.* 2007 Feb;19(1):117-28, vii.
- [30] Carmichael FA, McGowan DA. Incidence of nerve damage following third molar removal: a West Scotland oral surgery research group study. *Br J Oral Surg* 1992-30:78-82.
- [31] Rood JP, Shehab BA. The radiological prediction of inferior alveolar nerve injury during third molar surgery. *Br J Oral Maxillofac Surg.* 1990 Feb;28(1):20-5.
- [32] Renton T, Hankins M, Sproate C, McGurk M. A randomised controlled clinical trial to compare the incidence of injury to the inferior alveolar nerve as a result of coronectomy and removal of mandibular third molars. *Br J Oral Maxillofac Surg.* 2005 Feb;43(1):7-12.
- [33] Howe GL, Poyton HG. Prevention of damage to the inferior dental nerve during the extraction of mandibular third molars. *Br Dent J* 1960;109:355-63.
- [34] Renton T, Yilmaz Z, Gaballah K. Evaluation of trigeminal nerve injuries in relation to third molar surgery in a prospective patient cohort. Recommendations for prevention. *Int J Oral Maxillofac Surg.* 2012 Dec;41(12):1509-18. Epub 2012 Sep 25.
- [35] Blackburn CW, Bramley PA. Lingual nerve damage associated with the removal of lower third molars. *Br Dent J.* 1989 Aug 5;167(3):103-7.
- [36] Hillerup S, Stoltze K. Lingual nerve injury in third molar surgery I. Observations on recovery of sensation with spontaneous healing. *Int J Oral Maxillofac Surg.* 2007 Oct;36(10):884-9. Epub 2007 Sep 4.

[37] Marciani RD. Complications of third molar surgery and their management. Atlas Oral Maxillofac Surg Clin North Am. 2012 Sep;20(2):233-51.

