3-Equitable Labeling in Context of the Barycentric Subdivision of Cycle Related Graph

I. I. Jadav¹, G. V. Ghodasara²

¹Research Scholar, R K University, Rajkot-360020, Gujarat-India

²H & H. B. Kotak Institute of Science, Rajkot-360001, Gujarat-India

Abstract: A function from the vertex set of a graph G to the set $\{0, 1, 2\}$ is called 3-equitable labeling if the induced edge labels are produced by the absolute difference of labels of end vertices such that the absolute difference of number of vertices of G labeled with 0,1 and 2 differby at most 1 and similarly the absolute difference of number of edges of G labeled with 0,1 and 2 differby at most 1[1]. In this paper, we discuss 3-equitable labeling in the context of barycentric subdivision of different cycle related graphs.

Keywords: 3-equitable labeling, barycentric subdivision

AMS Subject classification number (2010): 05C78

1. Introduction

We consider simple, finite, undirected graph. If the vertices of the graph are assigned values subject to certain conditions, then it is known as a graph labeling [5]. A survey on graph labeling is given by Gallian[3].

We follow Gross and Yellen [5] for the graph theoretical terminology and notations. The vertex set and the edge set of graph G are denoted by V(G) and E(G) respectively. A mapping f from V(G) to $\{0,1,2\}$ is called ternary vertex labeling of G[1]. Aternary vertex labeling of a graph G iscalled 3-equitable labeling if the induced edge labeling function f^* from E(G) to the set $\{0,1,2\}$ is defined as $f^*(e = uv) = |f(u) - f(v)|$ such that the absolute difference of number of vertices of G with label0,1 and 2 differ by at most 1. A graph which admits 3-equitable labeling is called a 3-equitable graph [1].

$C_n \odot K_1$	Crown obtained from the corona of C_n with K_1 .	
$C_n \odot 2K_1$	Double crown obtained from the corona	
	of C_n with $2K_1$.	
AC _n	Armed crown obtained from $C_m \oplus P_n$.	
C'	Obtained by duplicating a vertex by an edge in a	
<i>C</i> _{<i>n</i>}	cycle C_n , where $\{v'_1, v'_2,, v'_{2n}\}$ are vertices	
	added to obtain $C^{'}_n$ corresponding to the vertices	
	$v_1, v_2,, v_n$ in C_n .	

Notation 1.1: Standard graphs used in this paper [2]: Let e = uv bean edge of a graph G and w is not a vertex of G. Then edge is said to be subdivided when it is replaced by edges e' = uw and e'' = vw[5]. If every edge of graph G is subdivided, then the resulting graph is called barycentric subdivision of the graph G. It is denoted by S(G). Vaidya et al.[6] proved that cycle with twin chords is cordial as well as 3-equitable. In [7] Vaidyaetal. proved that the barycentric subdivision of cycle with one chord, cycle with twin chords and cycle with triangle are cordial. G. V. Ghodasra and I.I. Jadav[4] proved that barycentric subdivision of cycle with twin chords, cycle with triangle, shell graph and wheel graph are3-equitable. In this paper, we prove that the barycentric subdivision of crown, double crown, armed crown and some standard graph are3-equitable graphs.

2. Main Results

Theorem 1: The barycentric subdivision of crown $C_n \odot K_1$ is 3-equitable. **Proof:** Let $S(C_n \odot K_1)$ denote the barycentric subdivision of crown $C_n \odot K_1$. Let $u_1, u_2, ..., u_n$ be pendent vertices of $C_n \odot K_1$ and $v_1, v_2, ..., v_{2n}$ are the vertices corresponding to C_n , where v_j is the vertex added due to the barycentric subdivision of edge $V_{j-1}V_{j+1}$, j = 2,4,6,...,2n-2 and v_{2n} is the vertex added due to the barycentric subdivision of edge $v_{2n-1}v_1$. Here, w_i is the vertex added due to the barycentric subdivision of edge $u_iv_{2i-1}, (i = 1,2,3,...,n)$. Note that $|V(S(C_n \odot K_1))| = 4n$ and $|E(S(C_n \odot K_1))| = 4n$. We define labeling function $f: V(S(C_n \odot K_1)) \rightarrow \{0,1,2\}$ as follows

$$f(v_i) = \begin{cases} 0; i \equiv 0, 3 \pmod{6} \\ 1; i \equiv 4, 5 \pmod{6} \\ 2; i \equiv 1, 2 \pmod{6}, 1 \le i \le 2n. \end{cases}$$

$$f(u_i) = \begin{cases} 1; i \equiv 1, 4 \pmod{6} \\ 2; i \equiv 0, 2, 3, 5 \pmod{6}, 1 \le i \le n. \end{cases}$$

$$f(w_i) = \begin{cases} 0; i \equiv 1, 2, 4, 5 \pmod{6} \\ 2; i \equiv 0, 3 \pmod{6}, 1 \le i \le n. \end{cases}$$

Hence $S(C_n \odot K_1)$ is 3-equitable, for all $n \in \mathbb{N}$, $n \ge 3$. Above defined labeling pattern satisfies the conditions of 3-equitable labeling as shown in Table 1.

Let
$$n = 6a + b$$
, where $b \in \mathbb{N}$ and $a \in \mathbb{N} \cup \{0\}$.

TADIC 1. TADIC IDI THEOREM I	Table 1:	Table for	Theorem	1
-------------------------------------	----------	-----------	---------	---

b	Vertex Conditions	Edge Conditions
0,3	$v_f(0) = v_f(1) = v_f(2)$	$e_f(0) = e_f(1) = e_f(2)$
1,4	$v_f(0) + 1 = v_f(1) + 1 = v_f(2)$	$e_f(0) = e_f(1) + 1 = e_f(2) + 1$
2,5	$v_f(0) = v_f(1) + 1 = v_f(2)$	$e_f(0) + 1 = e_f(1) = e_f(2)$

Example 1.3-equitable labeling of the graph obtained by barycentric subdivision of crown $C_5 \odot K_1$ is shown in Figure 1. It is the case related to $n \equiv 5 \pmod{6}$.



Theorem 2. The barycentric subdivision of double crown $C_n \odot 2K_1$ is 3-equitable.

Proof. Let $S(C_n \odot 2K_1)$ denotes the barycentric subdivision of double crown $C_n \odot 2K_1$. Let $u_1, u_2, ..., u_{2n}$ be the pendant vertices of $C_n \odot 2K_1$ and $v_1, v_2, ..., v_{2n}$ are vertices corresponding to C_n , where v_i is the vertex added due to barycentric subdivision of edge $v_{j-1}v_{j+1}$ j = 2, 4, 6, ..., 2n - 2 and v_{2n} is the vertex added due to the barycentric subdivision of edge $v_{2n-1}v_1$. Here, W_i is the vertex added due to the barycentric subdivision of edge for odd \mathbf{i} and $u_i v_{i-1}$ for even $u_i v_i$ i (i = 2, 4, 6, ..., 2n). Note that $|V(S(C_n \odot 2K_1))| = 6n$ and

 $|E(S(C_n \odot 2K_1))| = 6n$. We define labeling function $f: V(S(C_n \odot 2K_1)) \rightarrow \{0,1,2\}$ as follows.

$$\begin{split} f(u_i) &= 1; if \ 1 \leq i \leq 2n. \\ f(v_i) &= 0; if \ 1 \leq i \leq 2n. \\ f(w_i) &= 2; if \ 1 \leq i \leq 2n. \end{split}$$

From the above labeling pattern, we get $v_f(0) = v_f(1) = v_f(2)$ and $e_f(0) = e_f(1) = e_f(2)$.

Hence $S(C_n \odot K_1)$ is 3-equitable graph.

Example 2.3-equitable labeling of the graph obtained by barycentric subdivision of double crown $C_4 \odot 2K_1$ is shown in Figure 2. It is the case related to $n \equiv 4 \pmod{6}$.



Theorem 3: The barycentric subdivision of armed crown AC_n is 3-equitable.

Proof: Let $S(AC_n)$ denote the barycentric subdivision of armed crown AC_n .Let u_i denote the pendant vertices of AC_n and $v_1, v_2, ..., v_{2n}$ are the vertices corresponding to C_n , where v_j is the vertex added due to the barycentric subdivision of edge $v_{j-1}v_{j+1}$, j = 2,4,6,...,2n-2 and v_{2n} is the vertex added due to the barycentric subdivision of edge $v_{2n-1}v_1$. Here w_i is the vertices of degree wo and w'_i is the vertex added due to barycentric subdivision of edge

Volume 5 Issue 7, July 2016

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

 $w_i v_{2i-1}, u'_i \text{ is the vertex added due to barycentric subdivision}$ of edge $w_i u_i, 1 \le i \le n$. Note that $|V(S(AC_n))| = 6n$ and $|E(S(AC_n))| = 6n$. We define labeling function $f: V(S(AC_n)) = 6n \rightarrow \{0,1,2\}$ as follows. Case 1: n is even $f(v_i) = \begin{cases} 2; i \equiv 0,1 \pmod{4} \\ 0; i \equiv 2,3, \pmod{4}, 1 \le i \le 2n. \end{cases}$ $f(w_i) = 1; if \ 1 \le i \le n.$ $f(w'_i) = 1; if \ 1 \le i \le n.$ $f(u_i) = 0; if \ 1 \le i \le n.$ $f(u'_i) = 2; if \ 1 \le i \le n.$

Case 2: n is odd.

$$f(v_i) = \begin{cases} 2; i \equiv 0, 1 \pmod{4} \\ 0; i \equiv 2, 3, (\mod{4}), 1 \le i \le 2n. \end{cases}$$

$$f(w_n) = 1, f(w'_n) = 2, f(u_n) = 0, f(u'_n) = 1$$

$$f(w_i) = 1; if \ 1 \le i \le n - 1.$$

$$f(w'_i) = 1; if \ 1 \le i \le n - 1.$$

$$f(u'_i) = 0; if \ 1 \le i \le n - 1.$$

$$f(u'_i) = 2; if \ 1 \le i \le n - 1.$$

From the above labeling pattern, we get $v_f(0) = v_f(1) = v_f(2)$ and $e_f(0) = e_f(1) = e_f(2).$ Hence $S(AC_n)$ is 3-equitable graph.

Example 3. 3-equitable labeling of the graph obtained by barycentric subdivision of armed crown**AC**₃ is shown in

Figure 3. It is the case related ton $\equiv 3 \pmod{6}$.



Theorem 4. The barycentric subdivision of C'_n is 3-equitable.

Proof. Let denotes the barycentric subdivision of C'_n . Let $u_1, u_2, ..., u_{2n}$ be the outer vertices of C'_n and $v_1, v_2, ..., v_{2n}$ are the vertices corresponding to C_n , where v_j is the vertex added due to the barycentric subdivision of edge $v_{j-1}v_{j+1}$, j = 2,4,6,...,2n-2 and v_{2n} is the vertex added due to the barycentric subdivision of edge $u_{2n-1}v_1$ and w_i is the vertex added due to barycentric subdivision of edge $u_{2n-1}u_{2i}$, i = 1,2,3,...,n. Here u'_i is the vertex added due to barycentric subdivision of edge u_iv_i for odd i and edge u_iv_{i-1} for even i, (i = 1,2,3,...,2n).

Note that $|V(S(C'_n))| = 8n$ and $|E(S(C'_n))| = 8n$

We define labeling function

$$f:V(S(C'_n)) = 6n \rightarrow \{0,1,2\}_{as \text{ follows}}$$
Case 1: $n \equiv 0,3 \pmod{6}$
 $f(v_i) = 1; if \ 1 \le i \le 2n.$
 $f(u_i) = 2; if \ 1 \le i \le 2n.$
 $f(u'_i) = 0; if \ 1 \le i \le 2n.$
(0: $i = 1 \pmod{6}$

$$f(w_i) = \begin{cases} 0, i \in 1, 4 \pmod{6} \\ 1; i \in 2, 5 \pmod{6} \\ 2; i \in 0, 3 \pmod{6}, 1 \le i \le n. \end{cases}$$

Case 2:
$$n \equiv 1,2,4,5 \pmod{6}$$

 $f(u'_{2n}) = 2.$
 $f(v_i) = 1; if \ 1 \le i \le 2n$
 $f(u_i) = 2; if \ 1 \le i \le 2n.$
 $f(u'_i) = 0; if \ 1 \le i \le 2n - 1$
 $f(w_i) = \begin{cases} 0; i \equiv 1,4 \pmod{6} \\ 1; i \equiv 2,5 \pmod{6} \\ 2; i \equiv 0,3 \pmod{6}, 1 \le i \le n. \end{cases}$

Above defined labeling pattern satisfies the conditions of 3equitable labeling as shown in Table4.

Let n = 6a + b, where $b \in \mathbb{N}$ and $a \in \mathbb{N} \cup \{0\}$.

Figure 3

Table 2: Table for Theorem 4

b	Vertex Conditions	Edge Conditions
0,3	$v_f(0) = v_f(1) = v_f(2)$	$e_f(0) = e_f(1) = e_f(2)$
2,5	$v_f(0) + 1 = v_f(1) = v_f(2)$	$e_f(0) + 1 = e_f(1) = e_f(2) + 1$
1,4	$v_f(0) + 1 = v_f(1) + 1 = v_f(2)$	$e_f(0) = e_f(1) + 1 = e_f(2)$

Volume 5 Issue 7, July 2016 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY Hence the barycentric subdivision of C'_n is 3-equitable.

Example 3.3: equitable labeling of the graph obtained by barycentric subdivision of C'_n is shown in Figure 4.



References

- I.Cahit, Oncordial and 3-equitable labeling of graphs, Util. Math., 37(1990),189–198.
- [2] R.Distel, Graph theory, Springer verlagheidelberg, (2005).
- [3] J.A.Gallian, Adynamic survey of graph labeling, The
- Electronics Journal of Combinatoric 16(2016),#DS6 1–389.
- [4] G. V.Ghodasra,I.I.Jadav, 3-equitable labelingin context of the barycentric subdivision of some special graphs, International Journal of Mathematics and Soft Computing, Vol5, No.1(2015).
- [5] J. Gross and J.Yellen, Graph theory and its applications, C RCPress, (1999).
- [6] S.K.Vaidya, S. Srivastav, V.J. Kaneria and G. V. Ghodasara, Cordial and 3-equitable of cycle with twin chords, Proceedings of the First International Conference on Emerging Technologies and Applications in Engineering, Technology and Sciences, 1(2008), 905-907.
- [7] S.K.Vaidya, K.K.Kanani, S.Srivastav, and G. V. Ghodasara, Barycentric subdivision and cordial labeling of some cycle related graphs, Proceedings of the First International Conference on Emerging Technologies and Applications in Engineering, Technology and Sciences, 1(2008), 1081-1083