

Synthesis and Study of Manganese Complexes of Benzilic Acid

Dr. M. Jerome Rozario¹, Dr. A. John Merina²

¹Associate Professor, Department of Chemistry, PRIST University, Thanjavur


²Principal, Govt., Arts College for Women (Autonomous), Kumbakonam

Abstract: Benzilic acid complexes with Mn(II) metal ions have been synthesised and characterised by electrical conductance, analysis of cation, magnetic susceptibility, electronic and infrared spectral studies. The analytical data show that the Chloride complexes of Mn(II) have the general formula $[ML_4Cl_2]$. The Perchlorate complexes of Mn(II) have the general formula $[ML_6][ClO_4]$. The conductance measurements show that the chloride complexes behave as a 1:2 nonelectrolyte. The magnetic moment measurement indicates paramagnetic behavior for the complex and the electronic spectral data suggest octahedral geometry for the complexes. The infrared spectral studies clearly indicate that the carbonyl group is coordinated to the metal ion and benzilic acid acts as monodentate.

Keywords: benzilic acid, Manganese complexes, Coordination Chemistry, Inorganic Chemistry

1. Introduction

Benzilic acid is soluble in hot water and in ordinary organic solvents. The structure of benzilic acid is



Since carbonyl group serves as a donor of electrons, Benzilic acid acts as a ligand. Hence it is of immense importance to study the coordinating ability of this ligand with the metal Manganese. The transition metal Manganese is chosen to study the coordination tendency. This work deals with the Synthesis and characterization of Manganese complexes of benzilic acid.

2. Experimental Techniques

1. Preparation of Manganese(II) Chloride complex¹

The chloride of Mn(II) was prepared by the action of HCl on the metal carbonate. Manganese(II) chloride and benzilic acid were mixed in the molar ratio 1:6 in the ethanol and refluxed for five hours. The solution was concentrated to half of the volume and cooled to crystallize out the complex which were filtered at the pump. It was washed with ether to remove excess ligand and dried in a desiccator over anhydrous calcium chloride.

2. Preparation of Manganese(II) Perchlorate complex

About 10gm of manganese carbonate was made into paste by adding water 14ml of Perchloric acid was then added and the mixture was concentrated on a water bath to obtain the crystals

of manganese Perchlorate. It was treated again with water and concentrated. The crystals were dried over anhydrous calcium chloride in a desiccator.

Analysis of the complexes

Estimation of Metals

Preparation of the complex solution²

About 0.5gm of the complex was weighed accurately and 10ml of concentrated nitric acid was added. The solution was evaporated and concentrated almost to dryness. Then few ml of nitric acid was added and concentrated to 1ml. Then the solution was made up to 100ml in Standard flask.

Estimation of Manganese

The percentage of the Manganese was estimated volumetrically using Erio-T as an indicator. 20 ml of the made up solution was pipetted out into a clean conical flask. Excess water ammonia solution and one Spatula of Erio-T indicator were added. Then it was titrated against 0.01M EDTA solution. The end point was the disappearance of red to blue colour. From this the percentage of manganese was calculated.

General properties of the complexes^{3,4}

All complexes were coloured solids, soluble in ethanol, stable at room temperature and nonhygroscopic in nature.

3. Instrumental Methods

1. Electrical Conductivity

Conductance measurements of the complexes in solution were made to verify the ionic formulation of complexes. Conductivity bridge with dip-type platinized platinum electrodes were employed for measurements. The cell constant was determined by measuring the resistance of 0.1N KCl whose specific conductivities were known at various

temperature from literature. The solution used for conductivity measurements has concentration around 10-3M. All the measurements were corrected for the conductance of the solvent subtracting the conductance of the pure solvent from that of the solution.

Molar conductance calculated by using the relation.
 Specific conductance (K) = Conductance × Cell constant

$$\text{Molar conductance} = \frac{1000 \times \text{Specific conductance}}{\text{Molar concentration}}$$

$$\Delta m = 1000K/C$$

The values in Acetonitrile are given in Table II.

2. Magnetic Susceptibility Measurements

Magnetic susceptibility and hence magnetic moments of various complexes can be determined using the Gouy balance. The magnetic susceptibility of the complexes were calculated using the relation

$$106 X = \frac{\alpha + \beta F}{W}$$

Where F' = corrected force = F in mg, β = calibration constant, α = 0.029 × 10⁻⁶ × volume of the tube, W = Weight of substance in gm, X = gram susceptibility.

To find out the effective magnetic moment the following relation was used.

$$\mu_{\text{eff}} = 2.84 \sqrt{(X' M T)} B.M$$

X'M = Molar susceptibility, T = Absolute temperature.

The weights of the gouy tube were taken in the absence and in the presence of the magnetic field. To evaluate the value of calibrant constant the tube filled with Hg [Co(CNS)₄] up to the mark and the weight was taken in the absence or in the presence of the magnetic field. The calibration was removed and the gouy tube was filled with various complexes and the weight were taken. Finally the weight of the gouy tube with water, filled upto the mark was noted in order to find out the volume of the sample, using the formula, magnetic susceptibility and effective magnetic moment were calculated after making proper diamagnetic corrections. Using Pascal's constant⁵. The magnetic susceptibility values are given in Table III.

3. Electronic Spectra

The electronic spectra of the complexes were taken in the ethanol medium for various wavelengths ranging from 300–700nm. Absorbance values were plotted against the corresponding wavelengths (nm). The maximum values corresponding to the peaks were noted.

4. Infrared Spectra

The infrared spectra of the free ligand and the complexes were taken in KBr pellets in the range of 4000-200cm⁻¹ by infrared spectrometer (pellet technique). The percentage transmission was recorded against wavelength(cm⁻¹). The infrared spectral bands along with their probable assignments are given below.


4. Results and Discussion

1. Behavior of Complexes in Solution⁶

The electrical conductance of these complexes in ethanol were measured. Molar conductance values (Table II), expected for 1:0, 1:1, 1:1, 1:2, 1:3, 1:4 in Electrolyte in Acetonitrile are given below.

Conductance values (Ohm ⁻¹ cm ² mole ⁻¹)	Type of electrolyte
<80	1:0
80-115	1:1
160 -220	1:2
290-350	1:3
>450	1:4

The molar conductance of the Perchlorate complexes of the divalent metals correspond to those of 1:2 electrolytes indicating that the Perchlorate groups are ionic in nature⁷. The low electrical conductance values of chloride complexes show that all of them are 1:0 nonelectrolytes. Based on the analytical and the molar conductance data of the complexes, the prepared complexes may be assigned the following compositions.



2. Magnetic Susceptibility⁸

The strong and weak field complexes of several transition metal ions differ in the number of unpaired electrons in the complex. This number can be found readily by a comparison between the measured magnetic moment and that calculated from the spin only moment. Hence it is possible to distinguish spin paired and spin free complexes. Determination of the number of unpaired electron can also give information regarding the oxidation state of a metal ion in a complex. It is also useful in establishing the structure of many complexes⁹. The magnetic moments, μ_{eff} of the complexes are given in Table III. The values indicate that they are all of high spin type. The magnetic moment of Mn(II) chloride and Perchlorate complex are 5.92B.M and 5.96B.M respectively indicating octahedral geometry.


3. InfraRed Spectra¹⁰

The IR spectra of the complexes are very useful in determining the coordinating groups of the ligand. The shifts in the bands indicate mode of linkage in the complexes. The comparison of IR spectrum of Benzilic acid and its complex confirmed the formation of their complex. There were expected shifts in the IR spectrum of the complexes especially in the C=O stretching frequencies of the acid group of the benzilic acid. The C=O stretching frequency of the ligand are shifted to the lower frequency in the spectrum of the complex. The peak at 1720cm⁻¹ which is due to C=O stretching of the carboxylic acid group in the spectrum of ligand is shifted to 1690cm⁻¹ in the spectrum of the complex, the peak at 1230cm⁻¹ which may be to C=O or OH deformation of the carboxylic acid group in the ligand spectrum shifted to 1170cm⁻¹ in the

complex spectrum. The absence of symmetric and asymmetric C=O stretching frequencies and the presence of anions indicate that the acid group of benzoic acid is not ionized and the carbonyl oxygen is coordinated to metal ion. Vandorne and Hannik¹¹ reported that the unionized acetic acid is coordinated to metal atom through carbonyl oxygen. The band at 3380cm⁻¹ which is due to OH stretching of alcoholic group is not affected in the complex spectrum. Hence it may be concluded that the hydroxyl group is not coordinated with the metal ion and the carbonyl oxygen is coordinated with the metal ion.

4. Electronic spectra¹²

In the electronic spectra of high spin octahedral complexes of Mn(II) one should expect three bands corresponding to the transitions.



The Manganese(II) perchlorate complexes of benzoic acid exhibited the band at 485nm suggesting octahedral geometry.

5. Structure of the complexes

The analytical data suggest that benzoic acid is a monodentate ligand in which C=O group is coordinated to the metal

atom. The analytical and magnetic moment data suggested octahedral geometry for both Manganese(II) chloride and Perchlorate complexes (Table I).

Table 1: Analytical data and colour of benzoic acid complexes


No	Complex	Colour of the Complex	Percentage of Metal	
			Calculated	Observed
01	Mn(BENZ) ₄ Cl ₂	Pink	5.21	5.25
02	[Mn(BENZ) ₆](ClO ₄) ₂	Pale Pink	3.68	3.27

Table 2: Molar conductance in acetonitrile

No	Complex	Molar conductance	Nature of the Electrolyte
01	Mn(BENZ) ₄ Cl ₂	61.7	1:0
02	[Mn(BENZ) ₆](ClO ₄) ₂	210.7	1:2

Table 3: Magnetic moments of complexes

No	Complex	μ _{eff}	Magnetic Behavior
01	Mn(BENZ) ₄ Cl ₂	5.92	Para
02	[Mn(BENZ) ₆](ClO ₄) ₂	5.96	Para



5. Conclusion

Benzoic acid complexes with Mn(II) metal ions have been prepared and characterized by electrical conductance, analysis of cat ion, magnetic susceptibility, electronic and infrared spectral studies. The analytical data show that the CO have the general formula [ML₄Cl₂]. The perchlorate complex have the general formula [ML₆](ClO₄)₂. The conductance measurements show that the chloride and perchlorate complexes behave as a 1:2 nonelectrolyte and 1:2 electrolyte respectively. The magnetic moment measurement indicates paramagnetic behavior for the complexes and the octahedral geometry. The electronic spectral data suggest octahedral geometry for the complexes. The infrared spectral studies clearly indicate that the carbonyl group is coordinated to the metal ion and benzoic acid acts as monodentate ligand.

References

- [1] Eggleston, D.S. and Jackels, S.C., "Tetrasubstituted [14]-1,3,8,10-Tetraene N₄ Macrocyclic Complexes: Synthesis, Organic Precursor, and Template Reaction Mechanism". *Inorg.Chem.*, vol.19, pp.1593-1599,1980.
- [2] Malik, W.U, Bembi, R and Singh, R.(1983). Preparation and Characterization of Copper, Cobalt and Manganese Complexes of Tetradentate N₆ Macrocyclic Ligand. *Polyhedron*, 2, 369-373.
- [3] Eggleston, D.S and Jackels, S.C.(1980). Tetrasubstituted [14]-1,3,8,10-Tetraene N₄ Macrocyclic Complexes: Synthesis, Organic Precursor, and Template Reaction Mechanism. *Inorg.Chem.*, 19, 1593-1599.
- [4] Hathaway, B.J., and Billing, D.E., *Coord. Chem.Rev.*, vol.5, pp.143,1970.
- [5] Hathaway, B.J., Duggan, M., Murphy, A., Mullane, J., Power, C., Walsh, A., and Walsh, B., *Coord.Chem. Rev.*, vol.36, pp.267,1981.

- [6] Figgis, B.N and Lewis, J., "Mordrencoordination chemistry"(Interscience, 1960).
- [7] Parekh, J., Inamdhar, P., Nair, R., Baluja, S., and Chanda, S., J. Serb. Chem.Soc., vol.70, pp.1161, 2005.
- [8] Geary, W.J Coord.chem.Rewiew, 7(1971), 81.
- [9] Bersuker, I. B., Coord.Chem.Rev., vol.14, pp.367, 1975.
- [10] Nyholm, R.S.J.CoordChem.Rewiew, 1958, 8, 401.
- [11] Vandoorne, W and Hanni K.M. signth, React inorganic met.org.chemistry 11 (I), 1981, 15.
- [12] Sutton, D. "Electronic spectra of transition metal complexes" McGrawHill, London, 1968, 186-192.

