

# Effectiveness of Buerger Allen Exercise on Lower Extremity Perfusion and Pain among Patients with Type 2 Diabetes Mellitus in Selected Hospitals in Chennai

MS.E.Selmar Mellisha, M.SC (N)

Nursing Tutor, Department of Medical Surgical Nursing, M.A. Chidambaram College Of Nursing, Chennai, India

**Abstract:** A quasi experimental study was conducted to assess the effectiveness of Buerger Allen exercise on lower extremity perfusion and pain among patients with type 2 Diabetes Mellitus admitted in selected hospitals, Chennai. It was conducted among 60 samples with type 2 diabetes mellitus by using non probability purposive sampling technique. Pre test was conducted to assess the level of pain and lower extremity perfusion and Buerger Allen exercise was administered for the experimental group for a period of five days and post test was conducted on the sixth day. Comparison of pre and post test findings showed that in the experimental group, the mean score of level of lower extremity pain was reduced from 4.33 to 1.30. The reduction of pain was statistically significant difference at 1% level of significance ( $p=0.001$ ). The mean score of level of lower extremity perfusion was increased from 44.50 to 52 and it showed a statistically significant difference at 1% level of significance ( $p=0.001$ ).

**Keywords:** Buerger Allen exercise; type 2 diabetes mellitus; lower extremity perfusion and pain

## 1. Introduction

Diabetes Mellitus is a metabolic disorder that is characterized by increased levels of glucose in the blood resulting from defects in insulin secretion, insulin action, or both. Diabetes Mellitus causes, between two and four times increased risk of peripheral vascular disease by causing endothelial and smooth muscle cell dysfunction in peripheral arteries. The risk of developing lower extremity peripheral vascular disease is proportional to the severity and duration of diabetes. Most of the complications are preventable. It requires a lifelong commitment to staying healthy, maintaining weight, exercising, taking medications, as prescribed by doctor. Exercise is the fundamental principle for preventing the Peripheral vascular diseases among diabetes patients. One of the exercises is Buerger Allen exercise, is an active postural exercise of the feet and legs for preventing peripheral vascular disease and promoting collateral circulation in lower extremities.

## 2. Statement of the Problem

A study to assess the effectiveness of Buerger Allen exercise on lower extremity perfusion and pain among patients with type 2 Diabetes Mellitus admitted in selected hospitals, Chennai.

### Objectives

- To assess the pre and post test level of lower extremity perfusion and pain among patients with type 2 Diabetes Mellitus in experimental and control group.
- To assess the effectiveness of Buerger Allen exercise on lower extremity perfusion and pain among patients with type 2 Diabetes Mellitus.
- To associate the post test level of lower extremity perfusion and pain with selected demographic variables and clinical variables among patients with type 2

diabetes mellitus in both experimental and control group.

## 3. Methodology

The research approach was experimental in nature and quasi experimental pre test and post test control group design was used. The study was conducted among 60 samples with type 2 diabetes mellitus. Thirty samples in experimental group and 30 samples in control group were selected by using non probability purposive sampling technique. Pre test was conducted using structured interview schedule, modified Wong Baker FACES pain assessment scale, modified Clarke enhanced foot assessment scale. For the experimental group, the investigator demonstrated the Buerger Allen exercise and were asked to do the exercise three times a day at four hours interval (8am, 12noon, and 4pm) for a period of five days under the supervision of investigator. Post assessment was done on the sixth day by using the same scale.

## 4. Results



Comparison of pre and post test findings showed that in the experimental group, the mean score of level of lower extremity pain was reduced from 4.33 to 1.30. The reduction of pain was statistically significant difference at 1% level of significance ( $p=0.001$ ). The mean score of level of lower extremity perfusion was increased from 44.50 to 52 and it showed a statistically significant difference at 1% level of significance ( $p=0.001$ ). There was a statistically significant association found between post test level of lower extremity pain with demographic variables such as age, dietary pattern. Regarding lower extremity perfusion, the post test result showed a statistically significant association with age and specific habits such as smoking and alcoholism.

Volume 5 Issue 7, July 2016

[www.ijsr.net](http://www.ijsr.net)

Licensed Under Creative Commons Attribution CC BY

Assessment of Pre and Post Test Level of Lower Extremity Perfusion and Pain for Experimental and Control Group



Comparison of Level of Lower Extremity Perfusion and Pain among Experimental and Control Group  
 N=60 O<sub>1</sub>=30, O<sub>2</sub>=30

Variables	Test	N	Experimental group		Control group		Mean difference	Student independent t test
			Mean score	SD	Mean score	SD		
Lower extremity pain	Pre test	30	4.33	1.88	4.00	1.46	0.33	t=0.76 P=0.44
	Post test	30	1.30	1.34	3.83	1.64	2.53	t=6.54 P=0.001***
Lower extremity perfusion	Pre test	30	44.50	4.61	44.23	4.96	0.27	t=1.89 P=0.06
	Post test	30	52.00	3.31	44.57	4.84	7.43	t=19.03 P=0.001***

O<sub>1</sub>=Experimental group O<sub>2</sub>= Control group \*\*\* denotes significant at P≤0.001

Association of Post Test Level of Lower Extremity Pain With Selected Demographic and Clinical Variables of Samples in Experimental Group and in Control Group, N=30

Demographic variables	Post test level of pain						Total	Chi square
	No pain		Mild pain		Moderate pain			
	F	%	F	%	F	%		
<b>Age</b>								
a) 41years-45years	1	100	0	0.00	0	0.00	1	$\chi^2=16.45$ P=0.03* S
b) >45years-50years	2	66.7	1	33.3	0	0.00	3	
c) >50years-55years	5	71.4	2	28.6	0	0.00	7	
d) >55years-60years	3	60.0	2	40.0	0	0.00	5	
e) >60years-65years	0	0.00	12	85.7	2	14.3	14	
<b>Dietary pattern</b>								
a) Vegetarian	9	64.3	5	35.7	0	0.00	14	$\chi^2=9.24$ P=0.01** S
b) Non vegetarian	2	12.5	12	75.0	2	12.5	16	
<b>Gender</b>								
a) Male	2	18.2	8	72.4	1	9.1	11	$\chi^2=2.56$ P=0.27 NS
b) Female	9	47.4	9	47.4	1	5.3	19	
<b>Specific habits</b>								
a) Smoking	-	-	6	85.7	1	4.8	7	$\chi^2=5.50$ P=0.23 NS
b) Tobacco chewing	1	50	1	50	1	14.3	2	
<b>Duration of type 2 Diabetes Mellitus</b>								
a) 5years-10years	6	85.7	1	14.3	0	0.00	7	$\chi^2=15.42$ P=0.05* S
b) >10years-15years	4	40.0	6	60.0	0	0.00	10	
c) >15years-20years	3	50.0	3	50	0	0.00	6	
d) >20years-25years	0	33.3	2	66.7	1	33.0	3	
e) >25years	0	0.00	3	75.0	1	25.0	4	
<b>Associated illness</b>								
a) Hypertension	3	23.1	9	69.2	1	7.7	13	$\chi^2=2.34$ P=0.67 NS
b) Cardiac problems	5	45.5	5	45.5	1	9.1	11	
c) Hyperlipidemia	-	-	-	-	-	-	-	
d) Renal and respiratory problem	3	50.0	3	50.0	-	-	6	
<b>Nail condition</b>								
a) Normal	10	52.6	9	47.4	0	0.00	19	$\chi^2=7.84$ P=0.02* S
b) Thickened	1	9.1	8	72.7	2	18.2	11	

NS-Not significant, S-Significant (\*denotes significant at 5% level)

Association of Post Test Level of Lower Extremity Perfusion with Selected Demographic and Clinical Variables of Samples In Experimental Group and in Control Group, N=30

Demographic variables	Post test level of perfusion				Total	Chi square test
	Mild impairment		Normal			
	N	P (%)	N	P (%)		
<b>Age</b>						
a) 41years-45years	0	0	1	100	1	$\chi^2=9.98$ P=0.04* S
b) >45years-50years	0	0	3	100	3	
c) >50years-55years	1	14.2	6	85.8	7	
d) >55years-60years	2	40	3	60	5	
e) >60years-65years	10	71.4	4	28.6	14	
<b>Gender</b>						
a) Male	6	54.5	5	45.5	11	$\chi^2=0.88$ P=0.34 NS
b) Female	7	36.8	12	63.2	19	
<b>Dietary pattern</b>						
a) Vegetarian	5	35.7	9	64.3	14	$\chi^2=0.62$ P=0.43 NS
b) Non- Vegetarian	8	50	8	50	16	
<b>Specific habits</b>						
a) Smoking	5	71.4	2	28.5	7	$\chi^2=6.72$ P=0.03* S
b) Tobacco chewing	2	100	0	0	2	
<b>Duration of type 2 Diabetes Mellitus</b>						
a) 5years-10years	0	0	7	100	7	$\chi^2=10.04$ P=0.04* S
b) >10years-15years	4	40	6	60	10	
c) >15years-20years	5	83.3	1	16.7	6	
d) >20years-25years	2	66.7	1	33.3	3	
e) >25years	2	50	2	50	4	
<b>Associated illness</b>						
a) Hypertension	6	46.2	7	53.8	13	$\chi^2=0.30$ P=0.85 NS
b) Cardiac problems	5	45.5	6	54.5	11	
c) Hyperlipidemia	-	-	-	-	-	
d) Renal and respiratory problem	2	33.3	4	66.7	6	
<b>Nail condition</b>						
a) Normal	5	26.3	14	73.7	19	$\chi^2=6.11$ P=0.01** S
b) Thickened	8	72.7	3	27.3	11	

NS-Not significant, S-Significant (\*denotes significant at 5% level)

## 5. Conclusion

The findings of the present study revealed that there was a significant improvement in lower extremity perfusion and reduction in pain. Therefore it was concluded that Buerger Allen exercise was found to be effective on improving the lower extremity perfusion and reducing pain among patients with type 2 diabetes mellitus.

## References

- [1] Agarwal, B.L. (2011). *Text book of statistics*. (3rd ed.). New Delhi: CBS publishers.
- [2] Allen, A.W. (1930). Recent advances in the treatment of circulatory disturbances of the extremities. *Annals of surgery*, 92,931-946.
- [3] American Diabetes Association Consensus, (2013). Peripheral arterial disease in people with diabetes. *Clinical diabetes*, 22(4), 181-189.
- [4] Aravind, S. et al., (2013). *Diabetology and metabolic syndrome*. Retrieved from DOI:10.1186/1758-5996-5-57.
- [5] Ashok, P. et al., (2013). Peripheral vascular disease- a silent assassin. *JIACM*, 14(2), 111-114.
- [6] Basavanthappa, B.T. (2007). *Nursing theories*. (2nd ed) New Delhi: Jaypee Brothers.
- [7] Black, M.J. (2001). *Medical surgical nursing*. (2nd ed). Philadelphia: W.B. Saundar Company.
- [8] Brunner., & Suddarth. (2011). *Text book of medical and surgical nursing*. (11<sup>th</sup>ed.). Philadelphia: J.B. Lippincott Company.
- [9] Buerger, L. (1926). The circulatory disturbances of the extremities. *Annals of surgery*, 83, 157.
- [10] Chang, C. F. et al., (2015). Effect of buerger's exercises on improving peripheral circulation, a systematic review. *Open journal of nursing*, 5, 120-128. Retrieved from <http://dx.doi.org/10.4236/ojn.2015.52014>
- [11] Davis, K & Raymond, M. (2014). Retrieved from DOI: 10.1371/journal.pone.0105211
- [12] Deepa, S & Srikanth, G. (2014). Macrovascular and microvascular complications in newly diagnosed in type 2 Diabetes Mellitus. *Indian journal of clinical practice*, 25(7), 644-648.
- [13] Edward, B. et al., (2009). Diabetes care, peripheral arterial disease in diabetic and non diabetic patients- a comparison of severity and complication. *Diabetes journal*, 24(8), 1443-47.
- [14] Faris, I. et al., (2005). Increased subcutaneous flow for 24 hours. *European journal of nuclear medicine and molecular imaging*, 8(4), 15-17. Retrieved from <http://www.springerlink.com/content>
- [15] Gupta, S.P. (2003). *Statistical methods*. (32<sup>nd</sup>ed.). New Delhi: Sultan Chant & Son's Education Publishers.
- [16] Hoe, J. et al., (2012). Predictors of decrease in ankle-brachial index. *Diabetic medicine*, DOI:10.1111/j.1464-5491.2012.03705.x
- [17] Jemcy, J & Rathiga, A. (2015). Effectiveness of buerger allen exercise. *Indian journal of current research and academic review*, 3(4), 358-366.
- [18] Joshua, A. et al., (2013). DOI: <http://dx.doi.org/10.1093/eurheartj/eh142.2444-2452>.
- [19] Judy, J. (2013). An experimental study to assess the effectiveness of buerger's Allen exercise on improving wound healing, Kerala (Unpublished dissertation).
- [20] Kawasaki, T. et al., (2013). The effect of different positions on lower limbs skin perfusion pressure. *Indian J plast surg*. Cited 2015 Feb 21, 46, 508-12. DOI: 10.4103/0970-0358.121995.
- [21] Lawrence, H. et al., (2005). Value of exercise in peripheral arterial disease, *JAMA*, 153(1), 10-12.
- [22] Lewis, et al., (2010). *Medical surgical nursing* (7th ed.). New Mexico: Mosby Publications.
- [23] Likhila, A. (2013). Effectiveness of structure teaching programme regarding knowledge on varicose vein and practice of buerger's Allen exercise in improving peripheral perfusion among workers, Mangalore (Unpublished dissertation).
- [24] Luckman, J. (1998). *Medical surgical nursing*, (2<sup>nd</sup>ed.). Philadelphia: W B Saunders.
- [25] Marso, P. et al., (2010). Prevalence of peripheral arterial disease among people attending diabetic clinics, *BMB*, 33(4), 1-9.
- [26] Matfin, P. et al., (2009). Peripheral vascular disease and diabetes. Retrieved from <http://diabetes.hiddk.nih.gov/dm/pubs/america>
- [27] Mohammed, E. et al., (2013). DOI:10.1007/s10753-013-9746-4.
- [28] Mohan, V. et al., (2013). Indian scenario, *Indian journal of medicine*, 125, 225-230. Retrieved from [www.indianjmedsci.org](http://www.indianjmedsci.org)
- [29] Mundet, X. et al., (2008). Prevalence and incidence of chronic complications and mortality in a cohort of type 2 diabetic patients in Spain, 2(3), 135-140.
- [30] Neil, J.S & Will, G. (2012). Effect of different mode of exercise programme on glucose control and risk factors for complication in type 2 diabetic patients. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc>
- [31] Pat, F & Diana, B. (2011). Peripheral vascular disease and type 2 diabetes. doi:10.1016/j.jacc.2012.09.065
- [32] Polit, D.F., & Hungler, B.P. (2004). *Nursing research principles and method*. (6<sup>th</sup>ed). Philadelphia: J.B. Lippincott, company.
- [33] Potter & Perry, (2010). *Fundamentals of nursing*. (6<sup>th</sup>ed). Philadelphia: Mosby publication.
- [34] Premalatha, S. et al., (2009). Prevalence and risk factors of peripheral vascular disease in south Indian population, *Diabetes care*, 23(9), 1295-1299.
- [35] Rajabharan, Y. et al., (2008). Risk factors and complications of type 2 DM. Retrieved from <http://www.niper.ac.in/riskfactor>
- [36] Sandeep, S. et al., (2013). *DMSO*, 6, 327-328. DOI:10.2147/DMSO.551325
- [37] Steven, P. et al., (2010). Peripheral vascular complications among type 2 Diabetes Mellitus patients, *Journal of diabetes and metabolism*, 2(1), 1-5.
- [38] Sharma, R. (2009). *Methodology of education research*. (3rd ed). New Delhi: Vora publishers.

- [39] Shen, Q. et al., (2010).Prevalence of PVD in diabetes and IGR subjects:a community based study, 86(22), 1530-33.
- [40] Sherin, H. & Mohamed, M.(2012).Comparison between two vascular rehabilitation program, Bull, *Faculty of physical therapy*, Cairo University, 17(1), 7-15.
- [41] Tota, K. et al., (2013).The effect of different positions on lower limbs skin perfusion pressure, *Indian J plast surg*, 46, 508-512.
- [42] William, Y.T& Wang, S.S. (2010). Effects of buerger allen exercises on peripheral vascular function in patients with peripheral arterial insufficiency, *National Science Council Report*, Taipei, 1-15.
- [43] World Health Organisation. (2012)The prevalence of diabetes Mellitus, Global health Estimates.
- [44] Zhaolan, L. et al., (2012). DOI: 10.1161/JAHA.112.002832.
- [45] Zimmermann, W.M et al., (2011). Course of occlusive arterial disease in diabetes, *Diabetes care*,13, 143-52.

