

Comparative Study of Lipid Alterations in Flight Muscles of *Leucopholis lepidophora* and *Oryctes rhinoceros*

S. B. More

Department of Zoology, P.V.P. Mahavidyalaya, Kavathe Mahankal, Dist. Sangli (M.S.), India 416 405

Abstract: The neutral lipids (NL) and phospholipids (PL) with their constituents were studied in the flight muscles of male and female adults of *Leucopholis lepidophora* and *Oryctes rhinoceros*, by employing thin layer chromatography (TLC) and bioassay technique. The quantity of neutral lipids in male and female flight muscles of *L. lepidophora* were measured 39.192 and 40.039 whereas the *O. rhinoceros* exhibits 57.501, 45.894 mg/gm wet weight of tissues respectively. The main component of neutral lipids was triacylglycerol. Similarly the phospholipid values in male and female flight muscles of former and later species were 23.163, 22.999, 17.367 and 15.58 mg/gm wet weights of tissues respectively. The NL: PL ratio in male and female adult flight muscles of *L. lepidophora* was recorded to 3:1 whereas in *O. rhinoceros* 4:1. The neutral lipids found in six forms. Triacylglycerol was the main component in both species. It exhibits 29.60 and 32.08 48.40, 40.05 mg/gm wet weight of tissues in male and female flight muscles of former and later species respectively. monoacylglycerol, diacylglycerol, cholesterol, were moderate and cholesterol ester and free fatty acids low in quantity. Phospholipids exhibited seven constituents; phosphatidyl-choline and phosphatidyl-ethanolamine in high concentration in both species. Lysophosphatidyl-choline and sphingomyelin moderate in concentration and phosphatidyl-inositol, phosphatidyl-serine and phosphatidic acid low in quantity.

Keywords: lipids, Male and female flight muscles, Thin layer chromatography, *Leucopholis lepidophora* and *Oryctes rhinoceros*

1. Introduction

The white grub larvae of *L. lepidophora* are important pest of sugar cane in general and south west Maharashtra in particular [2] The Larval development persists of 225-295 days. It damages the roots of different crops. So it becomes a problem of sugarcane cultivators, especially in Maharashtra. The white grub have become known polyhagus and found in a particular type of agro ecosystem. In India Lefray [25] Ghosh [19] studied the white grubs of sugarcane. White grubs are most destructive insect pest all over the world. *Oryctes rhinoceros* is the pest of coconut trees. Nirula made a thorough inspection of one million coconut palms heavily infected by the rhinoceros beetle and estimated the annual loss as of \$ 6, 00,000 to the coconut planters. Recent report of the damage caused by this beetle in Assam and Andaman Islands are very alarming [47] According to government of Assam, rhinoceros beetle presents a colossal problem to the coconut industry attacking both young and adult trees. In many insects such as *Manduca sexta*, lipids are major fuel for flight which is mobilized as diacylglycerol (DG) into the circulation [3], [8]. DG is derived from triacylglycerol (TG) stores in the fat body and constitutes the main form in which fatty acids are mobilized to the site of utilization, for example, flight muscle. The mobilization of triglyceride stores during flight is controlled by adipokinetic hormone [18], [40] which increases the rate of triglyceride hydrolysis and promotes a concomitant release of diacylglycerol into the hemolymph. Increases in lipase activity of fat body homogenates after initiation of flight or AKH injection have been observed in several insects [5], [6]. Lipid is bio-chemically important component of the insect. Recently it has been observed that, many of the pesticides and the insecticides were accumulated in lipid. The role of sterol in insect development and metamorphosis was described by

Madariaga, *et al*; [29] and Dwivedy [16],[17] The significance of phospholipid with PC and PE was explained by Turunen [38], [27], [32] The laboratory of the Department of zoology, Shivaji University Kolhapur is actively engaged in research on the white grubs. Bhanot [10] studied the biology of the white grub in Kolhapur region; while Patil [34] investigated biochemical aspects like proteins, carbohydrates and enzymes. The literature survey indicated that, lipid of this species have not been studied. Hence in the present study the lipid were investigated in the flight muscles of *L. lepidophora* and *O. rhinoceros* Lipids are bio-chemically important components of insect. Lipid performs a variety of functions in insect physiology. Triacylglycerol is utilized for biological energy [14] The significance of phospholipids with PC and PE was explained by Locke and Krishnan [27]. Flight is important for insect dispersal and flight pushes the envelope of organism design [12]. The main insect muscles are the leg muscles. Flight muscle is one of the most widely studied light-related tissues in insects mainly because of the variation in the metabolic activity of the different types of muscle [36], [39], [30]. The present report describes the comparative study of lipids in flight muscles of *L. lepidophora* and *O. rhinoceros*.

2. Methods and Materials

The male and female adults of *L. lepidophora* were collected from sugarcane field of Sangrul village, 18 k.m. away from Kolhapur city (M.S.). Adults were used for flight muscle lipid extraction. Similarly the prepupae and adults of *O. rhinoceros* were collected from dunghill of Agriculture College, near the Shivaji University, campus Kolhapur, (M.S.). They were reared at laboratory and adults were used for flight muscle lipid extraction.

A) Extraction of Lipids

The flight muscles of male and female adults of above both species were weighed and homogenized with 20 ml of chloroform-methanol (2:1 v/v) at room temperature. The homogenates were allowed to stand for 2-3 hours at 4⁰c and filtered. The filtrate was washed according to Floch *et.al*; [21] and evaporated in vacuum at 40⁰c. The lipid samples were weighed and preserved at -20⁰c until further use. The total lipid in the sample was determined gravimetrically.

B) Separation of Neutral Lipids and Phospholipid

The neutral lipids and phospholipids were separated by thin layer chromatography (TLC) using silica gel G and about 200 mesh containing CaSO₄, as a binder, (E Merck Germany). The TLC plates (20 X 20 cm) were prepared according to Wagner *et.al* [43]. The known quantities of samples dissolved in chloroform were applied with Hamilton's micro syringe (No.8206-B) on activated plates. For neutral lipid the plates were developed in hexane (B.P. 65-70⁰c) diethyl ether-acetic acid (85:15:2 v/v) as recommended by Gloster and Flecker [20]. The phospholipid plates were developed in chloroform-methanol-ammonia (115:45:5 v/v) as recommended by Barwal and Kalra [7]. The standards of neutral lipids and phospholipids (Sigma,U.S.A.) were co-chromatographed in each respective run and then plates were kept in iodine chamber for identification of individual spots of lipids.

C) Estimation of Neutral Lipids and Phospholipids.

The iodine was allowed to evaporate and the silica gel from the individual spots of glycerides was scraped and eluted in 1 ml of diethyl-ether and assayed according to Viogue and Holman [42]. The cholesterol and its ester were estimated according to Abell *et.al* [1]. The rest of the neutral lipid components were assayed titrometrically by the method of Skipski *et.al* [35]. The phospholipid was determined by the method of Marinetti [28].

3. Results

I) Neutral Lipids

The TLC separation of various neutral lipid components are illustrated in plate No.1, Fig. A; whereas, Table No.1 exhibits quantitative variations in the neutral lipid components. The neutral lipids in male and female flight muscles of *L. lepidophora* (herein after called as former species) were measured 39.192 and 40.039 mg/gm.wet weight of tissues respectively; whereas the neutral lipids in male and female flight muscles of *O. rhinoceros* (herein after called as later species) were measured 57.501 and 45.894 mg/gm.wet weight of tissues respectively. The neutral lipids consist of six components. Triacylglycerol (TG) being the major component in above two species. Monoacylglycerol (MG), diacylglycerol (DG) and cholesterol (CHO) were found moderate in concentration; whereas free fatty acids (FFA) and cholesterol ester (CE) were occurred low in quantities. The TG concentration in male and female flight muscles of *L. lepidophora* was about 29.60 and 32.08; whereas in *O. rhinoceros* 48.40 and 40.05 mg/gm wet weight of tissues respectively.

II) Phospholipids

The phospholipids are illustrated in plate No.1, Fig B and Table 2. The phospholipids in male and female flight muscles of former and later species were measured 23.163, 22.999 and 17.367, 15.58 mg/gm.wet weight of tissues respectively. The TLC separation of phospholipids constitute following constituents; phosphatidyl-choline (PC), phosphatidyl-ethanolamine (PE), Lysophosphatidyl-choline (LPC), sphingomyelin (SPG), phosphatidyl-inositol (PI), phosphatidyl-serine (PS) and phosphatidic acid (PA). Among the phospholipids PC and PE were predominant. In male and female flight muscles of former and later species they were measure about 376.1, 345.3, 345.1, 370.5 and 282.1, 260.3 , 275.1 , 250.5 $\mu\text{g -P/ gm}$ wet weight of tissues respectively. The LPC and SPG were estimated in moderate concentration, whereas PI, PS and PA less in amount.

PLATE - 1




Table 1

Alterations in total lipids, neutral lipids and its components in the male and female Flight muscles of *L. lepidophora* and *O. rhinoceros*

Lipid Type	<i>Leucopholis lepidophora</i>		<i>Oryctes rhinoceros</i>	
	Male Flight Muscles	Female Flight Muscles	Male Flight muscles	Female Flight muscles
Total Lipids	62.355 ± 2.44	63.038 ± 2.30	74.868 ± 2.74	61.474 ± 2.60
Neutral Lipids	39.192 ± 4.15	40.039 ± 1.23	57.501 ± 4.15	45.894 ± 2.23
MG	2.458 ± 0.20	1.465 ± 0.07	1.615 ± 0.15	1.335 ± 0.06
CHO	1.626 ± 0.18	1.868 ± 0.09	1.826 ± 0.09	0.868 ± 0.06
FFA	0.890 ± 0.08	1.324 ± 0.03	0.591 ± 0.04	1.639 ± 0.04
DG	3.346 ± 0.22	2.334 ± 0.10	3.867 ± 0.16	1.834 ± 0.06
TG	29.60 ± 2.79	32.08 ± 1.10	48.40 ± 3.832	40.05 ± 1.30
CE	1.272 ± 0.05	0.968 ± 0.05	1.202 ± 0.05	0.168 ± 0.01

The values for total lipids, neutral lipids and its components are expressed as mg/gm.wet weight of tissues.

Table 2

Alterations in phospholipids and its constituents in male and female flight muscles of *L. lepidophora* and *O. rhinoceros*

Lipid Type	<i>Leucopholis lepidophora</i>		<i>Oryctes rhinoceros</i>	
	Male Flight Muscles	Female Flight Muscles	Male Flight Muscles	Female Flight Muscles
Phospholipids	23.163 ± 0.32	22.999 ± 0.93	17.367 ± 0.66	15.58 ± 0.83
PI	15.70 ± 0.75	20.70 ± 0.68	15.70 ± 0.75	10.70 ± 0.62
LPC	66.22 ± 2.79	55.47 ± 3.12	56.21 ± 2.76	35.47 ± 2.12
SPG	78.36 ± 2.89	70.22 ± 2.73	51.31 ± 2.56	30.22 ± 1.73
PS	23.87 ± 0.79	30.28 ± 0.99	16.18 ± 0.75	10.24 ± 0.51
PC	376.1 ± 12.57	345.1 ± 14.9	282.1 ± 4.57	275.1 ± 13.9
PE	345.3 ± 14.01	370.5 ± 14.7	260.3 ± 13.01	250.5 ± 13.7
PA	20.97 ± 0.68	27.70 ± 2.03	12.87 ± 0.64	10.70 ± 1.03

The values of phospholipids are expressed as mg/gm. wet weight of tissues; whereas, values of individual

constituents are expressed as $\mu\text{g-P/gm}$. wet weight of tissues.

4. Discussion

The male flight muscles of *L. lepidophora* and *O. rhinoceros* exhibited high amount of total lipids than the female.

The NL: PL ratio in male and female adult flight muscles of *L. lepidophora* was recorded to 3:1 whereas in *O. rhinoceros* 4:1. Indicated that the neutral lipids were dominated over the phospholipid in both species. Among the neutral lipids TG constitute the major component. The TG of the neutral lipids in male and female flight muscles of both species was 29.60, 32.08 and 48.40, 40.05 mg/gm wet weight of tissues respectively. These findings agreed with the findings on *T. castaneum* [7]. Domroese and Gilbert [15] and Lawrence et.al [24], studied some aspects of lipid catabolism in moth *H. cecropia*. He reported that, the utilization of lipids as a fuel for flight. Lipid is the available substrate as well as the preferred substrate in flight-muscle metabolism in male moths. TG exhibits high concentration in male and female flight muscle and for utilization as energy source [22], [13]. Phospholipids are expressed as mg/gm wet weight of tissues and their values in male and female flight muscles of *L. lepidophora* were 23.163 and 22.999 and in *O. rhinoceros* were 17.367 and 15.58 mg/gm wet weight of tissues respectively. In the present investigation in both species male flight muscles exhibited high phospholipid contents than female flight muscles. Among the phospholipids the value PC and PE were major constituents in both species. Kallapur et al [23] studied effect of environmental temperature on lipid composition of flight muscle mitochondria in *S. gregaria*, and observed that elevated temperature resulted in depressed level of PC and increased level of PE. Our findings stated that, PC: PE ratio was 1:1 which indicated that the PC and PE are equal in their values in both species. Beenackers [9] studied transport of fatty acids in *L. migratoria* during sustained flight. He reported that, during flight a considerable increase in fatty acid content was indication of an efficient transport of various species of insects by many investigators, such as Van Handel [41] Locust *migratoria migratorioides*, Mwangi and Gondsworthy [31] *Nilaparvata lugens*, Chi [11] and *Rhaphalosiphum maidis* Liquido and Irwan, [26] Cheng -Jijiang [12] *G. firmus*., Zera et.al. [44] Zhang et.al [45] and Zaho and Zera [46] In insects flight activity usually depends on lipids [13]. Insect flight muscles do not directly store lipids and require an efficient system to transport hydrophobic fatty acids to the muscle cell which in locusts is mediated by apolipoproteins. Lipid utilization during sustained flight has been studied on lepidopteran and orthopteran species by Norbert [33] He reported that, in migrating lepidopteran and orthopteran insects, lipid is the preferred fuel for sustained flight activity. Diacylglycerol is delivered by lipophorin to the flight muscle and hydrolyzed to free fatty acid and glycerol. Lipid is the available substrate as well as the preferred substrate in flight-muscle metabolism in male moth [15] Active fatty acid activating enzymes is present in flight muscle, and fatty acid oxidation in *H. cecropia* is discussed in relation to vertebrate and other invertebrate systems. [17]. Phospholipid synthesis during

flight muscle development in American silkmoth *H. cecropia* was studied by Thomas and Gilbert [37]. They found that the major phospholipid fraction of flight muscles and sarcosomes were PC and PE. Atkins studied lipid loss of flight muscle in the beetle *Dendroctonus pseudotsugae* and reported that the flown beetles contained significantly less lipids [4].

5. Acknowledgements

Author thanks to Prof. V A. Sawant Dept of Zoology Shivaji University Kolhapur for his encouragement.

References

- [1] Abell, L.L., Levy, B.B., Brodie B.B., and Kendall, F.F. : A simplified method for the estimation of total cholesterol in serum and demonstration of its specificity., *J. Biol. Chem.* **195**, 357-359 (1952).
- [2] Abraham V. A. and Rajendran T. P. White GrubS Newsletter, 2; 1978.
- [3] Arrese E. L., Wells M. A.. 1997. *Adipokinetic hormone-induced lipolysis in the fat body of an insect, Manduca sexta: synthesis of sn-1,2-diacylglycerols.* *J. Lipid Res.* **38**: 68–76.
- [4] Atkins M. D. : Lipid loss with flight in the Douglasfir beetle. *D. pseudotsugae*, *Can. Entomol.* **101**, 164-165.1969.
- [5] Auerswald L, Gade G. Endocrine control of TAG lipase in the fat body of the migratory locust, *Locusta migratoria*. *Insect Biochem Mol Biol.* 2006; **36**:759–68.
- [6] Auerswald L, Siegert KJ, Gade G. Activation of triacylglycerol lipase in the fat body of a beetle by adipokinetic hormone. *Insect Biochem Mol Biol.* 2005;**35**:461–70.
- [7] Barwal, R.N. and Kalra, R.L. : Lipids lindane susceptible and resistant strains of *T. castaneu.*, *Indian J. Expt. Biol.*, **26**, 228-234 (1988).
- [8] Beenackers A. M. T., Van der Host D. J., Van Marrewijk W. J. A. . 1985. *Insect lipids and lipoproteins and their role in physiological processes.* *Prog. Lipid Res.* **24**: 9–67.
- [9] Beenackers, A. M. T. : Transport of fatty acids in *L. migratoria*. *J. Insect Physiol.* **15**, 353-361.
- [10] Bhanot P. K. :Studies on some aspects of the Biology of white Grubs , A Ph. D. thesis , Shivaji University Kolhapur(India), 1993.
- [11] Chi. C.R. : Lipid a fuel of flight in brown plant hopper *Nilaparvata lugens* . *ACTA, Biol. Sin.* **26**. 42-48.
- [12] Cheng-Jijiang, Bao-Changzang, Wen-Fengchen, Qing-Wenzhang, Zhang-Wuzhao, Chun-Juan., and Jie-Pingli: Variations in the ultrastructure of the flight muscles of the polymorphic cricket, *Gryllus firmus*(Orthoptera: Gryllidae *Eur. J. Entomol.* **109**: 579–586, 2012.
- [13] Dickinson M. H. and Tum S.: The function of dipteran flight muscle – turning axis and gearbox of the wing base (Insecta, Diptera). *Compar. Biochem. Physiol. (A)* **116**: 223–238.
- [14] Downer RGH, Physiological and environmental consideration in insect bioenergetics, In energy metabolism in Insects, Edited By R.G.H. Downer, 1981, 1-17, Plenum Press New York, 1997.

- [15] Domroese K. A. and Gilbert L. I. : The role of lipid in adult development and flight muscle metabolism in *H. cecropia*, *J. Exp. Biol.* 41, 573-590, 1964.
- [16] Dwivedy A. K. Dietary cholesterol requirements of housefly *M. domestica* larvae, *J. Insect Physiol.*, 21, 1985-1990. 1975.
- [17] Dwivedy A. K.: Effect of cholesterol deficiency on the composition and fatty acids of the house fly, *M. domestica* *Insect Physiol*, 23, 549-557; 1977.
- [18] Gade G. Regulation of intermediary metabolism and water balance of insects by neuropeptides. *Annu Rev Entomol.* 2004;49:93-113.
- [19] Ghosh G. L. : *Ind. J. Agri. Sci.*, 7 907-931; 1937.
- [20] Gloster, J. Flectar, R.F., Quantitative analysis of serum lipids with thin layer chromatography., *clin. Chim. Acta.*, **13**, 235-240 (1966).
- [21] Folch, J., Lees, M. and Sloane-Stanley A simple method for the isolation and purification of the total lipid from animal tissue. *G.H. J.Biol. Chem* **226** (1957) 497-509.
- [22] Hoffman, A.G.D. and Downer, R.G.H., : End product specificity of TG lipases from intestine fat body muscle and hemolymph of the American cockroach, *P. Americana., Lipids.*, **14**, 893-899 (1979).
- [23] Kallapur V. L. , Downer R. G. H. , George J. C. and Thompson J. E. : Effect of environmental temperature on the lipid composition of flight muscle mitochondria of *S. gregaria* , *Insect Biochem.* 12, 115-122. 1982.
- [24] Lawrence A. Domroese and Lawrence I. Gilbert: The Role of Lipid in Adult Development and Flight-Muscle Metabolism in *Hyalophora Cecropia* *Journal of Experimental Biology* 1964 41: 573-590;
- [25] Lefroy H. M. : *Ent. Memories*, 11, 143-146; 1900.
- [26] Liquido N. J. and Irwin M. E. : Lipid content and utilization during sustained flight of corn leaf aphid *R. maidis* , *Ann. Appl. Biol.* 108, 449-460.
- [27] Locke M and Krishnan N, J. Phospholipids and fatty acid composition of *T. granarium* and biosynthesis of fatty acids during development, *Comp. Biochem. Physiol.*, 39, 1971, 183-194.
- [28] Marintti, G.V., : ed. "Lipid Chromatography Analysis". Vol I *Marcel Dekker. Inc.*, New York. (1967).
- [29] Madariaga, M. A. Mata F. Municio A. M. and Ribera A. : Effect of lipid composition of the larval diet on the fatty acid composition during development of *C. capitata*. *Insect Biochem.* 2, 249-256, 1972.
- [30] Menitel T., Duche., Stypah., Mulleru. and Wegenerg. :Central modulatory neurons control fuel selection in flight muscle of migratory locust.*J. Neurosci.* 23: 1109-1113., 2003
- [31] Mwangi R. W. and Goldsworthy G. J. : Diacylglycerol-transporting lipoproteins and flight in *L. migratoria migratorioides*. *J. Insect Physiol.*, 27, 47-50.1981
- [32] Niemierko S., Woldawer P. and Wojtaczak A. F. : Lipid and phosphorus metabolism during growth of the silkworm . *ACTA Biol. Biol. Exper.* 17, 255-276; 1956.
- [33] Norbert H. Haunerland : Transport and Utilization of Lipids in Insect Flight muscles, *Comp.Biochem*, Vol. 11 B7, No. 4 PP.475- 482, 1997.
- [34] Patil S. B. Studies on some systems of *H. serrata* F. Coleoptera: Scarabaeidae; A Ph. D. thesis , Shivaji University Kolhapur(India), 1997.
- [35] Skipski, V.P., Barclay, M., Reichman, E.S. and Good, J.J., : *Biolchem, Biophys. Acta.*, **137**, 80 (1967).
- [36] Sasktor B. . : Regulation of intermediary metabolism with special reference to the control mechanisms in insect flight muscle. *Adv. Insect Physiol.* 7: 267-347., 1970
- [37] Thomas K. K. and Gilbert L. I.: Phospholipid synthesis during flight muscle development in the American silkmoyh *H. cecropia* , *Comp. Biochem. Physiol.* 21, 279-290, 1967.
- [38] Turunen S. : The role of labeled dietary fatty acids and acetate in phospholipids during the metamorphosis of *P. brassicae*. *J. Insect physiol.*, 2327- 2340. 1973.
- [39] Usher Wood P.N.R. (ed.) 1975: *Insect Muscle*. Academic Press, London, 621 pp 1975.
- [40] Van der Horst DJ. Insect adipokinetic hormones: release and integration off light energy metabolism. *Comp Biochem Physiol B.* 2003;136:217-26.
- [41] Van.Handel E. and Nayer J. K. : Direct use of carbohydrate during sustained flight in yhe moth, *S. frugiperda*. *Insect Biochem.*, 2, 203-208.
- [42] Viogue, E, and Holaman, R.T., : Quantitative estimation of esters by thin layer chromatography., *J.Am.oil. chemistis. Soc.*, **39**, 68-66 (1962).
- [43] Wagner, H., Horhammer, L. and Wolfe, P., : Thin layer chromatography of phospholipids and glycolipids., *Biochem.*, **2**, 175-184 (1961)
- [44] Zera A. J., Sall J. & OtoO K. : Biochemical aspects of flight and flightlessness in *Gryllus*: flight fuels, enzyme Activit ies and electrophoretic profiles of flight muscles from flight-capable and flightless morphs. *Insect Physiol.* 45: 275-285. 1999.
- [45] Zhang B.C., Jiang C.J., Zhang Q.W. and Zhao Z.W. : Varia-tions in fuel use in the flight muscles of wing-dimorphic *Gryllus firmus* and implications for morph-specific dispersal. *Enviorn. Entomol.* 40: 1566-1571., 2011.
- [46] Zhao Z.W. & Zera A.J. : Differential lipid biosynthesis underlies a tradeoff between reproduction and flight capa-bility in a wing-polymorphic cricket. *PNAS* 99: 16829-16834.2002.
- [47] Nirula K. K.: Investigations on the pests of coconut palm. *Indian Coconut Journal.* **9**, 30-37, 1956.