

Singular Two-Point Boundary Value Problems of Finite Element method

Weiwei Zhang

University of Sciences, Hebei University of Technology, Beichen district, Tianjin 300401, China

Abstract: In this paper, a class of singular two-point boundary value problems using the Galerkin method, using hierarchical grid method in discrete space, using one yuan for discrete function respectively, is obtained the general grid on the norm of the error convergence.

Keywords: singular, two-point boundary value problems, Galerkin method, super convergence, Finite Element method

1. Introduction

In this paper a yuan in hierarchical grid numerical solving the following singular two-point boundary value problems.

$$\begin{cases} -u''(x) - \frac{b}{x}u'(x) + q(x)u(x) = f(x), & 0 < x < 1 \\ u'(0) = 0, \quad u(1) = 0 \end{cases} \quad (1.1)$$

Here is the constant, $b > 0$ is responsible $q(x)$ for the function, $f(x)$ for a given function. [1], [2], [3] know if there is the only solution of this problem and smooth.

The singular boundary value problems can be caused by high dimensional non simplified spherically symmetric problems, the first side value problem of three dimensional elliptic equations:

$$\begin{cases} -\nabla(a\nabla u) = f & \text{in } \Omega, \\ u|_{\partial\Omega} = 0 \end{cases}$$

Through coordinate transformation into this problem.

This equation numerical Xie Ben Wen from the following several aspects: one, layered grid; Second, the equation of variational form; Three, equations of discrete; Four, error estimate.

Duran [7]in 2006 for the first time put forward the hierarchical grid, the grid to adjacent cell changes slowly, without considering the change of the thickness of the grid. Chuan-miao Chen[13]on special grid discussed the convergence of the differential error norm and superconvergence. Zhang xu, [1],[12] on the basis of chuan-miao Chen made a general grid model and the error of the maximum modulus difference of convergence.

In this paper, on the basis of zhang xu, do the general layered grid on the grid on the norm of the error convergence and convergence.

2. The Discrete

By the Green points male, can get the variational equation

$$a(u, v) = \int_{\Omega} x^b f v dx, \quad \forall v \in H^1(\Omega) \quad (2.1)$$

there

$$a(u, v) = \int_0^1 x^b u'(x)v'(x) + x^b u(x)v(x)dx \quad v \in H^1$$

$$\langle f, v \rangle = \int_0^1 x^b f v dx \quad v \in H^1$$

By [5] the problem (1) and (2.3), and other equivalent. The interval are partitioned into coarse grid and fine grid:

$$\Omega_1 = [0, \frac{1}{n}], \Omega_2 = [\frac{1}{n}, 1]. \text{The fine grid } \Omega_2 \text{ for subdivision:}$$

$$\text{average composition, } \frac{1}{n} = x_1 < \dots < x_n = 1.$$

The i remember as a unit $I_i = [x_{i-1}, x_i]$, $h_i = \frac{1}{n}$. The same

$$\text{as } \Omega_1, 0 = x_{00} < x_{01} < \dots < x_{0n} = \frac{1}{n}. \text{The 1i remember as a}$$

$$\text{unit } I_{1i} = [x_{1i-1}, x_{1i}], h_i = \frac{1}{n^2}.$$

Define the finite dimensional test space V_h is piecewise polynomial in hierarchical grid space,

$$V_h = \{v \in H_0^1(\Omega) : v|_{\Omega_i} \in P_{k-1}(I_i), 1 \leq i \leq n\}$$

Here $P_{k-1}(I_i)$ is polynomial representative the i remember as

$$\text{a unit } I_i, V_h \subset H^1.$$

The equation can be turned into

$$a(u_h, v) = \int_0^1 x^b f v dx \quad v \in H_0^1 \quad (2.2)$$

$$\sum_{i=1}^n a(u_i, v_i) = \int_0^1 x^b f v_i dx, \quad i = 1, 2, \dots, n$$

$$\text{Norm: } \|v\|_{m,p,\Omega} = \left\{ \sum_{|\alpha| \leq m} \|D^\alpha v\|_{0,p,\Omega}^p \right\}^{1/p}$$

$$\text{Semi-norm: } |v|_{m,p,\Omega} = \left\{ \sum_{|\alpha|=m} \|D^\alpha v\|_{0,p,\Omega}^p \right\}^{1/p}$$

$$\text{Weight norm: } \|v\|_{m,p,\Omega} = \left\{ \sum_{|\alpha| \leq m} x^\alpha \|D^\alpha v\|_{0,p,\Omega}^p \right\}^{1/p}$$

$$\text{Weight semi-norm: } |v|_{m,p,\Omega} = \left\{ \sum_{|\alpha|=m} x^\alpha \|D^\alpha v\|_{0,p,\Omega}^p \right\}^{1/p}$$

3. The Convergence of Solution

3.1 Preliminary Knowledge

Lemma1[5] We assumption $u(x)$ and $u_h(x)$ are the solution of vanational problem and the solution of the discrete problem. Here, $\exists C = const > 0$, can be used

$$\|u - u_h\| \leq C \inf_{v_h \in V_h} \|u - v_h\|$$

Lemma2[8],[9],[10] Construct the interpolation postprocessing operator Π_{2h}^k , here $u_I \in V_h$ is the finite element interpolation for u , k is the number of interpolation polynomia, so

$$\begin{aligned} C &= const > 0, \\ \|\Pi_{2h}^k u - u\|_l &\leq Ch^{r+1-l} \|u\|_l \\ \|\Pi_{2h}^k v\|_l &\leq C \|v\|_l, \quad \forall v \in V_h \\ \Pi_{2h}^k u_I &= \Pi_{2h}^k u \end{aligned}$$

When $l = 0, 1, 0 \leq r \leq k$.

Lemma3[6] Assumption $u(x)$ and $u_I(x)$ is the solution of variational problem and linear interpolation,

$$a(u - u_I, v) \leq O(h^2) \|u\|_3 \|v\|_1, \quad v \in V_h$$

$\forall p \geq 2$ and $b > 0$.

3.2 Error estimation for $u - u_I$

Theorem1 Assumption $u(x)$ and $u_I(x)$ is the solution of variational problem and linear interpolation,

$$|u - u_I|_1 \leq Ch^2 \|u\|_3$$

$\forall p \geq 2$ and $b > 0$.

From Lemma3 :

$$\begin{aligned} a(u - u_I, v) &= \int_0^1 [x^b (u - u_I)' v'(x) + x^b (u - u_I) v(x)] dx \quad v \in H^1 \\ \int_0^1 x^b (u - u_I)' v'(x) dx &< \int_0^1 [x^b (u - u_I)' v'(x) + x^b (u - u_I) v(x)] dx \quad v \in H^1 \end{aligned}$$

$$a(u - u_I, v) \leq CO(h^2) \|u\|_3 \|v\|_1$$

so:

$$|u - u_I|_1 \leq Ch^2 \|u\|_3.$$

proof: In order to discuss on local unit, we introduce the following auxiliary function

$$\phi(x) = \begin{cases} \int_{x_j}^{x_{j+1}} v'(t) dt & 0 \leq x \leq x_{j-1} \\ \int_{x_j}^x v'(t) dt & x_{j-1} \leq x \leq x_j \\ 0 & x_j \leq x \leq 1 \end{cases}$$

The auxiliary function $\phi(x) \in V_h$,

Use the affine transformation $x = x_{i-1} + h_i \xi$, here $\xi \in [0, 1]$ in I_i .order $v_h = \phi(x)$, then

$$x^b = (x_{i-1} + h_i \xi)^b = (x_i \xi + (1-\xi)x_{i-1})^b \geq x_i^b \xi^b$$

so

$$\begin{aligned} \int_0^1 x^b (u - u_I)' v'_h dx &= \int_0^1 x^b (u - u_I)' \phi' dx \\ &= \int_{x_i}^{x_{i+1}} (v'(x))^2 dx \\ &\geq h_i x_i^b \int_0^1 t^b (v'(x))^2 dx \\ &\geq Ch_i x_i^b \max |v'|_1 \end{aligned}$$

and because

$$\begin{aligned} &\int_0^1 x^b (u - u_I)' v'(x) dx \\ &= \sum_{i=0}^n \int_{x_i}^{x_{i+1}} x^b (u - u_I)' v'(x) dx \\ &= \sum_{i=0}^n [x^b (u - u_I) v(x) \Big|_{x_i}^{x_{i+1}} - \int_{x_i}^{x_{i+1}} b x^{b-1} (u - u_I)' v'(x) dx] \\ &= \sum_{i=0}^n -b x^{b-1} \int_{x_i}^{x_{i+1}} (u - u_I)' v'(x) dx \\ &\leq C x^{-1} \sum_{i=0}^n \int_{x_i}^{x_{i+1}} (u - u_I)' v'(x) dx \\ &\leq C x^{-1} O(h^2) \|u\|_3 \|v\|_1 \end{aligned}$$

3.3 Error estimation for $u_h - u_I$

Theorem2 Assumption $u_h(x)$ and $u_I(x)$ are discrete problems and the solution of a linear interpolation,
 $\forall p \geq 2$ and $b > 0$, then

$$|u_h - u_I|_1 \leq Ch^2 \|u\|_3$$

proof: Take coefficient of variable $x = x_e$, because (2.1),
 (2.2)

$$a(u, v) = \int_0^1 x^b u'(x)v'(x) + x^b u(x)v(x)dx \quad v \in V_h$$

$$a(u_h, v) = \int_0^1 x^b u_h'(x)v'(x) + x^b u_h(x)v(x)dx \quad v \in V_h$$

Two type is presupposed regularity,

$$a(u - u_h, v) = 0, \quad \forall v \in V_h$$

so

$$a(u - u_I + u_I - u_h, v) = 0, \quad \forall v \in V_h$$

$$a(u_h - u_I, v) = a(u - u_I, v) \quad \forall v \in V_h$$

Because theorem1 then

$$a(u - u_I, v) = O(h^2) \|u\|_3 |v|_1, \quad \forall v \in V_h$$

order $v = u_h - u_I \in V_h$

$$C \|u - u_I\|^2$$

$$\leq a(c, u_h - u_I)$$

$$\leq a(u - u_I, u_h - u_I)$$

$$= O(h^2) \|u\|_3 |u_h - u_I|_1, \quad \forall v \in V_h$$

Then

$$|u_h - u_I|_1 \leq Ch^2 \|u\|_3.$$

3.4 The postprocessing for the Finite Element Method

Theorem3 Assumption u_h and $u_I(x)$ the solution of discrete problem and the solution of the linear interpolation, and meet the superclose property theorem2,then

$$\|\Pi_{2h}^2 u_h - u\|_1 \leq Ch^2 \|u\|_3$$

proof: Through the variable type for $\Pi_{2h}^2 u_h - u$

$$\Pi_{2h}^2 u_h - u = (\Pi_{2h}^2 u_h - \Pi_{2h}^2 u_I) + (\Pi_{2h}^2 u_I - u)$$

Fist

$$\|\Pi_{2h}^2 u_h - \Pi_{2h}^2 u_I\|_1 \leq \|\Pi_{2h}^2 (u_h - u_I)\|_1$$

From Lemma2, $\Pi_{2h}^k u_I = \Pi_{2h}^k u$ so

$$\|\Pi_{2h}^2 u_h - \Pi_{2h}^2 u_I\|_1 \leq \|\Pi_{2h}^2 (u_h - u_I)\|_1 \leq C \|u_h - u_I\|_1$$

From theorem3

$$\|\Pi_{2h}^2 u_h - \Pi_{2h}^2 u_I\|_1 \leq \|\Pi_{2h}^2 (u_h - u_I)\|_1 \leq C \|u_h - u_I\|_1 \leq Ch^2 \|u\|_3$$

Second,from Lemma2

$$\|\Pi_{2h}^k u - u\|_1 \leq Ch^{r+1-l} \|u\|_r,$$

$$\|\Pi_{2h}^k v\|_1 \leq C \|v\|_r, \quad \forall v \in V_h, \text{then}$$

$$\|\Pi_{2h}^2 u_h - u\|_1 \leq \|\Pi_{2h}^2 u - u\|_1 \leq Ch^2 \|u\|_3.$$

3.5 Error estimation for $u - u_h$

Theorem4 Assumption $u(x)$ and $u_I(x)$ is the solution of variational problem and the solution of discrete problem,
 $\forall p \geq 2$ and $b > 0$,

$$|u - u_h|_1 \leq Ch^2 \|u\|_3$$

proof: From theorem1,3

$$|u - u_I| \leq O(h^2) \|u_h\|_3 \|v\|_1, \quad \forall v \in V_h$$

From theorem2and the triangle inequality

$$\begin{aligned} |u - u_h|_1 &= |u - u_I + u_I - u_h|_1 \\ &\leq |u - u_I|_1 + |u_I - u_h|_1 \leq C(h^2) \|u\|_3. \end{aligned}$$

4. Conclusion

Chuan-miao Chen and zhang xu, is presented in this paper on the basis of the numerical solution of one dimensional singular perturbation problems do the global superconvergence analysis, and got the expected results.

References

- [1] X Zhang , “A class of singular nonlinear two-point boundary value problems of error estimate of Galerkin solution,” Journal of computational mathematics, XXIII (4), pp. 195-205, 2010. (journal style)
- [2] D. Jespersen, “Ritz-Galerkin Methods for Singular Boundary Value prob lems,” Journal of Siam Journal on Numerical Analysis, pp. 813-834, 1978. (journal style)
- [3] K. Eriksson, V Thomée, “Galerkin methods for Singular Boundary Value Problems in One Space Dimension,” Journal of Mathematics of Computation, pp. 345-367, 1984. (journal style)
- [4] Chen, C. M, “Superconvergence of galerkin solutions for singular nonlinear two-point boundary value problems,” Journal of Mathematica Numerica Sinica, pp. 113-123, 1985. (journal style)
- [5] Wang, Xu, R. Caves, The mathematical basis of the finite element method, Science Press, 2004. (book style)
- [6] Lin, Yan, Efficient finite element structure and analysis, Efficient finite element structure and analysis, Hebei university press, 1996. (book style)
- [7] R G Dur á n, A L Lombardi, “Finite element approximation of convection diffusion problems using graded meshes,” Journal of Applied Numerical Mathematics, pp. 1314-1325, 2006. (journal style)
- [8] Lin, “Global error expansion and superconvergence for higher order interpolation of finite elements,” In Proceedings of Applied Numerical Mathematics, 1992. (journal style)
- [9] Q Lin, “A rectangle test for galerkin approximation to nonlinear hyperbolic problems,” Journal of the Shandong Normal University, 1997. (journal style)
- [10] Q Lin, “The finite element interpolated correction method for elliptic eigenvalue problems,” Chinese J.numer.math.appl, pp. 78-93, 1992. (journal style)
- [11] D Jespersen , “Ritz-Galerkin Methods for Singular Boundary Value Problems[J]. Siam Journal on

Numerical Analysis," Journal of Siam Journal on Numerical Analysis, pp. 813-834, 1978. (journal style)

[12] X Zhang, "A class of singular two-point boundary value problems of finite element method," Journal of Higher school, pp. 193-205, 2013. (journal style)

[13] M C Chen, "TGalerkin solution of singular nonlinear two-point boundary value problem of super convergence," Journal of computational mathematics, pp. 113-123, 1985. (journal style)