

Effect of Temperature and Time on Assessment of Methane Production from Microbial Degradation of Solid Waste in Port Harcourt, Rivers State, Nigeria

Mbata A. U. C¹, Nyenke C. U²

Medical Laboratory Science Department, Rivers State University of Science & Technology Nkpolu-Oroworukwo, Port Harcourt

Abstract: This research was to investigate the effect of temperature and time on emission of Methane from breakdown of solid waste by Methanogens. The assessment was carried out in three (3) dumpsites at Rumuokoro, Rumuosi and Eliozi, in the morning, afternoon and evening using lamcorn III which measured both atmospheric temperatures. At Rumuokoro dumpsite by 8.00hrs, 0.20ppm of Methane was emitted at 28°C, by 14.00hrs, 0.10ppm of Methane was emitted at 31°C and by 18.30hrs, 0.09ppm was emitted at 28°C. In Rumuosi dumpsite at 20°C 0.1ppm of Methane was emitted by 8.20hrs, 0.09ppm at 31°C by 13.40hrs and 0.09ppm at 28°C by 18.12hours. Similarly at Eliozi dumpsite by 10.30hrs, 0.09ppm was emitted at 30°C, 0.09ppm at 30°C by 13.00hours and 0.1ppm at 31°C by 17.40hours. There was no significant relationship between Methane emitted and atmospheric temperature ($p > 0.05$). There was also no significant relationship between the time and emission of Methane ($p > 0.05$). Consequently, Methane gas absorbs heat in the afternoon and release same in the morning and evening when the Methanogens would have degraded the waste overnight. Methane serves as alternative source of energy and production of ceramics if harnessed instead of acting as greenhouse gas which aids global warming.

Keywords: Methanogens, Degradation, Landfilling, Biogas, Decomposition.

1. Introduction

Microbial degradation of solid waste implies the breaking down of organic components of waste to inorganic form by Microorganisms which can readily serve as nutrient to variety of organisms and source of energy to humans. Waste can also be defined as materials which though may no longer be needed but may be used for other purposes (Oducha, 1994).

When waste is dumped on land, Methanogens readily colonize them and carry out degradation or transformation of degradable (organic) materials in the waste; provided the right environmental conditions are present. One major problem of urbanization is solid waste generation (Onibokun, 1989).

When anaerobic bacteria decompose waste, the end product is Methane (Morns, 1996). Waste generation started from the early man, who used natural forest resources as food and shelter and subsequently discard the remnant either as food remains or as human waste (Wachukwu and Eleanya, 2007). The rate of decomposition depends on the temperature, carbon/nitrogen ratio, PH, moisture, oxygen and chemical composition of the waste (Medina, 1997).

Methane is also called "swamp gas". It is a greenhouse gas formed as a result of decomposition of carbon containing substances found in oxygen free environments by Methanogens in landfills (Taralo and Taralo, 1999). Also Ruminating animals such as cattle, sheep and goat belch Methane into the air as a byproduct of digestion (Nester *et al.*, 2001). Methanogens produce Methane from the oxidation of hydrogen gas and the reduction of carbon dioxide as follows.



This is example of anaerobic respiration. Methane is lighter than air, colourless odorless and flammable. It has a melting point of 182°C and boiling point of 161.5°C. It is used as fuel in India to drive steam generators. Also used as raw materials in the manufacturing of chemicals, baking of bricks and ceramics tiles.

Methane is a primary constituent of landfill gas and a potent greenhouse gas when released to the atmosphere (Ackerman, 2000). Increase production of Methane contributes to greenhouse effect, with gradual global warming that could have dire results (EPA, 2008).

Globally, landfills are the third, largest human induced emission source of Methane accounting for about 12% of global Methane emission (Oyegun and Ologunorisa, 2002).

According to an environmental protection agency report, (2008) an estimated global landfill Methane emission is put at 747.38 million metric tons of carbon dioxide equivalent. The major factors driving landfill gas emission are the amount of organic material deposited in landfills, type of landfill practices, extent of anaerobic decomposition and the kind of landfill Methane recovery and decomposition (Ayoade, 2003).

The study was carried out to ascertain the level of Methane gas emitted from the waste dump as a result of microbial degradation by Methanogens and also to ascertain if temperature and time have effect on the emission of Methane gas.

2. Materials and Methods

2.1 Study Area

The research was carried out in the city of Port Harcourt, Rivers State of Nigeria. The occupants are people from

different ethnic group and beyond and they include civil servants, farmers, fishermen, traders, oil and gas workers and expatriates from other countries. Port Harcourt has an international airport, seaport and a lot of migrant workers from other parts of the world. The city has quite a number of markets and sundry businesses and as such there is beehive of commercial and industrial activities which lead to an abundant generation of waste.

2.2 Study Sites

The study was carried out in seven waste dumpsites which were within the metropolis, the include Rumuokorodumpsite, Elioizu dumpsite and Rumuosi dumpsite.

2.3 Analysis

A total of 3 major dumpsites situated at Rumokoro, Rumuosi and Elioizu were visited in the morning, afternoon and evening hours at different times. The Methane gas and the atmospheric temperature were estimated at different times using Lamcam III gas analyzer which was suspended into the atmosphere or point source for about 10 minutes. The figures were displayed on the automated LCD screen of the equipment and recorded.

3. Result

The Methane emission at Rumuokoro, Rumuosi and Elioizu according to temperature and time is shown. There was no significant relationship between Methane production and atmospheric temperature ($p > 0.05$). As shown in table 1.

The time also has no significant effect on Methaneemission $p > 0.05$ as shown in table 2.

Figure 1 shows the graphical representation of Methane gas emitted in comparison to time and temperature. This shows no significant difference in emission of Methane in relation to time and temperature.

Table 1: Methane gas emission according to temperature and time at different locations

S/N	Location	Methane Result in ppm (%)	Temperature in °C (%)	Time in hours (%)
1	Rumuokoro	.20(21)	28.00(10.50)	8.00(6.60)
2	Rumuokoro	.10(10.50)	31.00(11.65)	14.00(11.60)
3	Rumuokoro	.09(9.50)	28.00(10.5)	18.30(15.15)
4	Rumuosi	.10(10.50)	29.00(10.90)	8.20(6.80)
5	Rumuosi	.09(9.50)	31.00(11.65)	13.40(11.10)
6	Rumuosi	.09(9.50)	28.00(10.50)	18.12(15.00)
7	Elioizu	.09(9.50)	30.00(11.30)	10.30(8.50)
8	Elioizu	.09(9.50)	30.00(11.30)	13.00(10.80)
9	Elioizu	.10(10.50)	31.00(11.65)	17.40(14.4)
	TOTAL	.95(100)	266.00(100)	120.72(100)

Table 2: Methane gas emission according to time of the day and dumpsite

Period of the day	Rumuokoro (o) (e)	Rumosi (o) (e)	Elioizu (o) (e)	Total	χ^2	P.value (0.05)	Comment
Morning Hours	0.2(0.16)	0.1(0.15)	0.09(0.115)	0.39	0.017	<0.05	Not Significant
Afternoon Hours	0.1(0.115)	0.09(0.083)	0.09(0.083)	0.28	0.003	<0.05	Not Significant
Evening Hours	0.09(0.115)	0.09(0.083)	0.10(0.082)	0.28	0.01	<0.05	Not Significant
TOTAL	0.39	0.28	0.28	0.95	0.04	<0.05	Not Significant




FIG. 4.7 MEAN VALUE OF METHANE ACCORDING TO TEMPERATURE AND TIME

4. Discussion

From the result obtained it was shown that Methane gas had the highest level in the morning hours, 0.20ppm, 0.1ppm and 0.09ppm than in the evening and afternoon. The reason being that Methane traps and absorbs more heat from the atmosphere during the day and starts emitting them by degradation of waste by Methanogens in the morning.

Therefore there was no significant relationship between the emission of Methane and temperature ($p > 0.05$). Also there was no significant relationship between the time and production of Methane ($p > 0.05$). The normal range of Methane is 70 parts per million (ppm). The low value obtained could be due to the fact that the waste are brought to the dumpsites every time and thus fresh refuse are used to cover the old ones when degradation could be taking place actively. Moreover most refuse dumps are fresh.

Since Methane gas production is increasing and as such the disruption of temperature balance is imminent as reported by Ackemann (2000), it becomes necessary to extract Methane gas for other purposes as fuel for steam generators as done in India and production of ceramics.

5. Conclusion

Methane is a primary constituent of landfill gas and a potent greenhouse gas when released to the atmosphere, excess Methane emitted can also be diverted to other meaningful areas.

Better management of our waste can significantly reduce emissions of greenhouse gas to the atmosphere. We can also turn waste materials into resources to be valued and also reduce the amount of waste we produce and the amount of energy we use.

References

- [1] Oducha, J.N.I.G (1994). Waste generation and management in a depressed economy. *A lecture delivered to students of law and environmental faculties in UNEC.*
- [2] Onibokun, A.G. (1989). Urban growth and management in Nigerian, in: African cities managing rapid urban growth. Westview press builder, p, 6911
- [3] Morns, F. (1996). Recycling versus incineration an energy conservation analysis. *Journal of hazardous material*, **47**,227-293
- [4] Wachukwu, C. K and Eleanya, E.U. (2007). Health impact assessment of solid waste disposal workers in Port-Harcourt Nigeria. *Journal of Applied science* **7**(22), 3562-3566.
- [5] Medina, M. (1997). Informal Recycling and collection of solid waste in developing countries, issues and opportunities. *Working paper 24 of the united nation universities institute of advance studies*, Tokyo, japan p 20-30
- [6] Taralo, K and Taralo, A. (1996). Atmospheric cycles, *Foundations in microbiology* WCB publishers, London, p 820-821

- [7] Nester, E.N., Anderson, D.G., Roberts, C.E., Persal, N.W, and Nester, M.T. (2001). Microbiology of solid waste, *Microbiology-A human perspective* vol. **31** McGraw Hill Boston, p 798-802
- [8] Ackerman; F (2000). Waste management and climate change local. *Environment* **5**,223-229
- [9] Environmental protection Agency, (2008). Global anthropogenic emission of non-CO₂ Greenhouse Gases, 1990-2020. EPA Report **6**, 24
- [10] Ayoade, J.O (2003). Mechanical *Biological pretreatment of msw Bioprocessing of solid waste and sludge*, **2**, 23-36