

Telomere Length a Biomarker for Ageing

Mritunjay Kumar Singh¹, Taruna Gupta²

¹Research Associate Allele Life Sciences Pvt. Ltd. Noida,U.P.

²Research Trainee, Allele Life Science Pvt. Ltd. Noida, U.P.

Abstract: *Telomere length in human is emerging as a biomarker of ageing because its shortening is associated with age-related disease. In this review we summarize our current knowledge on the role of telomeres shortening in humans by comparative analysis of TL by using blood samples of different person with different age groups, smokers and non-smokers & sports and non-sports of same ages with the help of different techniques conventional PCR and RT-PCR.*

Keywords: Telomere, ageing, senescence, TL (Telomere Length), RT-PCR (Real Time-Polymerase Chain Reaction)

1. Introduction

Ageing is the process of becoming older. In broader sense, ageing can refer to single cell within an organism. In human, ageing represents the accumulation of changes in a human being over time, encompassing physical, psychological, and social change. Ageing is among the largest known risk factors for most human diseases. The rate of ageing varies substantially across different species, and this, to a large extent, is genetically based. Every nucleus of a eukaryotic cell is packed with chromosomes, the long, thread-like DNA molecules that carry the genes of living organisms. In the end of the chromosomes there is a region of repetitive nucleoprotein structure called telomeres, which prevent chromosomal ends from end to end fusion and degradation.

Telomeres consist of stretches of repetitive DNA with high guanine and cytosine content. In humans, the telomere terminus consists of 4-15 kbp (bp = base pairs) of the hexanucleotides 5'-TTAGGG-3'. The G-rich DNA strand runs 5' to 3' toward the terminus and protrudes 100-150 nucleotides beyond the complementary C-rich strand.

Telomere shortening rates can be modified by several lifestyle factors that include mental stress levels, cigarette smoking and dietary intake. Baseline telomere shortening due to the end-replication problem will always exist, but efforts to live a healthy and balanced lifestyle can reduce any additional loss of telomeres. Even poor sleep quality has been associated with shorter telomere lengths in leukocytes among midlife women, independent of perceived psychological stress. There are several factors that causes telomere shortening in a cell. The two most common causes are due to the end-replication problem, and oxidative damage to DNA.

2. Material and Methods

1) Sample Collection, Storage

Blood samples of Smoker & Non-Smoker Person; Sports & Non-sports person, Normal person with different age groups varying from 13 to 55 yrs. Most in vivo studies of TL in vertebrates' uses blood, for which small samples can be taken routinely and repeatedly from the same individuals, usually with little adverse effect. Importantly, blood cells generally have a high turnover rate

Leading to the expectation and widespread observation of TL loss in blood cells with age and over time. However, blood cells comprise a rich composite of cell types with very different functions, proliferation patterns and levels of telomerase expression.

2) Genomic DNA Isolation from Blood Samples

Preparation of Chemicals

a) Saturated Phenol

Materials Required:

- Redistilled Phenol, molecular biology grade. (Stored in aliquots at -20°C).
- 8-hydroxyquinoline
- 0.5 M Tris.Cl buffer (pH 8.0)
- 0.1 M Tris.Cl buffer (pH 8.0)

Note: Phenol is volatile and caustic. .

Protocol for preparation of Saturated Phenol:

- Heat a water bath to 65°C in the fume hood. Place the bottle of phenol in the fume hood to warm to room temperature.
- Place the bottle of phenol in the 65°C water bath to melt the crystals.
- Add 8-hydroxyquinoline to a final concentration of 0.1 % w/v to the phenol. Mix to dissolve the 8-hydroxyquinoline.
- Add an equal volume of 0.5 M Tris. Cl (pH 8.0) to the phenol. Mix for 15 minutes. Return the bottle to the 65°C water bath. Allow the phases to separate. Siphon off the top layer and discard. Repeat the procedure as in Step 4 twice
- Add an equal volume of 0.1 M Tris. Cl (pH 8.0) to the phenol. Repeat the procedure as in Step 4.
- Repeat the extractions with 0.1 M Tris. Cl (pH 8.0) until the aqueous phase is ~pH 7.8 (measure with pH paper). Repeat the procedure as in Step 4. Leave approx. ~1 cm layer of 0.1 M Tris.Cl (pH 8.0) over the phenol. Add 2-mercaptoethanol to a final concentration of 0.2% w/v to the 0.1 M Tris.Cl (pH 8.0).
- The buffer saturated phenol may be stored at 4°C for 1 month for DNA extraction. Test the pH periodically and do not use if the pH is <7.5.

b) Lysis Buffer

- 50M Tris.cl
- 20 mM EDTA
- 1% SDS
- 10 µL β-merceptoethanol

c) TE Buffer

- 10mM Tris
- 1 mM EDTA

Protocol for Genomic DNA Isolation:

- Approximately 250µl of previously separated blood cells were taken in a 700µl of lysis buffer. The mixture was incubated at 55°C overnight for digestion. Incubated for 37°C for 6 hrs with occasion are mixing by gentle swirling in water bath.
- Dissolve solution was centrifuge to separate cell debris at 10000 rpm for 10 minutes. Aqueous phase was transfer to another tube.
- Equal volume of s-phenol was added to precipitate the protein and centrifuge at 10,000rpm for 10 minutes. Aqueous phase was taken and transfer to another tube, without disturbing protein complex precipitated as an interphase.
- Equal volume of s-phenol: Chloroform (1:1) was added and centrifuged at 10,000 rpm for 10 minutes to ensure emulsification of the phases. Aqueous phase was transfer to another tube.
- Equal volume of chilled chloroform were added and centrifuged at 10,000 rpm for 10minutes. Aqueous phase was taken and transfer to another tube.
- 150-170µl of 5M NaCl was added to the mixture and double the volume of chilled absolute isopropanol and is allowed to stand for incubation at -20°C for 45 minutes.
- DNA pellete was collected by centrifugation at 10,000 rpm for 10 minutes.
- The tube were inverted and drained on a paper towel. The pellete was washed with 500 µl of 70% ethyl alcohol and centrifuged at 10,000rpm for 10 minutes. Discard the supernatant and air dried the pellete.
- The pellete was resolved in 50µl of TE buffer.

d) Real Time PCR

1st step:

- RNA 5 ul
- ddH2O 15 ul
- Incubate at 45°C for 600 C
- Incubate at 95°C for 5 minutes.
- Dilute it with 30ul Nuclease free water.

2nd Step:

RT PCR ABI7000

Temperature	Time
95°C	5 minutes
Loop[35 cycles
95°C	30 seconds
55°C	1 minute
72°C	30 seconds
Loop]	
72°C	5 minutes

Primers used for RT-PCR

36b4 gene	
Forward Primer	5'- TCGAACACCTGCTGGATGAC -3'
Reverse Primer	5'- CCACGCTGCTGAACATGCT -3'

Cocktail for RT -PCR


Components	Quantity (micro-litre)
Sybr Green	12.5
Template cDNA	1.0
Primer forward	1.0
Primer reverse	1.0
Double distilled water	9.5

Real time thermal cycler program

Temperature	Time
95°C	10 minutes
Loop[40 cycles
95°C	15 seconds
60°C	1 minute
Loop]	

3. Result and Discussion

Qualitative analysis of DNA:



Sample DNA (Ageing)		Sample DNA (Sports and Non-Sports)	
L1	AG1 (19 years)	L6	SP1 (25 years)
L2	AG2 (23 years)	L7	NSP1 (25 years)
L3	AG3 (33 years)	L8	SP2 (35 years)
L4	AG4 (46 years)	L9	NSP2 (35 years)
L5	AG5 (56 years)	L10	SP3 (45 years)
		L11	NSP3 (45 years)

Sample DNA (Smokers and Non-Smokers)	
L12	SM1 (27 years)
L13	NSM1 (27 years)
L14	SM2 (38 years)
L15	NSM2 (38 years)
L16	SM3 (47 years)
L17	NSM3 (47 years)

*AG: Ageing blood sample

*SP, NSP: Sports person and Non-sports person respectively

*SM, NSM: Smokers and Non-smoker blood sample

Quantitative analysis of DNA:

Sample	Abs @ 260	Abs @ 280	Ratio (OD)	Yield of DNA (ug/ul)
AG1 (19 years)	0.396	0.246	1.60	1.72
AG2 (23 years)	0.325	0.192	1.69	1.76
AG3 (33 years)	0.252	0.148	1.70	1.26
AG4 (46 years)	0.286	0.166	1.72	1.62
AG5 (56 years)	0.185	0.111	1.66	1.44
SP1 (25 years)	0.121	0.078	1.55	1.72
NSP1 (25 years)	0.194	0.121	1.60	1.76
SP2 (35 years)	0.235	0.131	1.79	1.26
NSP2 (35 years)	0.298	0.185	1.61	1.62
SP3 (45 years)	0.321	0.179	1.79	1.44
NSP3 (45 years)	0.285	0.158	1.80	1.44
SM1 (27 years)	0.245	0.147	1.66	1.72
NSM1 (27 years)	0.421	0.245	1.71	1.76
SM2 (38 years)	0.325	0.198	1.64	1.26
NSM2 (38 years)	0.215	0.121	1.77	1.62
SM3 (47 years)	0.315	0.196	1.60	1.44
NSM3 (47 years)	0.184	0.105	1.80	1.44

Real Time PCR analysis of Tel1b and Tel2b genes.

Standard curve for 36B4 (Reference gene)

36B4 gene is used as single copy gene. SCG (single copy gene) is used as control for amplification of each sample. 36B4 gene encodes acidic ribosomal phosphoprotein P0. For standard curve of 36B4 gene DNA was diluted in different concentration and the reaction was run in real time PCR generating different Ct value for each dilution. Ct value increases with decrease in DNA concentration. Concentration of synthesized Oligomer of 36B4 was.....which was diluted to 1 ng for first dilution and it was further diluted to 2X, 4X, 8X and 16X for standard curve analysis.

Standard curve of 36B4 showed a slope of -3.34 and PCR efficiency of 99.08%.




Figure: 36B4 oligomer standard

On the x-axis the logarithm of the genome copy number is displayed, on the y-axis the corresponding Ct values. The trend line equation is used for calculating 36B4 genome copy number for telomere Ct values.

Primer sequence for real time PCR

4. Conclusion

Although relying on earlier experiment we performed direct conventional PCR to check comparative length of telomere, we fail to do so.

In conventional PCR, which DNA sequence will be targeted is a matter of concern. From various literature we have come to know that 36b4 gene can be targeted to design primer.

Data collected from Real time PCR suggest variations but only in first case i.e. different age group.

Although variation is observed in rest two cases also, data interpretation is not quite satisfactory.

Therefore, we conclude telomere can be used as for biomarker in ageing.

References

- [1] Aviv, A., Hunt, S.C., Lin, J., Cao, X., Kimura, M., and Blackburn, E. (2011). Impartial comparative analysis of measurement of leukocyte telomere length/DNA content by qPCR. *Nucleic Acids Res* 39, e134.
- [2] Haussmann, M.F. & Mauck, R.A. (2008) new strategies for telomere-based age estimation. *Molecular Ecology Resources*, 8,264–274.
- [3] Pfaffl, M.W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, 29, e45.
- [4] Zee, R.Y., Castonguay, A.J., Barton, N.S., Germer, S., and Martin, M. (2010). Mean leukocyte telomere length shortening: a case-control study. *Transl Res* 155, 166-169.
- [5] O'Callaghan, N.J. & Fenech, M. (2011). A quantitative PCR method for measuring absolute telomere length. *Biological Procedures Online*, 13:3