

Pre-Prosthetic Preparation in the Treatment with Fixed Prosthetic Constructions

Desislava Konstantinova, Anna Nenova-Nogalcheva

Abstract: The authors' aim was to determine the role of pre-prosthetic periodontal preparation in the life expectancy of dental bridges. **Materials and methods:** A retrograde screening study was performed on 58 patients with bridge restorations, subsequently divided into two groups: Group 1 involved patients who had undergone pre-prosthetic periodontal preparation and Group 2 where patients had not had any pre-prosthetic periodontal preparation. Inclusion criteria were subjective complaints, life span of the bridge, evaluated according to the three medico-biological criteria and according to the periodontal pocket depth. Changes were followed up in four successive examinations: after 1 week, 1 year, 2 years and 5 years. **Results:** The results pointed towards a significant correlation between participants' affiliation to a particular group (Group 1 or Group 2) and the progression of the periodontal pocket depth ($P=0.20$). A linear correlation was established between the number of patients in Group 2 and any discrepancies in the preventive ($R=0.792$, $P < 0.01$), functional ($R=0.742$, $P < 0.01$) and aesthetic criteria ($R=0.881$, $P < 0.01$). At the first control examination, Group 2 patients had subjective complaints which over time (after 3 years and 5 years) intensified. Such observations were not found among Group 1 participants. **Conclusion:** The longevity of dental bridge prostheses is contingent on the state of the hard and soft tissue of the oral cavity. Pre-prosthetic periodontal preparation is essential in achieving long-term functionality, lack of subjective complaints and good aesthetics.

Keywords: pre-prosthetic preparation, fixed constructions

1. Introduction

Periodontitis is defined as inflammatory destructive lesions induced by infectious agents [1, 2]. Shortly after fitting the prosthesis there is an early bacterial colonization and biofilm formation occurring in the prosthetic construction [3]. There are studies supporting that areas affected by periodontal infection may serve as a reservoir for pathogenic microorganisms that can pass from infected areas to healthy tissue, multiply and cause disease [4]. Hence, pre-prosthetic preparation is fundamental for the successful prosthetic treatment of the masticatory apparatus using fixed prosthetic devices [5, 6]. In modern dentistry, that can be easily achieved when an interdisciplinary approach is applied [7, 8].

2. Aim

To determine the role of pre-prosthetic periodontal preparation in the longevity of fixed dental constructions for the rehabilitation of the masticatory apparatus.

3. Material and Methodology

A retrograde screening study was performed on patients with bridge restorations, referred to specialized prosthetic and periodontal treatment by their general practitioners. The target of the study were 88 prosthetic devices (48 in the maxilla and 40 in the mandible), fixed onto 489 abutment teeth. (Table 1)

Table 1: Frequency distribution of abutment teeth

Type of dental construction	Maxillary dental bridges	Mandibular dental bridges	Total number of abutment teeth
3-unit dental bridge with 2 abutment teeth	3	10	26
4-unit dental bridge with 4 abutment teeth	5	10	45
5-unit dental bridge with 4 abutment teeth	10	8	72
7- or 8-unit dental bridge with 6 abutment teeth	14	7	136
Cross-arch stabilizing bridge with 10 abutment teeth	16	5	210
Total	305	184	489

58 patients signed an Informed Consent for the purposes of the study, divided into two groups:

- Group 1 comprised 28 patients (15 women and 13 men, mean age of 55 ± 8 years), who sought dental assistance in complete aesthetic rehabilitation of the masticatory apparatus using fixed bridge constructions. The patients consented to a pre-prosthetic periodontal and surgical preparation of the prosthetic field.
- Group 2 consisted of 30 patients (16 women and 14 men, mean age of 58 ± 7 years), who underwent prosthetic treatment without prior preparation of the prosthetics field.

All patients were selected based on the following inclusion criteria, suggested by the authors of the present paper: lack of common diseases, indications for prosthetic rehabilitation using fixed bridge constructions and presence of intraosseal bone defects measuring more than 5 mm in depth. The study included medical history taking, extra- and intraoral examinations, panoramic imaging and documentation of the patient's periodontal status on a specially developed Dental Diagnostic Card. (Figure 1)

Dental Diagnostic Card

Patient's name : Age:
 Telephone:

Dental Status

			55	54	53	52	51	61	62	63	64	65				
18	17	16	15	14	13	12	11	21	22	23	24	25	26	27	28	
48	47	46	45	44	43	42	41	31	32	33	34	35	36	37	38	
			85	84	83	82	81	71	72	73	74	75				

C - Caries P - Pulpitis G - Periodontitis R - Root F - fracture O - Obturation
 E - Missing K - Crown X - Artificial

Gingival Inflammation Index - Papilla Bleeding Index

I PalatinaI no	II Vestibularno	0-0.5 - healthy periodontium
		0.5-1 - mild gingivitis
IV Vestibularno	III P PalatinaI no	1-2 - moderate gingivitis
		3-4 - severe gingivitis

Oral hygiene index by Greene and Vermillion (DI): debris (plaque)

16	14	11	21	24	26	0 - No debris or stain present 1 - Soft debris covering not more than 1/3 of the tooth surface 2 - Not more than 2/3 of the exposed tooth surface. 3 - More than 2/3.
46	44	41	31	34	36	

Figure 1: Dental Diagnostic Card

The pre-prosthetic preparation for Group 1 involved open flap curettage in the following sequence: dissection of the vestibular and oral mucoperiosteal flap, performing a crevicular vestibular and lingual incision as well as vertical incisions beyond the mucogingival line, which provided visibility and access to the base of the periodontal pockets. Calculus (tartar) and unhealthy granulation were removed from the bottom of the periodontal pockets, followed by smoothing and polishing of teeth root surfaces. Necessary tooth extractions were carried out. After gentle and careful hemostasis the flaps were adjusted and sutured with a 4-0 silk suture. An interim construction was fitted and a periodontal dressing was inserted for 7 days. (Figure 2)



Figure 2: Patient M.Z. (62 years old) – an intraoral view after placing a periodontal dressing

Surgical dental crown lengthening (elongatio corone dentis) proved necessary to perform, using gingivectomy or apically positioned flap surgery (APF). Depending on the size of the recession, gingival recessions affecting one or several teeth were treated by covering the exposed teeth roots or by microsurgical periodontal intervention using palate mucosa autotransplantation. Bone-grafting technique was applied with some of the patients in order to increase the bone level.

The size and type of bone-grafting granules were selected in view of each patient's individual characteristics.

Group 1 patients had their final fixed prosthesis fitted only upon full recovery of the periodontium after 5 ± 1 months. Group 2 patients had their prostheses mounted immediately after removal of the previous construction and following initial therapy. Key evaluation criteria included subjective complaints, such as cold sensitivity, spontaneous gum bleeding and bad breath, as well as the dental bridge life span, evaluated according to the three medico-biological criteria (preventive, functional and aesthetic criteria). The periodontal pocket depth around the abutment teeth was also assessed. All subjective complaints, the state of the prosthesis and the periodontium were checked and analyzed in 4 follow-up checks – after 1 week, 1 year, 2 years and 5 years. The data were analyzed using SPSS software for epidemiological and clinical research (Windows, V 16.0.0, Nov. 2007). The following statistical methods were applied: frequency and percentage distribution of data, graphical representation of data and Pearson's correlation coefficient, etc.

4. Results and Discussion

The data indicated that 82.14% of all participants had previous prosthetic treatment 5 or more years before that, which did not meet the basic medico-biological criteria. (Figure 3)



Figure 3: Patient M.B. (58 years old) – an intraoral view prior to treatment



Figure 4: Patient K.K. (57 years old) – an intraoral view prior to treatment



Figure 5: Patient E.M. (48 years old) – a panoramic image prior to treatment



Figure 7: Patient S.S. (55 years old) – an intraoral view prior to treatment

17.86% of participants did not have any prosthetic devices but had missing teeth or intraosseal bone defects of varying depth. (Figures 6 and 7).



Figure 6: Patient L.P. (48 years old) – an intraoral view prior to treatment

At the initial examination 100% of patients had symptoms of bleeding gums, discomfort when chewing and increased sensitivity to cold.

The results demonstrated that 100% of Group 1 patients, who received prior periodontal and surgical treatment, showed initial periodontal pocket depth measuring 5 ± 2 mm, later observed to be reduced to 2 mm as early as the first control examination before the prosthetic treatment. The same results were recorded 1 week after the prosthesis was fitted, and were retained in 75% of patients after 1 year. (Figure 8)

Periodontal pocket depth (in mm)




Figure 8: A graphical representation of the distribution of Group 1 participants according to the periodontal pocket depth (in mm)

The observations from the control examination showed that 1 week after the treatment, no patients in Group 2, who underwent pre-prosthetic preparation, had periodontal pocket depth of 2 mm, and only 16.67% of participants in this group had pocket depth measuring 2 ± 1 mm. (Figure 8) The results observed at later examinations (after 3 years and 5 years) indicated that 71.43% of Group 1 patients did not have any change in the periodontal pocket depth, whereas in the majority of Group 2 patients (76.67%) the pocket depth measured 3 ± 1 mm, which could potentially compromise the bridge construction. 5 years after the treatment all patients in Group 2 showed indications for new prosthetic rehabilitation, as opposed to only 7.14% in Group 1. There was significant correlation between participants' affiliation to a particular group (Group 1 or Group 2) and the progression of the depth of the periodontal pockets ($P=0.20$).

The data obtained from the examinations of the prosthetic devices after 3 years revealed that according to the three basic medico-biological criteria there were no deviations from the norm in Group 1. After 5 years one patient showed discrepancy in the functional criterion, so did two patients in the preventive criterion, while four patients demonstrated discrepancy in the aesthetic criterion, which could be due to external factors (e.g. smoking, unsatisfactory oral hygiene, etc.). The observations showed that after 3 years in 50% of the patients in Group 2 there were discrepancies in the basic medico-biological criteria. Moreover, after 5 years the results were more alarming and there were indications for the need of new prosthetic rehabilitation. (Table 2) The linear correlation between the number of patients in Group 2 and the discrepancies in the preventive ($R=0.792$, $P<0.01$), functional ($R=0.742$, $P<0.01$) and aesthetic criteria ($R=0.881$,

P<0.01) was calculated using Pearson's correlation coefficient.

Table 2: Percentage distribution in groups according to the three basic medico-biological criteria

Medico-biological Criteria/ Assessment	Group 1 patients – after 7 days	Group 2 patients – after 7 days	Group 1 patients – after 1 year	Group 2 patients – after 1 year	Group 1 patients – after 3 years	Group 2 patients – after 3 years	Group 1 patients – after 5 years	Group 2 patients – after 5 years
Compliance with the Preventive Criterion	100%	100%	100%	86.67%	100%	50%	92.86%	10%
Discrepancy in the Preventive Criterion	0%	0%	0%	13.33%	0%	50%	7.14%	90%
Compliance with the Functional Criterion	100%	100%	100%	86.67%	100%	46.67%	96.43%	10%
Discrepancy in the Functional Criterion	0%	0%	0%	13.33%	0%	53.33%	3.57%	90%
Compliance with the Aesthetic Criterion	100%	96.67%	100%	86.67%	92.86%	40%	85.71%	33.33%
Discrepancy in the Aesthetic Criterion	0%	3.33%	0%	13.33%	7.14%	60%	14.29%	66.67%

Discrepancy in any of the three basic medico-biological criteria is considered an indication for a new prosthetic treatment plan [8]. The permanence of prosthetic

constructions is often determined by the subjective perception of patients. (Table 3)

Table 3: Percentage distribution in groups according to subjective complaints

Subjective complaints	Group 1 patients – after 7 days	Group 2 patients – after 7 days	Group 1 patients – after 1 year	Group 2 patients – after 1 year	Group 1 patients – after 3 years	Group 2 patients – after 3 years	Group 1 patients – after 5 years	Group 2 patients – after 5 years
Sensitivity to cold	0%	0%	0%	26.67%	3.57%	46.67%	3.57%	46.67%
No sensitivity to cold	100%	100%	100%	73.33%	96.43%	53.33%	96.43%	53.33%
Spontaneous gum bleeding	0%	3.33%	0%	40%	0.00%	53.33%	3.57%	60%
No spontaneous gum bleeding	100%	96.67%	100%	60%	100%	46.67%	96.43%	40%
Bad breath	0%	0.00%	0%	33.33%	0.00%	50%	3.57%	60%
No bad breath	100%	100%	100%	67.67%	100%	50%	96.43%	40%

The observations showed that Group 2 patients had subjective complaints as early as the first check-up, which intensified over time (after 3 and 5 years). Such complaints were not observed among Group 1 patients. The results only emphasized the difference between contaminated periodontal prosthetic field and a prosthetic field following periodontal treatment. (Figures 9 and 10)



Figure 9: Patient M.Z. (62 years old) – an intraoral view 5 years after periodontal and prosthetic treatment



Figure 10: Patient S.S. (61 years old) - an intraoral view 5 years after prosthetic treatment

5. Discussion

The proper planning of prosthetic treatment with fixed dental constructions demands a thorough study and assessment of the anatomical characteristics and the clinical condition of both hard and soft tissues [9]. Biological and biophysical aspects play a huge role in the outcome of the prosthetic treatment [6, 10]. Negligence of the state of the periodontal tissues, especially in the cases of upcoming prosthetic treatment, can dramatically affect the treatment success and its prognosis bona [6, 11]. Discrepancy in any the basic medico-biological criteria is considered an indication for new prosthetic rehabilitation [8, 12].

6. Conclusions

The longevity of dental bridge prostheses is contingent on the state of the hard and soft tissue of the oral cavity. Pre-prosthetic periodontal treatment is essential for the purpose of long term functionality, lack of subjective complaints and good aesthetics of the treatment with fixed bridge constructions. Overlooking the need for pre-prosthetic periodontal preparation, where there are indications for it, can only lead to short-term benefits of the prosthetic treatment. Best long-term prognosis of the treatment with fixed prostheses can be achieved through an interdisciplinary approach.

References

- [1] Liljenberg B, Gualini F, Berglundh T, Tonetti M, Lindhe J. Composition of plaque-associated lesions in the gingiva and the peri-implant mucosa in partially edentulous subjects. *J Clin Periodontol.* 1997; 24: 119-123.
- [2] Van Winkelhof AJ, Goene RJ, Benschop C, Folmer T. Early colonization of dental implants by putative periodontal pathogens in partially edentulous patients. *Clin Oral Implants Res.* 2000; 11:511-552
- [3] Manev G, Popova C. Clinical diagnosis of inflammatory / destructive peri-implant diseases. *Problems of dental medicine.* 2013;1:39.
- [4] Passariello C, Alessandra MP, Gigola VP. Microbiological and host factors are involved in promoting the periodontal failure of metaloceramic crowns. *Clinical Oral Investigations .* 2012; 16.(3): 987-995.
- [5] Ephros H, Klein R, Sallustio A. Preprosthetic Surgery. *Oral Maxillofacial Surg Clin.* 2015; 27: 459–472.
- [6] Jansson L, Ehnevid H et al. Proximal restorations and periodontal status. *J Clin Periodontol.* 1994; 21:577-82

- [7] Lindhe J, Nyman S, Lang NP. Treatment planning. In Lindhe J, Karring T, Lang N P (Ed). *Clinical periodontology and implant dentistry.* Oxford: Blackwell Munksgaard . 2003;4(19): 414–431.
- [8] Filtchev A, Ralev R. *Propedeutics of prosthetic dental medicine.* Sofia. 2010: 213
- [9] Yung-Ting H, Nan-Chieh H, Hom-Lay W. Relationship between periodontics and prosthodontics: The two-way street. *Journal of Prosthodontics and Implantology.* 2015; 4: 4 -11
- [10] Gašperšič R, Petelin M, Kopač I, Marion L, Skalerič U. The relationship between periodontal tissues and the gingival margin of prosthetic restorations. *Zobozdrav Vestn.* 2005; 60: 182- 192
- [11] Uhač I, Kuiš D, Kavčič R, Lajnert V, Šoškić MŠ, Antonić R et al. Fixed prosthodontic restorations and periodontal health. *Medicina Fluminensis.* 2014; 50(3): 279-287.
- [12] Lang N. Periodontal consideration in prosthetic dentistry. *Periodontology 2000.* 1995; 9: 118-131.

Author Profile



Dr. Desislava Konstantinova obtained her Master's degree in Dentistry in 1997 from the Medical University –Plovdiv. She has been working as an Assistant Professor at the Prosthetic Dentistry Department at the Medical University –Varna since 2008. She specialized in Prosthetic Dentistry in 2106 and was awarded a PhD degree in 2016.



Dr. Anna Nenova-Nogalcheva obtained her Master's degree in Dentistry in 2006 from the Medical University – Plovdiv, Bulgaria. She has been working as an Assistant Professor at the Oral Maxillofacial Surgery Department at the Medical University –Varna since 2009. She specialized in Oral Surgery in 2012.