

Study of Reproductive Behavior of Adult Female (Crocodile) in Kotmi Sonar of Janjgir Champa (C.G.)

Kavita Gabel¹, Manoj Gabel²

¹M. Phil. Scholar, Dr. C.V. Raman University Bilaspur (C.G)

²Assistant Professor, Dr. C.V. Raman University Bilaspur (C.G)

Abstract: Crocodile has been categorized as a vulnerable species in the Red List of IUCN and is placed under schedule 1 of the Wild Life Protection Act, 1972. To protect Crocodiles a Crocodile Conservation Park has been established by The Govt. of Chhattisgarh in Munda pond of Kotmi Sonar, District Janjgir - Champa, (C. G.) India. The physicochemical conditions of Munda pond provide an optimal conditions for, growth and vital activities for this species, *Crocodylus palustris*. The population of *Crocodylus palustris* is gradually increasing up to 378. The establishment of incubation centre, artificial hatchery and other technical facilities helps in the increasing population of *Crocodylus palustris*. The potentialities of Crocodile Park showed explored from point of view of knowledge, research activities and ecotourism as well.

Keywords: Conservation, Eco tourism, vulnerable species.

1. Introduction

The Marsh Crocodile, Mugger or *Crocodylus palustris* is a common and widespread crocodylian species in India [Vyas, 2012]. *Crocodylus palustris* is ecto and heterothermic, seasonal breeder, gonochoric, dioecious, hole-nesting and oviparous species [Jacobson, 1999; Da Silva and Lenin, 2010]. The population of *Crocodylus palustris* is reportedly declining [Whitaker and Andrews, 2003; Oza, 1975]. The population is reportedly declining due to illegal hunting for skin and indigenous medical purpose, habitat destruction. Lack of appropriate 'rescue and release' protocols and man made causes are another causes for shrinking population of *Crocodylus palustris* [Vyas, 2012; Da Silva and Lenin. 2010; Whitaker and Whitaker, 1984; Joshi et al., 2011; Vyas and Vyas, 2002; Vyas, 2010]. Their global population is tentatively estimated as 5400 to 7100 non-hatching [Da Silva and Lenin, 2010]. To protect and conserve this endangered species, it has been categorised as a vulnerable species in the Red List of IUCN and Red List of threatened Reptilian species. It is placed under schedule 1 of Wild Life Protection Act, 1972 [Joshi et al., 2011; Choudhury and Chowdhury, 1986; Bharos and Kanoje, 2007]. To provide a higher degree of protection to *Crocodylus palustris*, a Crocodile Conservation Park is established in Munda pond at Kotmi Sonar of Janjgir-Champa District of Chhattisgarh, India [Fig. no. 1]. Crocodiles were observed here during basking from 08.00 A.M. to 05.00 P.M. The presence of 16 - 17 transverse rows of dorsal shield, usually two median rows are considerably broader [Fig. no. 2], 4 - 6 longitudinal series of scutes, broadest snout with 19 upper teeth and an externally visible cone shaped 4th tooth on each side are the characteristics of *Crocodylus palustris* [Da Silva and Lenin, 2010] [Fig. no. 2].




Figure 1: Showing Study area (*Crocodylus palustris*) Kotmi Sonar, Dist. Janjgir Champa, Chhattisgarh



Figure 2

The present study was conducted in Kotmi Sonar. The Kotmi Sonar is an ancient village of District Janjgir - Champa (Chhattisgarh) India. It lies at 22° 01'; 44.8" north latitude

and 820, 21', 13.1" east longitude [Bharos and Kanoje, 2007]. It is situated in Mumbai Howrah railway line between Bilaspur and Champa junctions of Chhattisgarh, India [Fig. no. 1]. The Munda pond is spread on 85 acre area. The terrain of Kotmi Sonar is almost plain with gentle slope. The Kotmi Sonar is situated in 620 meters above M.S.L. The underlying rocks are granite, schist and limestone [Bharos and Kanoje, 2007]. In the present study, four study sites [at four corners] were pointed out.



Figure 3: Area of Kotmi Sonar (Munda pond)

Our qualitative analysis focuses primarily on the behaviours of adults during three phases of reproduction :-

- 1) Defense of Territory and courtship,
- 2) Nesting and Incubation,
- 3) Hatching and Post hatching.

2. Material and Method

The visual ground survey method was used to identify the morphological characteristics and study of behavioral responses. 'Direct Counting Method' and 'Capture and Release Technique' were used to estimate the population of Crocodiles. The indirect evidences such as foot prints, sign of body impressions were also studied. Useful Tools is Technical, camera, interview, visual observation.

Statistical analysis

$$\text{Mean } X = \frac{\sum x}{n}$$

$$\text{Standard or S.D. or } \sigma = \frac{\sqrt{\sum b^2}}{n-1}$$

$$\text{Standard Error. SE} = \frac{SD}{\sqrt{n}}$$

$$\text{Student's test } t = \frac{x_1 - x_2}{\sqrt{SE_1^2 - SE_2^2}}$$

3. Observation

Population estimated by "Direct counting method" and "Capture and release technique".

Visual observation, interaction with local people and Departmental field staff were made for the behavioural study of *Crocodylus palustris*.

Table 1: Population Survey of *Crocodylus Palustris* (Male+Female+Juveniles) in Kotmi Sonar in Janjgir-Champa (C.G.)

Sample Site	Male	Female	Juveniles	Total
A	06	09	50	65
B	06	07	35	48
C	08	11	30	49
D	05	08	45	58
Total	25	35	160	220

Table 2: Observation periods divided into Pre-mating, Mating and Post-mating period based on the timing of the natural mating season

Activities	Observation Period	Method	Behaviour
Pre-mating	1 January 2016 – 31 January 2016	Direct	Nesting (after 1 month)
Mating	1st February 2016 – 30 April 2016	Direct	Egg laying (After 55-75 days)
Post-mating	1st May to August 2016	Direct	Hatching-Behaviour

Table 3: Sound, Signaling, Courtship, Mating, Nesting, Incubation, Pre-hatching and Post-hatching of *Crocodylus palustris*

Sno.	Reproductive Behaviour			
	Activities	Month	Weeks	Observation
1	Pre-mating	January 2016	1 st	Normal activities
			2 nd	Normal activities
			3 rd	Normal activities
			4 th	Some crocodiles between fighting
2	Mating	February 2016	1 st	Normal activities Showing by crocodiles
			2 nd	Pair were showing copulation
			3 rd	Normal activities
			4 th	Normal activities
		March 2016	1 st	Pair were showing copulation
			2 nd	Normal activities
			3 rd	Normal activities
			4 th	Normal activities
		April 2016	1 st	Eggs were laying by female
			2 nd	Normal activities
			3 rd	Eggs were laying by female
			4 th	Normal activities
3	Post-mating	July 2016	1 st	Normal activities
			2 nd	1 female showing Hatching activities
			3 rd	Normal activities
			4 th	Some juveniles into water
		August 2016	1 st	Normal activities
			2 nd	Juveniles were come out from eggs.
			3 rd	Normal activities
			4 th	Some juveniles into water

Table 4: Study Site (A) area of female *Crocodylus palustris* Nesting, Incubation, hatching and Post hatching. Values are mean ± SE of 7 animals

S. No.	Female Length (Meter)	Nesting No.	Egg. Laying in One dutch	Hatching out from egg	Success juveniles
1	3.14	1	25	20	5
2	5.65	1	28	19	6
3	5.00	1	26	21	5
4	8.00	1	27	20	8
5	10.00	1	28	28	6
6	9.00	1	25	20	6
7	3.65	1	27	20	8
Σx	44.44/7	7/7	186/7	138/7	44/7
$\frac{\Sigma x}{N}$					
\bar{X}	6.35	1.00	26.57	19.71	6.28
σ	±2.47	±0.00	±1.17	±0.88	±1.16
SE	±0.93	±0.00	±0.44	±0.33	±0.43

Table 5: Study Site (B) area of female *Crocodylus palustris* Nesting, Incubation, hatching and Post hatching. Values are mean ± SE of 6 animals

S. No.	Female Length (Meter)	Nesting No.	Egg. Laying in One dutch	Hatching out from egg	Success juveniles
1	1	4.56	1	26	15
2	2	4.41	1	25	20
3	3	3.00	1	28	19
4	4	3.85	1	26	19
5	5	3.49	1	27	20
6	6	5.56	1	25	15
Σx	24.96/6	6/6	159/6	108/6	37/6
$\frac{\Sigma x}{N}$					
\bar{X}	4.16	1.00	26.16	18.00	6.16
σ	± 0.84	± 0.00	± 1.06	± 2.16	± 1.34
SE	± 0.34	± 0.00	± 0.43	± 0.88	± 0.55

Table 6: Study Site (C) area of female *Crocodylus palustris* Nesting, Incubation, hatching and Post hatching, Values are mean ± SE of 6 animals

S. No.	Female Length (Meter)	Nesting No.	Egg. Laying in One dutch	Hatching out from egg	Success juveniles
1	8.22	1	28	20	8
2	4.64	1	27	18	6
3	3.89	1	26	20	5
4	3.40	1	28	19	8
5	6.00	1	27	18	5
6	9.64	1	25	15	8
Σx	35.79/6	6/6	161/6	110/6	40/6
$\frac{\Sigma x}{N}$					
\bar{X}	5.96	1.00	26.83	18.33	6.66
σ	± 2.28	± 0.00	± 1.07	± 1.69	± 1.37
SE	± 0.93	± 0.00	± 0.43	± 0.69	± 0.56

Table 7: Study Site (D) area of female *Crocodylus palustris* Nesting, Incubation, hatching and Post hatching, Values are mean ± SE of 6 animals

S. No.	Female Length (Meter)	Nesting No.	Egg. Laying in One dutch	Hatching out from egg	Success juveniles
1	5.89	1	26	19	6
2	5.22	1	28	20	5
3	5.10	1	26	21	8
4	6.45	1	26	19	5
5	6.85	1	26	18	8
6	7.00	1	28	15	7
Σx	36.51/6	6/6	160/6	112/6	39/6
$\frac{\Sigma x}{N}$					
\bar{X}	6.08	1.00	26.66	18.66	6.50
σ	± 0.74	± 0.00	± 0.94	± 1.88	± 1.25
SE	± 0.30	± 0.00	± 0.38	± 0.76	± 0.51

Table 8: Comprative study of A, B, C,D area Female *Crocodylus Palustris*, Nesting, Incubation hatching and post hatching, Values are mean ± SE the figure in parentheses indicate number of animal

Study Site	Female Length (Meter)	Egg Laying in One clutch	Hatching out from egg.	Success juveniles
A	6.35± 0.93 (7)	26.57± 0.44 (7)	19.71± 0.33 (7)	6.28± 0.43 (7)
B	4.16± 0.34* (6)	26.16± 0.43 (6)	18.0± 0.88 (6)	6.16± 0.55 (6)
C	5.96± 0.93 (6)	26.83± 0.43 (6)	18.33± 0.69 (6)	6.66± 0.56 (6)
D	6.08± 0.30 (6)	26.66± 0.38 (6)	18.66± 0.76 (6)	6.50± 0.51 (6)

4. Results

Reproductive Behavior – A adult female Reproductive behavior is well in Mundapond. So *Crocodylus Palustris* (Male+Female+Juveniles) is gradually increasing The Environmental & Physico chemical-condition is well of Munda pond for reproductive so this area is very good for Species of *Crocodylus Palustris*.

The result in this chapter is supported by the statistical analysis presented in the form of table given under different sections.

Population survey of *Crocodylus Palustis* in Kotmi Sonar of Janjgir-Champa (C.G.) (January 2016 – August 2016) show total no of female + male + juveniles.

I observe *Crocodylus Palustris* in Munda pond in this time species show different activities in different month. . My weekly observation show in (January 2016 – August 2016) female *Crocodylus Palustris* so different activities in every week.

Study site A,B,C,D area in Munda pond female *Crocodylus Palustris*, Nestling, Incubation, hatching and post hatching, Values are mean ± SE of 7,6,6,6 animals in Male *Crocodylus Palustris*, nestling, incubation, hatching and post hatching comparative Account of A,B,C,D of site area .The physicochemical factors effect the distribution and population of crocodile [Sahato, 2004]. It was reported that

crocodile population is threatened in low water quality [Joshi et al. 2011]. The variation and access limits of physicochemical parameters were the causes of harmful effects on the population of crocodiles [Chang et al., 2013]. Rubbery of snout and hunch back conditions were occurring in the high level of calcium [Singh et al., 2001]. Low water level without replacement is another cause of bacterial infection in hatchling of Crocodiles [Misra et al., 1993]. The poor water quality and extreme humidity in monsoon causes some myotic infection and death of hatchlings Crocodiles [Lal, 1982; Maskey, 1989].

5. Conclusion

C.P. *Crocodylus palustris* is a Red listed species so conservation is required our nearest place village Kotmi sonar Distt. Janjgir-Champa(C.G.) a big conservation area for crocodile species (*Crocodylus Palustris*). This place soil, water and environmental condition is very perfect for reproduction by *Crocodylus palustris*. Their number increase in this place juveniles and adult growth is well and reproduction condition is well. Female is healthy and strange and their breeding states is very well.

Crocodylus palustris number and population very important for biodiversity and nature and other crocodilians. So save crocodiles you know crocodiles threaten species, so their conservation is very important. Crocodile very important, animal for every think Ex-food biodiversity, medical purpus, ecosystem, food chain, food wed, for fashion, for goods and this is very different another species.

References

- [1] Apha, 14th ed. Standard methods for the examination of water and waste water, Washington, D.C. (1998)journal style)
- [2] Bharos, A. M. K. and Kanoje, R. S., Wetlands of Kotmi Sonar, an abode of marsh Crocodile, Proceedings of Taal, The 12th world lake conference, 1796-1997, (2007).
- [3] Chang, M. S., Gachal, G. S., Qadri, A. H., Sheikh, M. Y., Physico-chemical assessment and its impacts on Marsh Crocodiles of Karachi Zoological Garden, International Journal of Advanced Research, 1, (3), 102-107, (2013).
- [4] Choudhury, B. C., and Chowdhury, Lessons from crocodile reintroduction projects in India, Indian forester 112, 881-890, (1986).
- [5] Da silva, A. and Lenin, J., Mugger Crocodile *Crocodylus palustris*. In: S. C. Manolis, C. Stevenson, (Eds.) Crocodile: Status survey and conservation action plan, Third edition. Crocodile specialist group : Darwin, 94-98, (2010).
- [6] Jacobson, C., Reintroduction of the Mugger Crocodile, *Crocodylus palustris*, in India, Restoration and Reclamation Review, Student on-line Journal, 4, (3), 1-7, (1999).
- [7] Joshi, R., Singh, R. and Negi, M. S., First record of mugger crocodile *Crocodylus palustris* (Lesson, 1831) from the Rajaji National Park, North India, International Journal of Biodiversity and Conservation vol. 3, (9), 444-450, (2011).
- [8] Lal, S., Diseases in Gharial Juveniles, Journal of Wildlife preservation Society of India, 23 (3), 38-40. (1982)
- [9] Maskey, T. M., Movement and survival of captive reared Gharials (*Gavialis gangeticus*) in Narayani River, Nepal, Ph. D. Dissertation, Graduate school, University of Florida, United State of America, (1989).
- [10] Misra, P., Kumar, R. D., Patanayak, G. M., Rahman, R. P., Singha, A., Bacterial isolates from apparently healthy and diseased crocodiles (*Gavialis gangeticus*), Indian Veterinary Journal, 70, 375-376, (1993).
- [11] Oza, G. M., Conservation of the crocodile in the Sayaji Sarovar Lake, Baroda, India. Biology Conservation, 7, 235-236, (1975).
- [12] Sahato, G. A., Lashari, K. H., Sahato, S. B., Hamdard Medicusvol, XL VII No (4), 100, (2004).
- [13] Singh, L. A. K., Observation on the movement of two captive reared Mugger Crocodiles, *Crocodylus palustris* when returned to the wild, Journal of the Bombay Natural history society, 80, 86- 90, 1983.
- [14] Singh, L. A. K., Srivastav, S. S., Mohanty A. P. and Raut, S. D., Prevention of rubbery snout and X-ray revelations on hunchback condition of Mugger crocodile (*Crocodylus palustris*), 220's print Journal 16, (4), 465-466, (2001).
- [15] Vyas, R., Current status of Marsh Crocodiles *Crocodylus palustris* (Reptilia: Crocodylidae) in Vishwamitri River , Vadodara City, Gujrat, India, Journal of threatened taxa, 4, (14), 3333-3341, (2012).
- [16] Vyas, R. and Vyas, R., Mugger survey in the Vishwamitri River of Gujrat, India. CSG Newsletter, 21(3), 9-110, (2002).
- [17] Vyas, R, Mugger (*Crocodylus palustris*) Population in and around Vadodara City, Gujrat State, India. Russian Journal of Herpetology, 17, (1), 43-50, (2010).
- [18] Whitaker, R. and Whitaker, Z., Reproductive biology of the Mugger, J. Bombay Natural history society 81, (2), 297-316, (1984).