

Assessment of Infant and Young Child Feeding Indicators with Special Emphasis on Practices and Knowledge of Mothers in Rural Areas

Taslima Arzu¹, Md. Ariful Kabir Sujan², Dr. Md. Sabir Hossain³

¹Jahangirnagar University, Department of Biochemistry and Molecular Biology, Savar, Dhaka, Bangladesh

²Nutrition Consultant, Nutrition International, Bangladesh

³Jahangirnagar University, Department of Biochemistry and Molecular Biology, Savar, Dhaka, Bangladesh

Abstract: *The IYCF practices have a great impact on the physical and mental development of the child. The objective of this study was to study the knowledge and compliance to IYCF practices along. The lactating mothers with the children in the age group 0-11 months were studied. Mother's knowledge level is average about IYCF components; 44.7% mothers have proper knowledge about IYCF practice. 61.2% mothers have knowledge on initiation of breast-feeding within an hour of birth while 50.7% of the mothers initiated breast-feeding within an hour after birth. Exclusive breast-feeding for 6 months was practiced by the mothers of 65.6%. 64% of the mothers told that complimentary feeding should start at 6 months and 59.8% of the mother will continue breast fed for 2 years of infants.*

Keywords: Infant and Young Child Feeding (IYCF), Breast feeding, Complementary feeding, Lactating mother

1. Introduction

The optimal infant and young child feeding practices during the first 2 years of life is of paramount importance as this period is the "critical window" for the promotion of health, good growth, behavioral and cognitive development. Optimal infant and young child feeding practices include initiation of breast-feeding within 1 hour of birth, exclusive breast-feeding for the first 6 months, and continuation of breast-feeding for 2 years or more, along with nutritionally adequate, safe, ageappropriate, responsive complementary feeding starting at 6 months.[1] Breast-feeding strengthens emotional security and affection creating a strong bond between the mother and the child, which in turn promotes psychosocial development of a child. To ensure good nutrition status of the infant as well as the mother, maternal nutrition plays a vital role. Breast-feeding is nature's way of nurturing the child. It provides learning and development opportunities to the infant. Breast milk also leads to increased intelligence quotients and better visual acuity due to the presence of special fatty acids in it.[2]

Approximately, 1.4 million deaths of children under the age of 5 years worldwide can be attributed to suboptimal breast-feeding. Almost 6% of under-five mortality can be prevented by the timely introduction of complementary feeding.[3] It was estimated that about one-fifth of overall under-five mortality can be averted if 90% infants are covered with an inclusive package of interventions to promote, protect, and support the optimal infant young child feeding (IYCF) practices.[3] A large proportion of children become vulnerable to stunting, poor cognitive development, and significantly increased risk of infectious diseases, such as, diarrhea and acute respiratory infection due to the poor complementary feeding practices.[4]

It has been established that because of the best bioavailable iron in breast milk, exclusive breast-feeding prevents anemia and infections particularly the diarrheal infections in the child. The need of introducing cereal-based foods in the diet of infant after the age of 6 months can be correlated with the fact that enzyme amylase appears in the seventh month of the infant.[5] The mother's risk for excess postpartum bleeding is decreased if breast-feeding is initiated early, which in turn lowers the risk for anemia. Exclusive breast-feeding delays next pregnancy boosts mother's immunity and reduces the insulin needs of diabetic mothers. Breast-feeding also provides protection from breast and ovarian cancers and osteoporosis.[6] This has an enormous impact in a developing country, like India, with a high burden of disease and low access to safe water and sanitation. The recent studies conducted even in developed countries have also emphasized the role of IYCF practices in reducing child mortality.[7] A global strategy for infant- and young child-feeding has been devised by the World Health Organization (WHO) and United Nations Children Fund. Based on these guiding principles, the Government of India, in collaboration with international agencies, has adopted the culturally acceptable IYCF guidelines, which were incorporated in the Integrated Management of Neonatal and Childhood Illness Programme. [8]

These guidelines recognize appropriate infant feeding practices to be crucial for improving nutrition status and decreasing infant mortality in all countries. WHO offers three recommendations for IYCF practices for children aged 6-23 months: continued breast-feeding or feeding with appropriate calcium-rich foods if not breast-fed; feeding solid or semisolid food for a minimum number of times per day according to age and breast-feeding status; and including foods from a

minimum number of food groups per day according to breast-feeding status.

Bangladesh Demographic and Health Survey (BDHS 2014) have provided useful national- and state-level information on the IYCF practices.[9] Available data showed a gross interstate variation. However, the BDHS was not designed to provide district-level data. According to the BDHS-2014 data Fifty-five percent of infants under age 6 months are exclusively breastfed. The Multiple Indicator Cluster Survey 2012-13 reported lower exclusive breastfeeding rates of 56 percent [10]. Overall, 26 percent of breastfed children age 6-23 months are given the recommended four or more food groups, and 63 percent are fed at least the minimum number of times.

According to the Multiple Indicator Cluster Survey 2012-13 (MICS-2012-13) early initiation of breast feeding (within an hour of delivery) is 57.4%, continued breast feeding up to 2 years 87.5%. [10]

With this background, this study was undertaken to assess the IYCF knowledge and practices with special emphasis on IYCF indicators, among children aged less than 1 year among 16 rural areas or villages at Maligasa Union, Pabna district, Bangladesh.

2. Methods and Materials

It was a convenient sample survey study conducted on April, 2017, at Maligasa Union, Pabna district, Bangladesh. This union caters population of about 25000 in the field practice area by providing primary and promotive health care. Study population comprised of mothers having children of age group 0–11 months. A total of 353 eligible mothers were approached through house-to-house visit to participate in the study by convenient sampling method. They were informed about the purpose of study and informed consent was obtained from the mothers. The data were collected by interview method using a pretested schedule. Data were analyzed through Epi info and SPSS software. WHO indicators for assessing infant and child feeding practices were used. A pretested questionnaire mainly based on the standard questionnaire on IYCF practices given by WHO was used for data collection. [11] These questions provide the information needed to calculate the key indicators of IYCF. As per WHO recommendations, information was collected about the child's diet in the previous 24 hours, which included the type of food items and the number of times they had consumed. Food items were categorized into seven types, that is, cereals, legumes and nuts, dairy products, meat products, egg, vitamin A-rich fruits and vegetables, and other fruits and vegetables. Children less than 12 months were included in the study after obtaining verbal informed consent from the mothers.

3. Results and Discussion

Out of the 353 children studied, 51.3% were boys and 48.7% were girls. Table 1 gives the complete information regarding the sociodemographic profile of the population studied.

Table 1: Socio demographic profile of the studied population

Parameter	Categories	Frequency (N=353)	(%)
Gender	Male	181	51.3
	Female	172	48.7
Age of the kid (in months)	0-5	164	46.5
	6-11	189	53.5
Religion	Muslim	349	98.9
	Hindu	04	1.1
Mother's educational qualification	B.A. or B.Sc or above	20	5.7
	Intermediate	64	18.1
	High School	126	35.7
	Primary School & below	143	40.5
Mother's knowledge assessment status	Excellent (100% knowledge)	35	9.9
	Good (60-80% knowledge)	123	34.8
	Average (40-60% knowledge)	189	53.6
	Below average (below 40% knowledge)	06	1.7

Table 2: Summary of the knowledge to key IYCF Practices

Component	Knowledge	Frequency	(%)
Timely Initiation of Breast Feeding (<1 hour from Birth)	Don't Know	7	2.0
	After 3/4 days	21	5.9
	After 8/10 hours	109	30.9
What is colostrum	Immediately	216	61.2
	Don't Know	71	20.1
Benefits of colostrum for baby (Multiple response)	Yellowish secretion after delivery	282	79.9
	Don't Know	71	20.1
Benefits of colostrum for mother & Family (Multiple response)	Good for health	162	45.9
	No cost for family	211	59.8
	Baby's first vaccine	179	50.7
Exclusive Breast Feeding (for 6 months)	Increase immunity	63	17.8
	Don't know	23	6.5
	Up to 5 months	76	21.5
	Up to 6 months	219	62.0
Timely Initiation of Complimentary Feeding (at 6 months)	Up to 1 year	35	10.0
	Don't know	37	10.5
	4-5 months	71	20.1
Continued Breast Feeding (for 2 Years)	After 6 months	226	64.0
	After 9 months	19	5.4
	Up to 2 years	211	59.8
As long as baby drink	Up to 1 year	70	19.8
	72	20.4	

Table 3: Summary of the compliance to key IYCF Practices

Component	Compliance	Frequency	(%)
Timely Initiation of Breast Feeding (<1 hour from Birth)	Within 1 hour	179	50.7
	Within 24 hours	111	31.4
	After 2/3 days	63	17.9
How often you breastfeed your children	8-12 times a day	243	68.8
	6-8 times a day	91	25.8
	When baby cry	19	5.4
Exclusive Breast Feeding (for 6 months) (N=189)	Up to 4-5 months	52	27.5
	Up to 6 months	124	65.6
	Up to 1 Years	13	6.9

Minimum diet diversity of Complementary feeding (at least 4 food groups) (N=189)	Acceptable diet	58	30.7
	Not acceptable diet	131	69.7
Minimum Meal Frequency (N=189)	2/3 times	96	50.8
	3/4 times	59	31.2
	4/5 times	28	14.8
	Others (one time)	06	3.2

3.1. Infant and Young Child Feeding (IYCF) knowledge of the mother

The knowledges of the mothers on Infant and Young Child Feeding (IYCF) components were average. Most of the mothers have knowledge on best practices of IYCF components lies in average categories (53.6% mothers with knowledge of 40-60% information about IYCF) and 44.7% mother have proper knowledge about IYCF practice (9.9% mothers know all information about IYCF and 34.8% mothers know 80% information). 61.2% of mothers have proper knowledge about timely initiation of breast feeding within one hour of birth. 79.9% Mothers knew that what is colostrum and the benefits of colostrum for the baby, mothers and family. Unfortunately, still 20.1% of mother don't know about colostrum and it's benefits for the children, mother and family. 62% mothers have knowledge about the continuation time of exclusive breast feeding up to 6 months of child's age and 38% mothers know wrong continuation time (don't know 6.5%, up to 5 months 21.5% and up to 1 year 10%). The knowledge about the timing of starting complementary feeding for the children of the mothers was quite good. 64% of the knew the exact timing of starting complementary feeding (after 6 months of baby's age). 36% mother had wrong knowledge about the starting time of complementary feeding.

3.2. Infant and Young Child Feeding (IYCF) practices of the mother

An epidemiological evidence of a causal association between early initiation of breast-feeding and reduced infection-specific neonatal mortality has also been documented. [12] The Survey result shows that 50.7% children were initiated breast feeding within one hour of delivery.

Exclusive breast-feeding was done by 65.6% of 189 children over 6 months of age. This was far better than the figures reported by BDHS-2014 data, at national level (55%). [9]

Minimum dietary diversity (MDD) indicator is the proportion of children of 6–23 months of age who receive foods from four or more food groups from a total of seven food groups, such as, dairy products, legumes and nuts, flesh foods, eggs, vitamin A-rich fruits and vegetables, cereals and tubers, and other fruits and vegetables.[13]This indicator reveals whether the child is receiving a complete and balanced diet or not. MDD was observed in only 30.7% children between 6 and 11 months age group.

Minimum Meal Frequency (MMF) indicator is the proportion of breast-fed and non-breast-fed children aged 6–23 months who receive solid, semisolid, or soft foods (but also including milk feeds for non-breast-fed children) the minimum number of times or more.[13]For breast-fed children, the minimum number of times varies with age (two times if 6–8 months and three times if 9–23 months). For non-breast-fed children, the minimum number of times does not vary by age (four times for all children aged 6–23 months). MMF was observed in the majority (96.8%) of children aged 6–11 months.

4. Recommendation

The IYCF practices are strongly influenced by what people know, think and believe and also affected by social circumstances and economic factors. Effective communication for behavioral change is necessary for ensuring optimal infant feeding. Awareness regarding IYCF practices and their benefits in Maternal and Child Health (MCH) is poor leading to poor compliance. It is important to educate mothers during the antenatal visits. The situation can be improved by training of grass root health workers on IYCF policies of WHO and MoHFW, Govt. of Bangladesh, stressing on the benefits of appropriate feeding practices by the hospitals, Health & Family Welfare Centre (H&FWC), Union Health Sub Centre (USC) and Community Clinic and making these services universally available along with intensive IEC (Information, Education & Communication) efforts to generate demand for these services. Most of the world's religions place particular emphasis on the total care of the child. In the context of the overwhelming evidence, the involvement of religious teachings in the promotion of breastfeeding is quite debatable. It is well established that religious ideologies influence the human mind and a person's way of living.

Health professionals traditionally encourage mothers to breastfeed by giving information on benefits of breastfeeding for the infant and the mother herself. The behavior of women can be easily modified through religious teachings in a positive way. Breastfeeding may be affected by religious ideologies using the doctrine in religious texts. Counseling the mothers by reinforcing the cultural and religious practices supporting breastfeeding can help enormously. Use of local religious teachings can bring positive changes in the implementation of health programs [14]. In addition, public nutrition education that promotes infant and young child feeding as defined by WHO, taking into account social-cultural factors is needed and recommended.

References

- [1] World Health Organization. Infant and Young Child Feeding: Model Chapter for Textbooks for Medical Students and Allied Health Professionals. France: World Health Organization, 2009.
- [2] Ten steps to successful Breastfeeding. UNICEF/WHO Baby Friendly Hospital Initiative (BFHI). Initiation of

- breastfeeding by breast crawl. Available at: <http://breastcrawl.org/10steps.shtml>.
- [3] Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet* 2003;361:2226–34.
- [4] WHO. Effect of breastfeeding on infant and child mortality due to infectious diseases in less developed countries: a pooled analysis. Collaborative Study Team on the role of breastfeeding on the prevention of infant mortality. *Lancet* 2000;355:451–5.
- [5] Butte NF, Lopez-Alarcon MG, Garza C. Nutrient Adequacy of Exclusive Breastfeeding for the Term Infant during the First Six Months of Life. Geneva: World Health Organization, 2002.
- [6] Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS. Bellagio Child Survival Study Group. How many child deaths can we prevent this year? *Lancet* 2003;362:65–71.
- [7] WHO. Implementing the Global Strategy for Infant & Young Child Feeding. Geneva, 3–5 Feb, 2003.
- [8] Government of India. National Guidelines on Infant and Young Child Feeding. Ministry of Human Resource development, New Delhi; Department of Women and Child development, 9–10, 2006.
- [9] Bangladesh Demographic and Health Survey 2014
- [10] The Multiple Indicator Cluster Survey 2012-13 (MICS-2012-13) by Bangladesh Bureau of Statistics and UNICEF, Bangladesh.
- [11] Indicators for Assessing Infant and Young Child Feeding Practices: Conclusions of a Consensus Meeting Held 6–8 November 2007 in Washington, DC, USA. World Health Organization, 2008. Available at: http://whqlibdoc.who.int/publications/2008/9789241596664_eng.pdf (last accessed on Apr 29, 2012).
- [12] Edmond KM, Kirkwood BR, Amenga-Etego S, Owusu-Agyei S, Hurt LS. Effect of early infant feeding practices on infection-specific neonatal mortality: an investigation of the causal links with observational data from rural Ghana. *Am J Clin Nutr* 2007;86:1126–31.
- [13] Indicators for Assessing Infant and Young Child Feeding Practices: Conclusions of a Consensus Meeting Held 6–8 November 2007 in Washington, DC, USA. World Health Organization, 2008. Available at: http://whqlibdoc.who.int/publications/2008/9789241596664_eng.pdf (last accessed on Apr 29, 2012).
- [14] Iftikhar Alam and Safia Begum (2014)
- [15] Advocating breastfeeding - can religious teachings bring a change? *Public Health* by Nova Science Publishers pp. 165-167.