# Comparing Different Bayes Estimator of Shape Parameter for Burr Type X

#### Maha Adil Alagaidi

Assistant Lecturer, University of Baghdad/ Post - Graduate Institute for Accounting & Financial Studies

Abstract: This paper deals with obtaining a new probability distribution from powering some given distribution to obtain a family that is more flexible for representing data, then we derive four Bayes estimators under exponential priors, also four Bayes estimators under Jeffery prior, then the comparison has been done through simulation, the results are compared using statistical measures (mean square error).

**Keywords:** A new family Burr type X, Generalized Rayleigh, Bayes I, Bayes II, Bayes D, Bayes P under exponential prior, Jeffery prior, MSE. Risk function

#### 1. Introduction

Many new probability distribution can be obtained from powering the CDF of any given distribution to a positive real number  $(\lambda)$ , this gives new family which may be more

flexible and fitting for the observation of random variable. The new CDF [G(z)] obtained from any given CDF [F(z)] is;  $G(z) = [F(z)]^{\lambda}$   $\lambda > 0$  (1)

We know that the generalized Rayleigh distribution, have the CDF;

$$F(x;\alpha,\lambda) = \left(1 - e^{-(\alpha x)^2}\right)^{\lambda} \quad x,\alpha,\lambda > 0$$
 (2)

We assume the scale parameter ( $\alpha = 1$ ), then;

$$F(x;\lambda) = \left(1 - e^{-x^2}\right)^{\lambda}$$
(3)

This CDF in equation (3) called (Burr type X) distribution with p.d.f;

$$f(x;\lambda) = 2\,\lambda\,x\,e^{-x^2} \left(1 - e^{-x^2}\right)^{\lambda-1} \,x,\lambda > 0 \qquad (4)$$

#### 2. Aim of Research

We work on comparing four Bayes estimators for parameter  $(\lambda)$  of the distribution (Burr type X), using different model

of prior distribution for  $(\lambda)$ , since the Bayes estimators

depend on the theory that the parameter to be estimated is considered random variable and having prior distribution  $\{g(\lambda), [\text{Thomas Bayes (1702)}], \text{ the comparison has been } \}$ 

done through simulation procedure using different sample size.

## **3. Theoretical Aspect**

Let  $(x_1, x_2, ..., x_3)$ , be *i.d.d* from  $[f(\underline{x}|\lambda)]$  defined in equation (4), then the Bayesian estimator depend on observations represented by conditional density, which consider the parameter  $(\lambda)$  as a random variable, this lead to find the posterior distribution of  $(\lambda)$  given data

 $(x_1, x_2, \dots, x_3);$ 

$$f(\underline{\lambda}|x) = \frac{f(\underline{x}|\lambda)g(\lambda)}{f(\underline{x})}$$
(5)

Where (in case of continuous);

$$f(\underline{x}) = \int_{\forall \lambda} f(x|\lambda) g(\lambda) d\lambda$$

Bayes estimator (*Bayes*) need loss function  $[L(\hat{\lambda}, \lambda)]$  which is the losses incurred by estimating  $(\lambda)$  by  $(\hat{\lambda})$ , then measuring the loss by difference between  $(\hat{\lambda} - \lambda)$ , the loss function  $[L(\hat{\lambda}, \lambda)]$  is defined as a real valued function satisfy; a.  $L(\hat{\lambda}, \lambda) \ge 0 \quad \forall \hat{\lambda} \neq \lambda$ 

$$D_{\lambda} L(\lambda, \lambda) = 0 \quad \forall \lambda = \lambda$$

We can give some types of loss function that may use in comparing the Bayes estimator  $(\lambda)$ ;

#### A. Squared error loss function

This is from symmetric loss function given by Mood, Graybill, and Boes (1974), which is;

$$L(\hat{\lambda},\lambda) = (\hat{\lambda} - \lambda)^{2}$$
(6)

The Byes estimator  $(\hat{\lambda}_{Bayesl})$  is obtained from minimizing risk function  $[R_s(\hat{\lambda}, \lambda)]$ .

$$R_{s}(\hat{\lambda},\lambda) = E(loss) = E[L(\hat{\lambda},\lambda)] = \int_{\forall\lambda} L(\hat{\lambda},\lambda)f(\lambda|\underline{x})d\lambda$$
$$R_{s}(\hat{\lambda},\lambda) = \int_{\forall\lambda} (\hat{\lambda}-\lambda)^{2}f(\lambda|\underline{x})d\lambda \quad (7)$$
$$R_{s}(\hat{\lambda},\lambda) = \int_{\forall\lambda} \hat{\lambda}^{2}f(\lambda|\underline{x})d\lambda - 2\int_{\forall\lambda} \hat{\lambda}\lambda f(\lambda|\underline{x})d\lambda + \int_{\forall\lambda} \lambda^{2}f(\lambda|\underline{x})d\lambda$$
$$R_{s}(\hat{\lambda},\lambda) = \hat{\lambda}^{2} - 2\hat{\lambda}E(\lambda|\underline{x}) + E(\lambda^{2}|\underline{x}) \quad (8)$$

Put 
$$R_s(\hat{\lambda}, \lambda) = 0$$
, this gives the first Bayes estimator;

$$\hat{\lambda}_{BayesI} = E(\lambda | \underline{x}) \tag{9}$$

Which is the conditional mean of posterior distribution  $[f(\lambda | \underline{x})]$ .

## Volume 6 Issue 10, October 2017

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

#### DOI: 10.21275/ART20177152

#### **B.** De Groot loss function

This loss function was proposed by De Groot (1970), the Bayes estimator for the positive parameter  $(\lambda > 0)$  by this

method obtained using the loss function;

$$L(\hat{\lambda}, \lambda) = \left(\frac{\lambda - \hat{\lambda}}{\hat{\lambda}}\right)^2 \tag{10}$$

Then the risk function and Bayes estimator  $(\hat{\lambda}_{Bayes 11})$  under

this loss function is;  

$$R_{D}(\hat{\lambda}, \lambda) = \hat{\lambda}^{-2} E(\lambda^{2} | \underline{x}) - 2\hat{\lambda}^{-1} E(\lambda | \underline{x}) + 1$$

$$\hat{\lambda}_{De\ Groot} = \hat{\lambda}_{D} = \frac{E(\lambda^{2} | \underline{x})}{E(\lambda | x)}$$
(11)

To derive the formulas Bayes estimators, we need to find the posterior  $[f(\lambda | \underline{x})]$  from equation (5), to simplify the work,

$$f(x,\lambda) = 2\lambda x e^{-x^2 + \ln(1 - e^{-x^2})} e^{-\lambda \ln(1 - e^{-x^2})}$$
(12)

Then using this equation according to two types of prior  $[g_1(\lambda),g_2(\lambda)]$ , we find posterior  $[f(\lambda|\underline{x})]$ , then using four

types of loss function to find  $[\hat{\lambda}_{Bayes}]$ .

#### **D.** Precautionary Loss Function

This type of loss function was given by Norstrom (1996), which use the loss function given in equation (10), then the risk function and Bayes estimator for this loss and risk function is;

$$R_{p}(\hat{\lambda},\lambda) = \hat{\lambda} - 2E(\lambda|\underline{x}) + \hat{\lambda}^{-1}E(\lambda^{2}|\underline{x})$$
(13)

Therefore the third Bayes estimator is;

$$\hat{\lambda}_{3 Bayes} = \sqrt{E(\lambda^2 | \underline{x})}$$
(14)

While the fourth Bayes estimator of  $(\lambda)$  is obtained under entropy loss function which was proposed by Calabria and Pulcini (1994), which is;

$$L(\hat{\lambda},\lambda) = \left[ (\hat{\lambda}|\lambda)^2 - t \ln(\hat{\lambda}|\lambda) - 1 \right]$$
(15)

Risk function according to this loss function is;  $R_{E}(\hat{\lambda}, \lambda) = \hat{\lambda}^{t} E(\lambda^{-t} | \underline{x}) - t \ln \hat{\lambda} + t E(\lambda^{2} | \underline{x}) - 1 \quad (16)$ 

Therefore the fourth Bayes estimator for  $(\lambda)$  is;

$$\hat{\lambda}_{4 \, Bayes} = E \left( \lambda^{-t} | \underline{x} \right)^{-\frac{1}{t}}$$
(17)

#### 4. Bayesian Analysis

 $L(x|\underline{\lambda}) = \prod f(x_i,\lambda)$ 

From the studied p.d.f as defined in equation (4) and rewritten as equation (12) we find;

Let;  

$$K = 2^{n} exp \left[ \sum_{i=1}^{n} \ln x_{i} - \sum_{i=1}^{n} x_{i}^{2} + \sum_{i=1}^{n} \ln \left( 1 - e^{-x_{i}^{2}} \right)^{-1} \right]$$
(19)  

$$q_{i}(x) = \sum_{i=1}^{n} \ln \left( 1 - e^{-x_{i}^{2}} \right)^{-1}$$
(20)

 $g_1(x) = \sum_{i=1}^{n} \ln(1 - e^{-x_i})$  (20)

First of all we find the posterior distribution  $[h(\lambda | \underline{x})]$ ;

$$h_1(\lambda|\underline{x}) = \frac{\prod_{i=1}^n f(x_i,\lambda)g(\lambda)}{\int_{\forall\lambda} \prod_{i=1}^n f(x_i,\lambda)g(\lambda)d\lambda}$$
(21)

Assuming that  $(\lambda)$  is random variable have exponential distribution with parameter (b);

$$g(\lambda) = \begin{array}{c} b e^{-\lambda b} & \lambda > 0\\ 0 & b > 0\\ 0 & o/w \end{array}$$
(22)

Then using *p.d.f* in equation (4), the prior in (22), applying (21) we find the equation of posterior distribution  $[h(\lambda | \underline{x})]$  under this assumed prior (22), i.e;

$$h_1(\lambda|\underline{x}) = \frac{[g_1(x) + b]^{n+1}}{\Gamma(n+1)} \lambda^n e^{-\lambda[g_1(x) + b]}$$
(23)

According to the posterior  $[h_1(\lambda | \underline{x})]$  and types of loss function selected, we find that  $[\hat{\lambda}_{Bayes}]$  formulas are;

 $\hat{\lambda}_{Bayesl}$  under squared error loss function is;

$$\hat{\lambda}_{Bayes1} = \frac{n+1}{g_1(x)+b} \tag{24}$$

Risk function under this estimator;

$$\hat{R}_{BayesI} = \frac{n+1}{[g_1(x)+b]^2}$$
(25)

While the second type of Bayes estimator  $(\hat{\lambda}_{2D})$  under De Groot loss function is;

$$\hat{\lambda}_{2D} = \frac{n+2}{g_1(x)+b}$$
 (26)

$$\hat{R}_{2D} = \frac{1}{n+2}$$
 (27)

While the third Bayes estimator of  $(\lambda \rightarrow \hat{\lambda}_{Bayes3})$  according

to precautionary loss function is;

$$\hat{\lambda}_{3P} = \sqrt{\frac{(n+1)(n+2)}{(g_1(x)+b)^2}}$$
(28)

$$\hat{R}_{3P} = \frac{2\sqrt{(n+1)(n+2)} - (n+1)}{(g_1(x) + b)}$$
(29)

The fourth Bayes estimator according to exponential prior and entropy function is;

$$\hat{\lambda}_{4E} = \sqrt[t]{\left[\frac{\Gamma(n+1-t)}{\Gamma(n+1)}(g_1(x)+b)^t\right]^{-1}}$$
(30)

Which can be simplified to;

$$L(\underline{x}|\underline{\lambda}) = 2^{n}\lambda^{n}exp\left[\sum_{i=1}^{n}\ln x_{i} - \sum_{i=1}^{n}x_{i}^{2} + \sum_{i=1}^{n}\ln\left(1 - e^{-x_{i}^{2}}\right)^{-1}\right]exp\left[-\lambda\sum_{i=1}^{n}\ln\left(1 - e^{-x_{i}^{2}}\right)^{-\hat{\lambda}^{1}}\right]\varepsilon = \frac{\sqrt[t]{\frac{\Gamma(n+1)}{\Gamma(n+1-t)}}}{(\underline{u}\underline{u}\underline{b}\underline{x}) + b)}$$
(31)  
$$\hat{R}_{4\varepsilon} = \ln\left[\frac{\Gamma(n+1-t)}{\Gamma(n+1)}(g_{1}(x) + b)^{t}\right] + t\left[\frac{(g_{1}(x) + b)^{n+1}}{\Gamma(n+1)}\sum_{i=1}^{n}\frac{(-g_{1}(x) - b)^{n+1}}{k!(n+1)}\right]$$

#### Volume 6 Issue 10, October 2017

#### <u>www.ijsr.net</u>

## Licensed Under Creative Commons Attribution CC BY

DOI: 10.21275/ART20177152

Now we assume second prior which is Jeffrey's prior defined by;

$$g_2(\lambda) = \frac{\kappa}{\lambda^3} \tag{33}$$

$$h_{2}(\lambda|\underline{x}) = \frac{\prod_{i=1}^{n} f(x_{i}\lambda)g_{2}(\lambda)}{\int_{\forall \lambda} \prod_{i=1}^{n} f(x_{i}\lambda)g_{2}(\lambda)d\lambda} = \frac{k\lambda^{n-3}e^{-\lambda g_{1}(\lambda)}}{\int_{0}^{\infty} k\lambda^{n-3}e^{-\lambda g_{1}(\lambda)}d\lambda}$$
(34)

The second posterior distribution of  $(\lambda | \underline{x})$  is;

$$h_2(\lambda|\underline{x}) = \frac{[g_1(x)]^{n-3}}{\Gamma(n-3)} \lambda^{n-3} e^{-\lambda g_1(x)} \qquad \lambda > 0 \quad (35)$$

Which is also Gamma  $(n - 3, \lambda)$ 

The fourth Bayesian estimators under Jeffrey's prior (using squared error loss function) are;

a. 
$$\hat{\lambda}_{BayesIJ} = \frac{n-3}{g_1(x)}$$
  
 $\hat{R}_{BayesIJ} = \frac{n-3}{[g_1(x)]^2}$ 

$$(36)$$

b. The second Bayes estimator under Jeffrey's using De Groot loss function is;

$$\hat{\lambda}_{BayesDJ} = \frac{n-2}{g_1(x)}$$

$$\hat{R}_{BayesDJ} = \frac{1}{n-2} \quad n > 2 \quad (37)$$

c. The third Bayes estimator under Jeffrey's and precautionary loss function is;

$$\hat{\lambda}_{3BayesJ} = \sqrt{\frac{(n-3)(n-2)}{[g_1(x)]^2}} \\ \hat{R}_{3BayesJ} = \frac{2\sqrt{n-3}(n-2)-n}{[g_1(x)]} \quad n > 2$$
(38)

Finally the fourth Bayes estimator for  $(\lambda)$  using entropy loss function and Jeffery's prior  $[h(\lambda) = \frac{k}{\lambda^3}]$  is;

$$\hat{\lambda}_{4BayesE} = \frac{t \sqrt{\frac{\Gamma(n+1)}{\Gamma(n+1-t)}}}{g_1(x)} \\ \hat{R}_{4BayesE} = \ln \left[ \frac{\Gamma(n-t)}{\Gamma(n)} (g_1(x))^t \right] + t \left[ \frac{(g_1(x))^n}{\Gamma(n)} \sum_{k=0}^n \frac{(-g_1(x))^k}{k! (n+1+k)^2} \right]$$
(39)

## **5. Simulation Procedures**

To find the estimator's  $(\hat{\lambda}_{Bayes1}, \hat{\lambda}_{Bayes2}, \hat{\lambda}_{Bayes2}, \hat{\lambda}_{Bayes3}, \hat{\lambda}_{Bayes4})$ 

we perform simulation experiments using Monte Carlo assuming that;

Ь	0.8	1	1.5
λ	0.9	1.7	3

Table 1: MSE Four Bayes ( $b = 0.8, \lambda$	= 0.9)
--	--------

n	λ <sub>Bayest</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.0724	0.0251	0.0246	0.0307	λ <sub>Bayes3</sub>
40	0.0281	0.0162	0.0168	0.0211	λ <sub>Bayes3</sub>
60	0.0186	0.0101	0.0102	0.0112	λ <sub>Bayes2</sub>
80	0.0107	0.0054	0.0055	0.0061	λ <sub>Bayes2</sub>

100	0.0052	0.0049	0.0050	0.0053	λ <sub>Bayes2</sub>	
-----	--------	--------	--------	--------	---------------------	--

**Table 2:** MSE Four Bayes ( $b = 1, \lambda = 0.9$ ).

n	λ <sub>BayesI</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.0341	0.0250	0.0249	0.0279	λ <sub>Bayes3</sub>
40	0.0216	0.0173	0.0163	0.0181	λ <sub>Bayes3</sub>
60	0.0107	0.0103	0.0101	0.0106	λ <sub>Bayes2</sub>
80	0.0063	0.0056	0.0055	0.0057	λ <sub>Bayes2</sub>
100	0.0045	0.0042	0.0042	0.0041	λ <sub>Bayes4</sub>

Table 3: MSE Four Bayes ( $b = 1.5, \lambda = 0.9$ ).

n	λ <sub>BayesI</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.1224	0.1040	0.1068	0.1348	λ <sub>Bayes2</sub>
40	0.0802	0.0682	0.0710	0.0853	λ <sub>Bayes2</sub>
60	0.0387	0.0354	0.0361	0.0407	λ <sub>Bayes2</sub>
80	0.0206	0.0200	0.0203	0.0216	λ <sub>Bayes2</sub>
100	0.0132	0.0121	0.0124	0.0138	λ <sub>Bayes2</sub>

Table 4: MSE Four Bayes ( $b = 0.8, \lambda = 1.7$ ).

n	λ <sub>BayesI</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.1507	0.1071	0.1038	0.1130	λ <sub>Bayes3</sub>
40	0.1030	0.0678	0.0688	0.0750	λ <sub>Bayes2</sub>
60	0.0432	0.0350	0.0355	0.0383	λ <sub>Bayes2</sub>
80	0.0228	0.0202	0.0201	0.0211	λ <sub>Bayes3</sub>
100	0.0138	0.0122	0.0123	0.0128	λ <sub>Bayes2</sub>

**Table 5:** MSE Four Bayes ( $b = 1, \lambda = 1.7$ ).

n	λ <sub>Bayesi</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.5228	0.4503	0.4605	0.5725	λ <sub>Bayes2</sub>
40	0.3007	0.2720	0.2754	0.3213	λ <sub>Bayes2</sub>
60	0.2010	0.1741	0.1782	0.2086	λ <sub>Bayes2</sub>
80	0.1201	0.1112	0.1134	0.1231	λ <sub>Bayes2</sub>
100	0.0833	0.0804	0.0803	0.0865	λ <sub>Bayes3</sub>

**Table 6:** MSE Four Bayes ( $b = 1.5, \lambda = 1.7$ ).

n	λ <sub>Bayesi</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.6363	0.3703	0.04484	0.4567	λ <sub>Bayes3</sub>
40	0.3478	0.2436	0.2710	0.2760	λ <sub>Bayes3</sub>

## Volume 6 Issue 10, October 2017

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

## International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

60	0.2103	0.1640	0.1752	0.1802	λ <sub>Bayes2</sub>
80	0.1281	0.1070	0.1118	0.1141	λ <sub>Bayes2</sub>
100	0.0884	0.0767	0.0805	0.0806	λ <sub>Bayes3</sub>

**Table 7:** MSE Four Bayes ( $b = 0.8, \lambda = 3$ ).

	ä	1	ň.	1	
n	A Bayest	A Bayes2	A Bavesa	A Baves4	Best
	Duycor	54,002	24,000	54,004	
20	0 1627	0 1651	0.1603	0.1682	2-
20	0.1027	0.1051	0.1005	0.1002	~ Bayess
40	0 1345	0.1330	0 1314	0 1377	2
-10	0.1545	0.1550	0.1514	0.1577	~ Bayess
60	0.1055	0.1041	0.1037	0.1072	2
00	011000	011011	011007	0.1072	- Bayess
80	0.0827	0.0818	0.0816	0.0837	2 Barrage
00	0.002/	0.0010	0.0010	0.0007	- Duyesa
100	0.0687	0.0680	0.0678	0.0704	2 Parrage
- 00	212007	212000	212070		- Dayess

**Table 8:** MSE Four Bayes ( $b = 1, \lambda = 3$ ).

n	λ <sub>Bayes1</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.1764	0.1485	0.1610	0.1557	λ <sub>Bayes2</sub>
40	0.1418	0.1246	0.1315	0.1302	λ <sub>Bayes2</sub>
60	0.1105	0.1002	0.1036	0.1032	λ <sub>Bayes2</sub>
80	0.0851	0.0807	0.0816	0.0814	λ <sub>Bayes2</sub>
100	0.0703	0.0666	0.0678	0.0678	λ <sub>Bayes2</sub>

n	λ <sub>Bayesi</sub>	λ <sub>Bayes2</sub>	λ <sub>Bayes3</sub>	λ <sub>Bayes4</sub>	Best
20	0.6023	0.5116	0.5235	0.6770	λ <sub>Bayes2</sub>
40	0.4080	0.3483	0.3631	0.4417	λ <sub>Bayes2</sub>
60	0.1877	0.1745	0.1773	0.2057	λ <sub>Bayes2</sub>
80	0.1160	0.1087	0.1107	0.1207	λ <sub>Bayes2</sub>
100	0.0863	0.0815	0.0830	0.0886	λ <sub>Bayes2</sub>

**Table 9:** MSE Four Bayes ( $b = 1.5, \lambda = 3$ )

## 6. Conclusion

1) According to different sets of initial values and different sets of sample sizes we find the best estimators are Bayes 2 and Bayes 3, where the second is called  $(\hat{\lambda}_{2D})$  under De

Groot loss function, the third Bayes is obtained under precautionary loss function.

2) The comparison between four estimators depends on statistical measure mean square error,  $\left[MSE(\hat{\lambda}) = \frac{\sum_{i=1}^{n} (\lambda - \lambda)^2}{R}\right]$ , where (R) is the replicate of

each simulation experiment.

## References

 Abbas Najim Salman, Maymona M. Ameen, (2015), "Estimate the Shape Parameter of Generalize RayleighDistribution Using Bayesian – Shrinkage Technique", IJISET - International Journal of Innovative Science, Engineering & Technology, Vol. 2 Issue 6.

- [2] Abd EL-Baset A. Ahmad, Magdy E. El-Adll, Tahani A. ALOaf, (2015), "Estimation under Burr type X distribution basedon doubly type II censored sample of dualgeneralized order statistics", Journal of the Egyptian Mathematical Society 23, 391–396.
- [3] A. M.Abd-Elfattah, M. H. Abu-Moussa, (2013), "Estimation of Stress-Strength Parameter for Burr Type XIIDistribution Based on Progressive Type-II Censoring", Basic and Applied Sciences, Vol 1(1).
- [4] Al Omari Mohammed Ahmed, Noor Akma Ibrahim, JayanthiArasan and Mohd Bakri Adam, (2011),"Extension of Jeffreys's Prior Estimate For Weibull Censored Data Using Lindley's Approximation", Australian Journal of Basic and Applied Sciences, 5(12): 884-889.
- [5] Dong-Ho Jang, Myoungjin Jung, Ji-Hoon Park, and Chansoo Kim, (2014), "Bayesian Estimation of Burr Type XII DistributionBased on General Progressive Type II Censoring", Applied Mathematical Sciences, Vol. 8, No. 69, 3435 – 3448.
- [6] Essam, A. Amin, (2012), "Bayesian and Non-Bayesian Estimation of P(Y<X) from Type I Generalized LogisticDistribution Based on Lower Record Values", Australian Journal of Basic and Applied Sciences, 6(3): 616-621.
- [7] FatihKızılaslan Mustafa Nadar, (13), "Bayesian and Non – Bayesian analysis for the Burr type XII distribution based on record values and inter – record times", The 7th International DaysofStatistics and Economics, Prague, September 19-21.
- [8] Feroze, N., Aslam, M. (2012a). Bayesian Analysis of Burr Type X Distributionunder Complete and Censored Samples, International Journal of Pure and Applied Sciences and Technology, 11(2), 16-28
- [9] Feroze, N., Aslam, M. (2012c). Bayesian Analysis of Burr Type XI Distributionunder Single and Mixture of Priors, Advances in Agriculture, Sciences andEngineering Research, 2(11), 487-502.
- [10] J.A.Reshi, A.Ahmed, K.A.Mir, (2013), "Bayesian Estimation of Parameter of Size Biased Generalized RayleighDistribution Under The Extension of Jaffrey Prior And New LossFunctions", International Journal of Mathematical Research & Science (IJMRS)Vol. 1 Issue 3.
- [11] Lanping Li, (2015), "Bayes Estimation of Shape Parameter of Minimax Distribution under DifferentLoss Functions", Research Journal of Applied Sciences, Engineering and Technology 9(10): 830-833.
- [12] Nada S. Karam, Ahmed K. Jbur, (2014), "Bayesian Analyses of the Burr Type XDistribution under DoublyType II censored samples using different Priors and Lossfunctions", Mathematical Theory and Modeling, Vol.4, No.14.
- [13] Nadia H. Al-Noor and Huda A. Abd Al-Ameer (2014), "Some Estimation Methods for the Shape Parameter and ReliabilityFunction of Burr Type XII Distribution / Comparison Study", Vol.4, No.14.
- [14] NavidFeroze, Muhammad Aslamand AzharSaleem, (2014), "Bayesian Estimation and Prediction of Burr Type XI Distributionunder Singly and Doubly Censored

# Volume 6 Issue 10, October 2017

<u>www.ijsr.net</u>

## Licensed Under Creative Commons Attribution CC BY

Samples", International Journal of Hybrid Information TechnologyVol.7, No.2, pp.331-346.

- [15] Nesar Ahmad, Sabiha Khan, and M. G. M. Khan, (2013), "Planning Accelerated Life Tests for BurrType X Failure Model with Type I Censoring ",Journal of Statistical Theory and Applications, Vol. 12, No. 3, 266-287.
- [16] Nasser A. Abou-Elheggag, (2013), "Estimation for Rayleigh Distribution Using ProgressiveFirst-Failure Censored Data", J. Stat. Appl. Pro. 2, No. 2, 171-182.
- [17] Ram Kishan, (2014), "Comparison Between MLE and Bayes Estimator of Scale parameter of Generalized Gamma Distribution with Known Shape Parameters under Squared Error Loss"Journal of Reliability and Statistical Studies; ISSN (Print): 0974-8024, (Online):2229-5666 Vol. 7, Issue 1 (2014): 43-50.
- [18] SankuDey, TanujitDey, and Sudhansu S. Maiti, (2015), "Bayes Shrinkage Estimation of the Parameter ofRayleigh Distribution for Progressive Type-IICensored Data", Austrian Journal of Statistics, Volume 44, 3–15.
- [19] TabassumNaz Sindhu, NavidFerozeb, and Muhammad Aslam, (2015), "Analysis of doubly censored Burr type IIdistribution: a Bayesian look", Electronic Journal of Applied Statistical AnalysisVol. 08, Issue 02, 154-169.
- [20] TabassumNaz Sindhu, NavidFeroze, and Muhammad Aslam, (2014), "Preference of Prior for Bayesian Analysis of theMixed Burr Type X Distribution under Type I Censored Samples", Pak.j.stat.oper.res. Vol.X No.1, pp17-39.
- [21] T. N. Sindhu, M. Aslam, (2014), "On the Parameter of the Burr Type X under Bayesian Principles", International Journal of Mathematics, Vol.8, No.1.
- [22] Yarmohammadi M. and Paziar H. (2010), "Minimax Estimation of the parameter of the Burr type XII distribution", Australian journal of basic and applied sciences, 4(12):6611 – 6622.
- [23] ZouaouiChikr el-Mezouar, (2010), "Estimation the Shape, location and Scale Parameters of the Weibull", RT&A, No. (04) (19)Vol.1. Principles", International Journal of Mathematical, Computational, Physical, Electrical and Computer Engineering Vol:8, No:1.