

Technical Sheet of Contribution to the Study and Valorization of the Therapeutic Potential of Pepper *Capsicum frutescens var. soudanais* used in the Traditional Pharmacopoeia in Côte d'Ivoire

Yao Kouadio¹, Dan CG², Kouassi KC³, Gonnéty TJ⁴

^{1,2,4}Department of Food Science and Technology, University of Nangui Abrogoua, Laboratory of Biocatalysis and Bioprocessing, 02 BP 801 Abidjan 02, Côte d'Ivoire

³Universities Jean Lorougnon Guede, Unit Training and Agroforestry Research, Laboratory of Microbiology, Bio-industry and Biotechnology BP150 Daloa, Côte d'Ivoire

Abstract: *Capsicum frutescens var. soudanais* is one of the most used peppers, especially in traditional medicine by healers to treat various infections and pathologies in Côte d'Ivoire. The objective of this study was to evaluate the nutritional and therapeutic potential of *Capsicum frutescens var. soudanais* (*C. frutescens var. soudanais*) to help show consumers and consumers the benefits of consuming this widely used vegetable. The determination of beta carotene (vitamin A) and ascorbic acid (vitamin C) (antioxidants) was carried out using standard extraction methods, thin layer chromatography (TLC) and high performance liquid chromatography (HPLC). The various extracts (petroleum ether, acetone, dichloromethane and methanol) showed contents ranging from 68.47 to 535.98 µg in beta-carotene and from 86.38 to 96.62 mg of ascorbic acid (antioxidants) per 100 g of fruits of *C. frutescens var. soudanais*. These extracts were also rich in bioactive compounds such as steroids, terpenes, flavonoids, polyphenols and quinones which are molecules endowed with preventive powers of various diseases, anti-inflammatory, analgesic, antimicrobial and strengthening of the immune system. Thus, the consumption and use of peppers *C. frutescens var. soudanais* in the treatment of certain diseases and infections of microbial origin would be beneficial because they could contribute more to the nutritional and health needs of consumers.

Keywords: *Capsicum frutescens var. soudanais*; Bioactive compounds; Beta-carotene; Ascorbic acid; Antioxidant; Abidjan

1. Introduction

The increasing growth of microbial infections and food-borne diseases encourage researchers and scientists to seek out new bioactive molecules that may be secondary metabolites such as phenolic compounds, terpenoids, steroids and alkaloids [1, 2]. Fruits and vegetables are important sources of bioactive compounds and antioxidants known for their potential effects on human infections and degenerative diseases [3, 4, 5].

The *Capsicum* pepper of the Solanaceae family is a popular vegetable of South American and central tropical agricultural crops, containing the species *Capsicum annum*, *Capsicum frutescens*, *Capsicum baccatum*, *Capsicum pubescens* and *Capsicum chinens* [2, 5]. The pepper is grown for economic needs but especially for its flavor, aroma, nutritional and sanitary effects [5, 6, 7]. The fruits of pepper can be used and eaten at different stages of maturation : unripe (green), ripe (red or yellow) or not too ripe.

In Côte d'Ivoire, many varieties and species of pepper are grown. However, the species *Capsicum annum* and *Capsicum frutescens* are widely known for their consumption, flavor, burning sensation or pungent effects conferred by components such as capsaicinoids and their wide range of applications in traditional therapies [5, 7, 8]. Fruits of the genus *Capsicum* are rich in carotenoids (Vitamins A) and antioxidants (Vitamin C). Carotenoids function as antioxidants at low oxygen pressure and can

protect against damage caused by free radicals and peroxidation [8, 10]. Carotenoids also play an important role in the elimination of free radicals, the prevention of certain types of cancers, cardiovascular diseases, ocular vision disorders, skin degeneration and aging [5].

The fruits of *Capsicum frutescens var. soudanais* (*C. frutescens var. soudanais*), in Côte d'Ivoire, are used extensively by traditional healers and households to treat infections and post-partum care through rectal or oral colon enemas.

The aim of this study is to show the bioactive compounds and vitamins with antioxidant activities of ripe and dry fruits, as well as the potential therapeutic and antimicrobial effects of the species of *Capsicum frutescens var. soudanais* used in Côte d'Ivoire in traditional therapy.

2. Material and Methods

2.1 Plant material

The plant material, that is to say, the fresh, dried and rendered fruits of *C. frutescens var. soudanais* was collected on four local markets in Abobo, Adjamé, Treichville and Koumassi communes in Abidjan, Côte d'Ivoire (Figure 1). The samples of *C. frutescens var. soudanais* were transported fresh to the laboratory and then identified by the National Floristic Center (Felix Houphouët-Boigny University, Abidjan-Côte d'Ivoire).



Figure 1: Organs of the pepper *C. frutescens var. soudanais* a) plant, leaves and fruits non-ripe (green color) ; (b) plant, leaves and fruits ripe (red color) ; c) fruits ripe and dry ; d) Packaging 100 g of ripe, dried fruit

The species of peppers of *C. frutescens var. soudanais* has several names according to certain ethnic groups in Côte d'Ivoire (Table 1).

Table 1: Different names of pepper of *C. frutescens var. soudanais* by ethnic group in Côte d'Ivoire

| Ethnic group | Local name |
|--------------|----------------|
| Abè | Makou |
| Abouré | Aissa mouklou |
| Abron | Téyssan |
| Agni | Mocloua |
| Baoulé | Boharé mancoun |
| Malinké | Frotofiè |

2.2 Methods

Extracts or mixtures of *C. frutescens var. soudanais* in traditional medicine are generally prepared according to the method described above (Figure 2). Fresh or dried ripe fruits of *C. frutescens var. soudanais* (10 to 30 g sometimes) are rinsed with potable water or soaked for a few minutes. Then the rehydrated fruits are crushed using a pebble (with the possibility of adding other ingredients such as leaves to therapeutic virtue). The paste obtained is suspended with warm water or cold water. It is also possible to slightly warm the mixture, an empirical form of pasteurization and increase the extraction of bioactive compounds. Finally, an enema bulb is used to administer the mixture rectally (Figure 2).

The research and the determinations of the bioactive molecules of the pepper *C. frutescens var. soudanais* were carried out by thin layer chromatography and then by high performance liquid chromatography (HPLC). The various extracts studied were obtained with solvents such as petroleum ether, dichloromethane, acetone and methanol. The beta-carotene of the sample was extracted according to the method described in [9, 12] and the ascorbic acid was extracted according to the method described in [13, 14].




Figure 2: Standard traditional method of preparing the suspension of *C. frutescens var. soudanais* for enema

3. Results and Discussion

Fruits of pepper *C. frutescens var. soudanais* are used in the traditional pharmacopoeia and in the human food in different forms of:

- Fresh fruit to maturity, ripe or unripe, cut or rendered in paste,
- Dried ripe fruit.

On the local markets, the fruits of *C. frutescens var. soudanais* are sold in dried form. However, the bag of 15 kilograms of dried fruits of *C. frutescens var. soudanais* is sold in bulk, at an estimated price of 25,000 F. CFA. Retail prices are shown in (Table 2).

Table 2: Economic and food value of stems sold

| Price of the tide | Mass (g) |
|-------------------|----------------------|
| 100 f. CFA | 20 g |
| 500 f. CFA | 100 g |
| 25 000 f. CFA | 15 Kg (15 000 g.) |

Bioactive compounds in varieties *C. frutescens var. soudanais*

Chromatographic processing methods have revealed many bioactive compounds. Total bioactive compounds were determined in the analyzed samples (Table 3). The extracts of *C. frutescens var. soudanais* analytes contain beta-carotenes, ascorbic acid and metabolites such as steroids, terpenes, flavonoids, polyphenols and quinones.

The levels of beta carotenes (pro vitamin A) and ascorbic acid (vitamin C) of *Capsicum frutescens var. soudanais* varied between 68.47 to 535.98 µg of beta-carotene and 86.38 to 96.62 mg of ascorbic acid per 100 g of fruit respectively.

Table 3: Bioactive metabolites from *C. frutescens* var. *soudanais* identified by HPLC

| Extraction solvents used | Bioactive metabolites identified from the variety of <i>Capsicum frutescens</i> var. <i>soudanais</i> | | | | | | |
|--------------------------|---|----------|---------|----------|----------|------------|-------------|
| | Alkaloids | Quinones | Tannins | Stéroids | Terpenes | Flavonoids | Polyphénols |
| Petroleum ether | + | + | - | + | + | + | + |
| Dichloro-methane | + | + | - | + | + | + | + |
| Acetone | + | + | - | + | + | + | + |
| Methanol | + | + | + | + | + | + | + |

+ : presence of the title compound ; -: absence of the compound title

Carotenoids behave like antioxidants and could protect the tissues against damage caused by free radicals and also intervene in the prevention of cardiovascular diseases, cancer, ophthalmic diseases and in the fight against aging [10]. As for ascorbic acid, it strengthens the immune system and plays a role in the state of health [11].

The presence of different bioactive metabolites would explain the wide use of the peppers of the genus *Capsicum frutescens* var. *soudanais* in the traditional Ivorian pharmacopoeia to treat many intestinal disorders and diseases such as gastroenteritis and also prevent infections in women after childbirth.

Apart from its use in oral feeding, the main route of administration in traditional therapy is the rectal route by suppositories or enemas. Indeed, bioactive compounds such as tannins, steroids, polyphenols, terpenes and quinones play a preventive role in diseases and possess antimicrobial activity [18]. Fruit consumption of *C. frutescens* var. *soudanais* would be beneficial for health as tannins would play an important role in the treatment and prevention of certain degenerative diseases such as cancer [3]. Alkaloids act as an anti-inflammatory, analgesic and also fight against asthma [15] while flavonoids in the diet also reduce the risk of cancers and heart disease as well as the symptoms of menopause [5,16, 17].

4. Conclusion

It is clear from this study that *Capsicum frutescens* var. *soudanais* is a major source of potential for bioactive compounds. The results obtained are important for informing actors (traditional healers) and promoters of traditional medicine in Côte d'Ivoire.

The study of the antimicrobial activity of extracts of bioactive compounds of this vegetable could contribute to the knowledge of new molecules derived from phytotherapy which is increasingly topical to help fight against the resistance of microorganisms to antibiotic therapy uncontrolled.

References

[1] Hervert-Hernandez D, Sayago-Ayerdi SG, Goni I (2010). Bioactive compounds of four hot pepper varieties (*Capsicum annum* L.) : antioxidant capacity, and intestinal bio accessibility. *Journal of Agricultural and Food Chemistry*, 58 : 3399-3406

[2] Zimmer AR, Leonardi B, Miron D, Schapoval E, Oliveira JR, Gosmann G (2012). Antioxidant and anti-inflammatory properties of *Capsicum baccatum*: From

traditional use to scientific approach. *Journal of Ethnopharmacology*, 139 (1) : 228-233

[3] Li F, Awale S, Tezuka Y, Kadota S (2008). Cytotoxic constituents from Brazilian red propolis and their structure-activity relationship. *Bioorganic & Medicinal Chemistry*, 16 : 5434-5440

[4] Meghvansi MK, Siddiqui S, Khan MH, Gupta VK, Vairale MG, Gogoi HK, Singh L (2010). Naga chilli: a potential source of capsaicinoids with broad-spectrum ethnopharmacological applications. *Journal of Ethnopharmacology*, 132 : 1-14

[5] Kouassi K C, Nanga YZ, Lathro SJ, Aka S, Koffi-Nevry R (2012). Bioactive compounds and some vitamins from varieties of pepper (*Capsicum*) grown in Côte d'Ivoire. *Pure Appl. Bio.*, 1(2) : 40-47.

[6] Cruz-Perez AB, Gonzalez-Perez VA, Soto Hernandez RM, Gutierrez-Espinosa MA, GardeaBéjar AA, Perez-Grajales M (2007). Capsainoid, vitamin C and heterosis during fruit development of Manzano hot pepper. *Agrociena*, 41: 627-635

[7] Kouassi KC, Koffi-Nevry R (2012). Evaluation de la connaissance et utilisation des variétés de piment (*Capsicum*) cultivées en Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*, 6 (1): 175-185

[8] Kollmannsberger H, Rodriguez-Burruezo A, Nitz S, Nuez F (2011). Volatile and capsaicinoid composition of aji (*Capsicum baccatum*) and rocoto (*Capsicum pubescens*), two Andean species of chile peppers. *Journal of the Science of Food and Agriculture*, 91: 1598-1611

[9] Guzman I, Hamby S, Romero J, Bosland PW, O'Connell MA (2010). Variability of Carotenoid Biosynthesis in Orange Colored *Capsicum* spp. *Plant Science*, 179 (1-2) : 49-59

[10] Rodriguez-Burruezo A, Gonzalez-Mas Mdel C, Nuez F (2010). Carotenoid composition and vitamin A value in aji (*Capsicum baccatum* L.) and rocoto (*C. pubescens* R. & P.), 2 pepper species from the Andean region. *Journal of Food Science*, 75: 446-453

[11] Walingo K.M. (2005). Role of vitamin C (ascorbic acid) on human health-a review. *African Journal of food, Agriculture, Nutrition and Development*, 5:1

[12] Amin I, Cheah SF (2003). Determination of vitamin C, β carotene and riboflavin contents in five green vegetables organically and conventionally grown. *Malaysian Journal of Nutrition* 9 (1) : 31-39

[13] Albuquerque B, Lidon CF, Leitao EA (2005). Ascorbic acid quantification in melon samples : the importance of the extraction medium for HPLC analysis. *General and Applied Plant Physiology*, 31 (3-4): 247-251

[14] Deepa N, Kaur C, Singh B, Kapoor HC (2006). Antioxidant activity in some red sweet pepper cultivars. *Journal of Food Composition and Analysis*, 19: 572-578

- [15] Staerk D, Lykkeberg AK, Christensen J, Budnik BA, Abe F, Jaroszewki JW (2002). In vitro cytotoxic activity of phenanthroindolizidine alkaloids from *Cynanchum vincetoxicum* and *Tylophora tanake* against drug-sensitive and multidrug-resistant cancer cells. *Journal of Natural Products*, 65 (9): 1299-1302
- [16] Ferguson LR (2001). Role of plant polyphenols in genomic stability. *Mutation Research*, 475 : 89111
- [17] Xu HX, Wan M, Dong H, But PP, Foo LY (2000). Inhibitory activity of flavonoids and tannins against HIV-1 protease. *Biological and Pharmaceutica Bulletin*, 23: 1072-1076
- [18] Hassan SW, Umar RA, Lawal M, Bilbis LS, Muhammad BY, Dabai YU (2006). Evaluation of antibacterial activity and phytochemical analysis of root extracts of *Boscia angustifolia*. *Journal of Biotechnology*, 5 (18) : 1602-1607

