

Pre-Harvest Treatments for Fruit Quality Improvement in Rainy Season Guava (*Psidium guajava*)

Swosti Debapriya Behera¹, Sanjay Pathak²

Department of Horticulture, College of Horticulture and Forestry, Narendra Deva University of Agriculture & Technology, Kumarganj, Faizabad, Uttar Pradesh, India (224 229)

Abstract: Rainy season or Ambe bahar guava crop gives low quality fruits having poor shelf life. The quality is poor due to infestation by insect pest like fruit fly and infection by disease like anthracnose. Various pre-harvest treatments viz. spraying of CaCl₂ 2%, spraying of CaSO₄ 2%, polythene bagging, brown paper bagging, spraying of CaCl₂ 2% + polythene bagging, spraying of CaCl₂ 2% + brown paper bagging, spraying of CaSO₄ 2% + polythene bagging, spraying of CaSO₄ 2% + brown paper bagging were tried at Faizabad, Uttar Pradesh, India during 2015-16 to find out ways for improving fruit quality. All treatments increased quality of guava fruits significantly. Among the treatments, CaCl₂ 2% + polythene bagging proved the best in enhancing post harvest quality attributes viz. fruit size, fruit weight, fruit firmness, organoleptic quality, total soluble solids (TSS), acidity, TSS: acid ratio, ascorbic acid, reducing sugar, non reducing sugar and total sugar.

Keywords: Ambe bahar, calcium chloride, calcium sulphate, polythene bag, post-harvest quality

1. Introduction

Guava (*Psidium guajava* Linn.) is one of the most common fruits in India. Among the three fruiting seasons of the crop, fruits harvested in rainy season are insipid, watery, poor in quality, attacked mostly by diseases and pests. Anthracnose (*Gloeosporium psidii*) is one of major disease that adversely affects quality of fruits in this season. Infestation with fruit fly (*Dacus dorsalis*) has been a major impediment to guava marketing. Keeping quality of ambe bahar guava fruits is very poor. Hence fruit production during this season is often minimized or removed by crop regulation or bahar treatment methods. It is a big hurdle in the annual production of guava. Therefore, to resolve all these problems a proper management programme should be incorporated.

2. Literature Survey

Several attempts have been carried out by various research workers to overcome this problem. Bagging is an effective eco-friendly and non-chemical method for the control of post harvest diseases and fruit fly infestation. It also improves the fruit quality, organoleptic quality in terms of appearance, uniform coloration, taste, flavour and overall acceptance at ripening and also prolongs the shelf life as reported in rainy season guava (Abbasi *et al.*, 2). Bagging may produce spot free, attractive and high quality fruit at harvest on ripening leading to export and better price for guava growers. Calcium plays an important role in a number of physiological and biochemical processes in the plant concerning membrane structure, function and enzyme activities. It helps in retaining fruit firmness, reducing respiration, decreasing storage breakdown, rotting and browning (Jakhar *et al.*, 8). It helps in creating a protecting layer on fruit surface which protects against pathogens and insects. Calcium nitrate is not useful for spraying at mature stage of guava fruits. Hence the present investigation was formulated with pre-harvest CaCl₂ and CaSO₄ spray and/or

fruit bagging with polythene bag and brown paper bag to enhance quality and appearance of rainy season guava fruits.

3. Materials and Methods

The experiment was carried out at the guava orchard at Main Experiment station of Horticulture and Post Harvest Technology laboratory, Department of Post Harvest Technology, College of Horticulture and Forestry, Narendra Deva University of Agriculture & Technology, Kumarganj, Faizabad (U.P.), India during the *kharif* season of 2015-16. Four year old bearing trees of guava cv. 'Lucknow-49', having uniform vigour and healthy fruits were selected for the study. The trees were spaced at 8m x 8m spacing and uniform crop management practices were followed for all the plants. The nine pre-harvest treatments viz., T₁ (CaCl₂@ 2%), T₂(CaCl₂ @ 2% + Polythene bag), T₃(CaCl₂ @ 2% + Brown Paper bag), T₄(CaSO₄ @ 2%), T₅ (CaSO₄ @ 2% + Polythene bag), T₆ (CaSO₄ @ 2% + Brown Paper bag), T₇ (Polythene bag), T₈ (Brown Paper bag) and T₉ (Control) were tried in randomized block design with three replications. Single tree was considered as an experimental unit. The nine treatments in each block were randomised. Altogether there were twenty seven trees. A total of 10 uniform sized fruits/tree present in all directions of tree canopy were selected and tagged for the study. Bagging of fruits with various bags was done about 30 days before harvesting of fruits. Spraying of CaCl₂ @ 2% alone in T₁ and CaSO₄ @ 2% alone in T₄ were done twice at 15 and 30 days before harvest of fruits. Bagging of fruits with polythene bag alone in T₇ and brown paper alone in T₈ were done one month before harvest of fruits. Bagging of fruits with polythene and brown paper bags was done in T₂ and T₃, respectively, in addition to single spraying with CaCl₂@ 2% 30 days before harvesting. Similarly, in case of T₅ and T₆, bagging with polythene bag and brown paper bag was done, respectively, in addition to single spraying with CaSO₄ @ 2%. In case of treatments where spraying and bagging both

were combined, the bagging was done immediately after the spraying.

Fruits marked for study in all treatments were separately harvested by hand carefully to avoid any damage to fruits in morning hours on 1st August, 2015. The field heat of harvested fruits was reduced by showing to stream of hand pumped water and kept in shade for drying of water sticking to surface of fruits. These fruits were transported from orchard to Post Harvest Technology laboratory with proper packing in CFB boxes to avoid physical damage including bruising. Individual weight of 4 fruits per treatment was recorded by physical balance. Length and width were measured by using Vernier callipers. Two fruits of each treatment were separately packed in polythene bags and bags were sealed properly. Six to eight small holes were made in polythene bags for air exchange. All bags were marked as per treatments and then stored at ambient temperature in the laboratory of Post Harvest Technology for further observations on changes during storage.

The fruit size was measured using Vernier callipers. Fruit weight (g), physiological loss in weight and spotted fruits were recorded using the standard procedure mentioned in (AOAC,1). The TSS(%) was determined with hand refractometer. Acidity (%) and ascorbic acid (mg/ 100g) were determined by the standard procedures. Fehling solution method was used for estimating sugar content. The TSS: acid ratio was calculated by dividing the TSS (%) to acidity (%) of fruits. The organoleptic evaluation for assessing sensory attributes of stored fruits was conducted by a panel of eight judges and samples were rated on the nine-point Hedonic Rating Scale.

4. Results and Discussion

4.1 Fruit Size

Among all treatments, application of CaCl₂@2% + polythene bag proved the best with fruit length and width of 4.8 cm and 5.1 cm, respectively (Table 1). The minimum fruit length and width of 3.5 cm and 3.8 cm, respectively, were obtained in control. Improvement in fruit size with application of CaCl₂ was reported in mango by Rubyrani and Brahmachari(10). Calcium plays a major role in cell elongation due to which size of fruit increases.

4.2 Fruit Weight

The treatment of CaCl₂@ 2%+ Polythene bag produced the heaviest fruit with average fruit weight of 155.33 g. The minimum fruit weight of 100.3g was recorded in control. The appreciable improvement in fruit weight in mango with application of CaCl₂ has been reported earlier by Rubyrani and Brahmachari(10).

4.3 Fruit firmness

The highest firmness was recorded in treatment of CaCl₂@ 2%+ Polythene bag (10.48 kg/cm²) followed by the treatment of CaSO₄ @ 2%+ Polythene bag (9.53 kg/cm²). The lowest fruit firmness (7.97 kg/cm²) was recorded in control. Several workers have also reported earlier regarding

the effect of pre harvest application of Ca⁺ on improvement in firmness of fruits (Tingwa and Young,15 ; Scott and Wallis,11; Singh *et al.*,12; Cheor *et al.*, 3).

4.4 Spotted fruit (%)

The minimum spotted fruits were noted with the treatment of CaCl₂@2% + Polythene bag(1.70%) followed by CaSO₄ @2%+ Polythene bag(5.30%).The maximum spotted fruits were noted in control(65.33%). Spots on fruits occur mainly due to high humidity condition creating congenial conditions for pathogen attack. This can only be minimised by protecting the fruits from adverse environment. Bagging creates a microenvironment safe for fruits and also avoids pathogen attack and spot formation.

4.5 Infested fruit (%)

No infested fruit was found with the pre harvest treatments of polythene bag alone and its combination with spraying treatments i.e. CaCl₂@ 2% + Polythene bag and CaSO₄ + Polythene bag. The highest infested fruits were recorded in control (13.5%). Infection with disease like anthracnose and infestation with insect pest like fruit fly mainly occur during rainy season due to availability of congenial micro-climate. Dieng *et al.* (5) have reported that the incidence of fruit fly infestation was observed to the tune of 70% on unbagged fruits as against 0.45% in bagged fruits.

4.6. Organoleptic quality

Calcium chloride+ polythene bag recorded the maximum organoleptic quality as indicated by the value of 8.80 (like very much) followed by 8.50 (like very much) with calcium chloride + brown paper bag. The minimum organoleptic quality with value of 6.20 (like slightly) was found in control. Abbasi *et al.* (2) reported that bagging of guava fruits in summer helped in improving organoleptic taste.

4.7 TSS content

The maximum TSS content was recorded in fruits subjected to treatment of CaCl₂@2% + Polythene bag (15.0 %) followed by the fruits treated with CaCl₂@2% + brown paper bag (14.5%) and the minimum TSS content was found in control (Table 2). The TSS content of fruits increased during the ripening and storage due to hydrolysis of insoluble starch into soluble sugars and loss of moisture (Koksal *et al.*, 9). Similar improvement in TSS content of fruits with pre harvest spray of calcium chloride was reported by Singh *et al.* (13) in Amrapalli mango and Goswami *et al.* (7) in Sardar guava.

4.8 Acid content

Rainy season guava are low in acid content because of leaching loss which causes very poor taste in fruits. It can be controlled by protecting fruits from such losses due to heavy rain. The maximum acid content was noted in treatment of CaCl₂@2% + Polythene bag (0.39%) because it creates a protective layer against leaching loss. The minimum was found in control (0.29%).

4.9 TSS: acidity ratio

The highest TSS: acidity ratio 39.67 was found in CaCl₂@2% + brown paper bag treatment followed by 38.26 in CaCl₂@2% + polythene bag treatment. The minimum was found in control (30.72).

4.10 Ascorbic acid content

The maximum ascorbic acid content was recorded in pre harvest treatment of CaCl₂@2% + Polythene bag(211.67mg/100g). The minimum ascorbic acid content was reported in control (113.33 mg/100g). This result conforms to the findings of Dhahiya *et al.*(4) in guava, Singh *et al.*(14) in ber fruits and Dutta (6) in guava.

4.11 Total sugars content

The pre harvest treatment of CaCl₂@2% + Polythene bag with total sugars content of 11.18% was found significantly superior to all other treatments(Table 3). The minimum total sugar content was observed in control (6.95%). Similar improvement in the total sugar content of fruits by pre harvest spray of calcium chloride was reported by Singh *et al.*(13) in Amrapali mango.

4.12 Reducing sugar content

The highest reducing sugar content was recorded in pre harvest treatment of calcium chloride 2%+ polythene bag (6.04%), while the lowest was recorded in control (3.80%). Similar improvement in the reducing sugar content of fruits by pre harvest spray of calcium chloride was reported by Singh *et al.* (13) in Amrapali mango and Goswami *et al.*(7) in guava fruits.

4.13. Non-reducing sugar content

The maximum non-reducing sugar content in fruits was recorded in pre harvest treatment of calcium chloride 2%+ polythene bag (5.23%), while the minimum was recorded in control (3.15%). Similar increase in non-reducing sugars content of fruits by pre harvest spray of calcium chloride have been reported by Singh *et al.*(13) in Amrapali mango and Goswami *et al.*,(7) in guava.

5. Conclusion

All the pre harvest treatments were found better than control in improving fruit quality while the pre harvest treatment of CaCl₂ 2% + polythene bag was found superior to increase the fruit quality than all other treatments. This treatment was found to have very low spots and no infestation. Hence it should be practised in rainy season guava crop to produce fruits with good quality.

6. Future Scope

Study of several bagging materials, calcium chloride and calcium sulfate will help for developing a good sustainable research programme as the materials and chemicals are non-

hazardous both for environment and human. It will provide a good rainy season guava crop to fruit growers and traders.

References

- [1] A.O.A.C 2000. Official methods of analysis. 17th edition, Association of Official Analytical Chemists, Washington, D.C. USA.
- [2] Abbasi, N. A., Chaudhary, M.A., Ali, M.I., Hussain A. and Ali, I. 2014. On tree fruit bagging influences quality of guava harvested at different maturity stages during summer. *International J. Agric. and Biol.*, **16** (3) : 543-49
- [3] Cheor, F., Wallemot, C., Arul, J., Desjardins, Y., Makhlauf, J., Charest, P.M. and Gosselin, A. 1990. Foliar application of calcium chloride delays post harvest ripening of strawberry. *J. Am. Soc. Hort. Sci.*, **115**: 789-92.
- [4] Dhahiya, D.K., Singh, S.K., and Singh, A. 2001. Studies on the foliar treatments on yield and quality of guava. *Acta Hort*, **628**: 121-127.
- [5] Dieng, E.O., Nadiaye, S., Diedhiou, P.M., Baol, A.B., Mbaye, N., Vaughan, L. 2012. Evaluation of the efficacy of mango fruit bagging on fruit flies punctures, economic profitability and impact on anthracnose and fruit quality in Senegal Niayes zones. *Acta Hort.*, **886**: 160-66.
- [6] Dutta, P. 2004. Studies on foliar nutrition in guava. *The Orissa J. Hort.*, **32**(1): 103-04.
- [7] Goswami, A.K., Sharma, R.M. and Kumar, R. 2008. Effect of pre and post harvest treatment of calcium on quality and shelf life of guava fruit cv. Sardar. *Acta Hort*, **682**: 1281-86.
- [8] Jakhar, M.S. and Pathak, S. (2014). Enhancing quality of mango (*Mangifera indica* L.) fruits cv. Amrapali with pre-harvest foliar spray and fruit bagging, *Annals of Agri-Bio Research*. **19**(3): 488-491.
- [9] Koksai, A.L., Dumanoglu, H. And Tuna, N.1994. The effect of Semper fresh on the storage of 'Williams' pear and 'Starkspur Golden Delicious' apple cultivars. *Acta Hort.*, **368**: 793-01.
- [10] Rubirani and Brahmachari, V.S. 2004. Effect of growth substances and calcium compound on fruit retention, growth and yield of Amrapali mango. *The Orissa J. Hort.*, **32**(1): 15-18.
- [11] Scott, K.J. and Wallis, R.B.H. 1977. Vacuum infiltration of calcium chloride: A method for reducing bitter pit and senescence of apples during storage at ambient temperature. *Hort. Sci.*, **12**: 53-57.
- [12] Singh, B.P., Singh, H.K. and Chauhan, K.S. 1981. Effect of post-harvest calcium treatments on the storage life of guava fruits. *Indian j. Agric. Sci.*, **51**: 44-47.
- [13] Singh, S., Brahmachari, V.S. and Jha, K.K. 1998. Effect of calcium and polythene wrapping on storage life of mango. *Indian J. Hort.*, **55**(3):218-22.
- [14] Singh, S., Singh, A.K. and Bagel, B.G.2002. Effect of foliar application of nutrients and growth regulators on quality and storage life of *ber* fruits. *The Orissa J. Hort.*, **36**(1): 126-30.
- [15] Tingwa, P.O. and Young, R.E. 1974. The effect of calcium on ripening of avocado (*Persia americana* Mill) fruits. *J. Am. Soc. Hort. Sci.*, **99**:540-42.

Table 1: Effect of pre-harvest treatments on fruit size, weight, firmness, spotted fruits, infested fruits and organoleptic quality

Treatments	Fruit length (cm)	Fruit width (cm)	Fruit weight (g)	Fruit firmness (Kg/cm ²)	Spotted fruits (%)	Infested fruits (%)	Organoleptic quality
T ₁ - CaCl ₂ 2%	3.6	4.0	108.6	8.57	(11.00)* 2.78	(4.95)* 2.33	7.8
T ₂ - CaCl ₂ 2% + Polythene bag	4.8	5.1	155.3	10.48	(1.70) 1.48	(0.00) 0.71	8.8
T ₃ - CaCl ₂ 2% + Brown Paper bag	4.5	4.5	139.6	9.40	(14.00) 3.80	(2.46) 1.58	8.5
T ₄ - CaSO ₄ 2%	3.5	3.9	105.0	8.62	(16.67) 4.14	(6.75) 2.61	7.0
T ₅ - CaSO ₄ 2% + Polythene bag	4.6	4.5	140.0	9.53	(5.30) 2.37	(2.46) 1.58	8.0
T ₆ - CaSO ₄ 2% + Brown Paper bag	4.3	4.4	133.3	9.30	(17.67) 4.24	(6.75) 2.61	7.8
T ₇ - Polythene bag	4.0	4.2	114.3	9.31	(8.17) 2.94	(0.00) 0.71	7.8
T ₈ - Brown Paper bag	3.8	4.0	110.3	8.50	(20.33) 4.56	(9.5) 3.16	7.4
T ₉ - Control	3.5	3.8	100.3	7.97	(65.33) 8.09	(13.5) 3.74	6.2
SEm±	0.1	0.1	4.72	0.15	0.19	0.01	0.15
CD at 5%	0.4	0.4	14.14	0.44	0.58	0.04	0.44

*√(X+0.5) transformed values. Original values in parentheses

Table 2: Effect of pre-harvest treatments on TSS, acidity, TSS : Acid ratio and ascorbic acid content

Treatments	TSS (%)	Acidity (mg/100g)	TSS:Acid ratio	Ascorbic acid (mg/ 100g)
T ₁ - CaCl ₂ 2%	12.3	0.34	36.60	175.00
T ₂ - CaCl ₂ 2% + Polythene bag	15.0	0.39	38.26	211.67
T ₃ - CaCl ₂ 2% + Brown Paper bag	14.5	0.37	39.67	183.00
T ₄ - CaSO ₄ 2%	11.0	0.33	33.38	164.64
T ₅ - CaSO ₄ 2% + Polythene bag	12.7	0.36	35.33	200.33
T ₆ - CaSO ₄ 2% + Brown Paper bag	12.0	0.35	34.65	151.67
T ₇ - Polythene bag	12.4	0.33	37.54	140.33
T ₈ - Brown Paper bag	11.2	0.31	36.20	131.67
T ₉ - Control	9.0	0.29	30.72	113.33
SEm±	0.19	0.01	1.16	1.70
CD at 5%	0.56	0.04	3.46	5.11

Table 3: Effect of pre-harvest treatments on reducing sugars, non-reducing sugars and total sugars

Treatments	Reducing sugars	Non-reducing sugars	Total sugars
T ₁ - CaCl ₂ 2%	4.91	4.31	9.22
T ₂ - CaCl ₂ 2% + Polythene bag	6.04	5.23	11.18
T ₃ - CaCl ₂ 2% + Brown Paper bag	5.82	5.07	10.88
T ₄ - CaSO ₄ 2%	4.43	3.85	8.28
T ₅ - CaSO ₄ 2% + Polythene bag	5.00	4.40	9.40
T ₆ - CaSO ₄ 2% + Brown Paper bag	4.83	4.23	8.87
T ₇ - Polythene bag	4.90	4.39	9.32
T ₈ - Brown Paper bag	4.53	3.93	8.44
T ₉ - Control	3.80	3.15	6.95
SEm±	0.12	0.21	0.31
CD at 5%	0.36	0.62	0.92

Author Profile



Swosti Debapriya Behera received B.Sc. (Hort.) degree and M.Sc. (Ag.) Horticulture degree in 2014 and 2016, respectively from Orissa University of Agriculture and Technology and Narendra Deva University of Agriculture and Technology, respectively. Currently she is working as a Ph.D. Research scholar in the department of Horticulture, Assam Agricultural University, Jorhat-13.



Dr. Sanjay Pathak : Professor and Head of Post harvest Technology, Department of Horticulture, Narendra Deva University of Agriculture and Technology, Faizabad, U.P.