

# Prevalence of Anaemia by Evaluating Various Haematological Parameters in Medical Students

Dr Vijay B S Gaur<sup>1</sup>, Dr Bindu Singh Gaur<sup>2</sup>

<sup>1</sup>Professor Physiology, HIMS Mau, Ataria, Sitapur

<sup>2</sup>Professor Pathology, LNMC Bhopal

**Abstract:** ***Objective:** To find out the prevalence of anaemia in medical students by evaluating their Haemoglobin content, PCV/hct, TRBC count and Absolute Indices. **Methods:** The present work has been undertaken on 150 cases between age group of 18 to 24 years. Two ml blood was taken from ante-cubital vein by vene-puncture by using 24-gauge needle under aseptic conditions using a disposable syringe. This blood was poured in EDTA vials and was used for the haematological investigation. **Summary:** The present study was undertaken in 150 medical students, which included 93 males and 57 females. The entire series was subjected to haematological investigation for anaemia prevalence. **Conclusion:** The overall prevalence of anaemia in our study was found to be 9.3%. But since the anaemic subjects were exclusively females it can also be concluded that the prevalence was zero percent in males and 24.56% in females.*

**Keywords:** Prevalence, Anaemia, Haematological parameters, Medical students

## 1. Introduction

Anaemia of various types affect large population groups in most of the developing countries. Therefore, it is reasonable to believe that anaemia impairs health and working capacity leading to mental, physical and economical loss. Anaemia is one of the major problems of medical practice in our country. Prevalence of anaemia in our country has been reported as being as low as 0.8% to 97.7% in various population. WHO sponsored studies indicate that anaemia is considered to be the late manifestation of nutritional deficiency believed to be result of malnutrition, chronic blood loss with particular attention to the relative roles of dietary deficiency such as iron, folic acid, vitamin B-12 in the pathogenesis of anaemia. In our country as reported by WHO and FAO dietary analysis of food consumed by anaemic person indicate that intake of iron is within recommended allowance but widespread anaemia is associated with less absorption and assimilation of iron or excessive losses. Anaemia due to iron deficiency is associated with many disabilities including learning capabilities. It is very essential that the exact prevalence of anaemia be studied. The present study aims at finding the same among medical students.

## 2. Prevalence

Iron deficiency is widespread throughout the world. It afflicts persons of all ages and economic groups, although it is more common among the very young, among those on poor diets and among women. Indeed, the worldwide prevalence of iron deficiency anaemia is such as to rank it among the commonest chronic maladies of humankind. Data of the WHO show that anaemia due principally to iron deficiency, affects approximately 30% of the world's population, or about 1.3 billion persons, and that approximately 43% of preschool children, 51% of pregnant women and 37% of school age children are anaemic. In parts of the world where there is a high frequency of intestinal helminthiasis and the population subsists on an iron poor diet, iron deficiency may be almost the norm. The screening

procedure used most often is simple blood haemoglobin determination. In such surveys, the usual procedure is to set a cut-off value on the basis of data from a normal population; values falling below the cut-off point are considered abnormal. The WHO cut-off points are 14gm/dl for adult men; 12gm/dl for adult women. The cut-off points are likely a bit too high and therefore the prevalence of anaemia has been exaggerated. Improved cut-off values of 11.7gm/dl in women and 13.2gm/dl in men are more reasonable. According to data from NHANES II (National Health And Nutrition Examination Surveys) in the United States Anaemia as defined with the improved cut-off points was noted in about 3% of adult men and in 4 to 6% of adult women. NHANES II was performed from 1976 to 1980 which involved the use of sampling techniques that allowed extrapolation of the data to the United States population as a whole, such variables as age, gender, race, geographic area and socio-economic status were considered. In women the disorder occurs most frequently during the reproductive years, where as in men the prevalence is relatively high in adolescence and low during young adulthood, it increases thereafter with advancing age.

## 3. Materials & Methods

The present work has been undertaken on 150 cases between age group of 18 to 24 years, with the object to study the haemoglobin, TRBC, PCV, Absolute Indices (MCV, MCH and MCHC) in medical students of AIMSSKS medical college Bhopal between November 2016 to April 2017. Institutional ethics committee permission was obtained. Each case was first subjected to a careful clinical examination to exclude the possibility of any systematic disorder, diseases or deficiency.

The following investigation were undertaken:

- 1) Hemoglobin by cyanmeth method spectrophotometrically.
- 2) TRBC by haemocytometer(Improved Neubauer's chamber)

- 3) PCV/Hct by centrifugation Wintrobe method @ 3000 rpm for 30minutes.
- 4) Absolute Indices derived by calculation with standard formulas.

The cyanmethemoglobin technique is the method of choice selected by the International Committee for Standardization in Hematology(ICSH).

**Collection of Sample**

Two ml of blood was taken from ante-cubital vein by venepuncture by 24-guage needle under aseptic conditions using a disposable syringe. This blood was poured in EDTA vials and was used for the haematological investigation as already mentioned.

**Equipments**

1. Drabkin’s solution for haemoglobin estimation
2. Photoelectric Colorimeter
3. Improved Neubauer’s Chamber
4. Automated Pipettes(5 & 20micro lt), tips, tissue paper
5. EDTA vials
6. Spirit swabs, disposable syringes
7. Test Tubes
8. Micro glass Slides
9. Field A and B stain
10. Methanol
11. Cover slip
12. Microscope –Light
13. Winrobe’s tube
14. Centrifuge Machine
15. Incubator
16. Distilled Water

**Precaution**

- 1) Hemolysis interferes with the test as the RBC’s have a very high iron content.
- 2) All glassware being used for the test were first rinsed with 1% or 0.1 N HCL and then with good quality deionised water before use.

Statistical analysis of the entire series, males and females was done. There mean values, standard error, t-values and p-values were calculated according to standard formula.


**4. Observations**

The present study was conducted on 150 medical students of which 93 were males and 57 were females. All the students were investigated for their anaemia prevalence. The students were between age of 18 to 24 years. Most of the cases had an average moderate type of constitution and few females were thin built. It was observed that majority of the subjects were sedentary while some where physically active. Non of our students were addict ed. Majority of them were vegetarians and a few were occasional non-vegetarians. All belonged to upper and lower middle socio-economic class.

The values for mean haemoglobin, TRBC count, PCV, and Absolute Indices(MCHC,MCH and MCV) with standard error in the case of 93 male and 57 female medical students have been presented in Table-1.

**TABLE -1**  
**SHOWING THE MEAN VALUES OF VARIOUS PARAMETERS AND THEIR STANDARD ERROR IN THE MALES, FEMALES AND ENTIRE SERIES**

Sex and Number	Hb.	TRBC	PCV/hct	MCHC	MCH	MCV
	Gm/dl	Millions /cu.mm	%	Gm%	Pg	Cu.μ
<b>Both sexes (150)</b>	13.2 ±0.12	4.97 ±0.05	44.04 ±0.36	29.8 ±0.16	26.7 ±0.26	89.62 ±0.75
<b>Male (93)</b>	13.9 ±0.10	5.14 ±0.06	46.21 ±0.35	30.07 ±0.19	27.35 ±0.33	91.06 ±0.95
<b>Female (57)</b>	11.9 ±0.17	4.68 ±0.08	40.49 ±0.45	29.32 ±0.27	25.6 ±0.4	87.28 ±1.16



The bar diagram No.1 shows the mean values obtained in males, females and entire series. When these values were compared it was observed that the values are higher in males than females.

On statistical analysis it was observed that there was a statistical significant difference in the mean values of haemoglobin, PCV and MCH in males and females. Table-2//.....

**TABLE 2**  
**SHOWING STATISTICAL ANALYSIS OF PARAMETERS OF TABLE-1**


Parameters	Males (93)		Females (57)		T-value	P-value
	Mean	S.E.	Mean	S.E.		
Hb gm/dl.	13.9	± 0.12	11.9	±0.17	10	<0.001
TRBC million/cu.mm.	5.14	± 0.06	4.68	±0.08	0.15	<0.8
PCV %	46.21	±0.35	40.49	±0.45	10	<0.001
MCHCgm %	30.07	± 0.19	29.32	±0.27	2.27	<0.02
MCH pg.	27.35	± 0.33	25.6	±0.4	3.37	<0.001
MCV cu.μ	91.06	±0.95	87.28	±1.16	2.52	<0.01

Although the mean values of TRBC, MCHC and MCV are higher in males than females but their difference were not statistically significant.

Table-3 shows mean values with their standard error of haemoglobin, TRBC count, PCV and Absolute Indices in case of 24 male and 32 female medical students who were having haemoglobin cut off less than 13.2gm/dl for males and 11.7gm/dl for females.

**TABLE -3**  
**SHOWING PARAMETERS OF 56 STUDENTS USING REVISED CUT-OFF POINTS (13.2 GM/DL FOR MALES; 11.7 GM/DL FOR FEMALES)**

Sex and Number	Hb	TRBC	PCV/hct	MCHC	MCH	MCV
	gm/dl	Millions/cu.mm	%	gm%	pg	Cu.μ
Both sexes (56)	11.7 ±0.14	4.71 ±0.09	40.94 ±0.53	28.64 ±0.24	25.26 ±0.44	88.3 ±1.4
Male (24)	12.6 ±0.08	4.88 ±0.13	43.8 ±0.7	28.91 ±0.4	26.3 ±0.67	91 ±2.06
Female (32)	10.9 ±0.13	4.59 ±0.123	38.8 ±0.52	28.43 ±0.32	24.5 ±0.57	86.2 ±1.9



The bar diagram No-2 exhibits the mean values of males and females who were below the revised cut- off points for

haemoglobin. The means are again higher for males when compared with females.

**Table 4**  
**SHOWING STATISTICAL ANALYSIS OF PARAMETERS OF TABLE-3**

Parameters	Males (24)		Females (32)		t-value	p-value
	Mean	S.E.	Mean	S.E.		
Hb (gm/dl.)	12.6	±0.08	10.9	±0.13	11.2	<0.001
TRBC (million/cu.mm.)	4.9	±0.13	4.6	±0.123	1.5	<0.1
PCV (%)	43.8	±0.7	38.8	±0.52	5.6	<0.001
MCHC (gm %)	28.91	±0.4	28.43	±0.32	1	<0.3
MCH (pg.)	26.3	±0.67	24.5	±0.57	2.34	<0.01
MCV (cu.μ)	91	±2.06	86.2	±1.9	1.71	<0.05

The statistical difference in the values of these parameters in males and females is depicted in Table-4. This Table depicts that haemoglobin content and PCV values in males are higher as compared to females and the differences are highly statistically significant. There were no significant difference in other parameters of males and females.

## 5. Discussion

The present work was undertaken in 150 medical students with the object of studying the prevalence of anaemia amongst them. There were 93 male and 57 female of age group 18 to 25 years. All the subjects were examined clinically. The haemoglobin estimation, TRBC count, PCV, Absolute Indices were done in all cases. The mean value with standard error of the haemoglobin content was 13.9+0.10 in males and 11.9+0.17 in females, TRBC count was 5.14+0.06 and 4.68+0.08, PCV was 46.21+0.35 and

40.5+0.45, MCHC was 30.07+0.2 and 29.32+0.27, MCH was 27.35+0.33 and 25.6+0.40, MCV was 91.06+0.95 and 87.28+1.16. All the parameters were less in females as compared to males. In all the parameters except TRBC count and MCHC, the differences were highly significant. On interpretation of the data, using the WHO criteria (cut-off points 14gm/dl for males and 12gm/dl for females) for reference placed the prevalence of anaemia at 49.3%. According to the WHO technical report 1968 to 1975 the prevalence of anaemia in India (Vellore) for males was 6% and for females was 35%. While it was 64% in females in India (Delhi), following the same criteria. However according to the data of NHANES II from 1976 to 1980, the prevalence of anaemia in the age group between 18-24 years in the United States for males was 2.7% and for females was 3.3% with the revised cut-off points (13.2gm/dl for males and 11.7gm/dl for females). When this study used the revised cut-off points, 56 students were found below the cut-

off points, out of which 16% were males and 21.3% were females.

## 6. Summary

The present study was under taken in 150 medical students, which included 93 males and 57 females. The entire series was subjected to haematological investigation for anaemia prevalence. In this study the case records of each subject were maintained. Out of 150 students, 56 were below revised cut-off points for haemoglobin i.e 13.2gm/dl for males and 11.7gm/dl for females. But on carrying out transferrin saturation studies only 14 female subjects had their transferrin saturation between 15 to 20%- the reference range for latent iron deficiency anaemia. Rest of the students (42) were found to have transferrin saturation above 20%.

## 7. Conclusion

The use of a single haemoglobin estimation is of limited value in determining the prevalence of anaemia. If the WHO protocol is used for reference the prevalence of anaemia is amplified due to higher cut-off values which are not applicable in our country. The revised cut-off values provided by NHANES(United States) for prediction of anaemia are better applicable in a developing country like India. The overall prevalence of anaemia in our study was found to be 9.3%. But since the anaemic subjects were exclusively females it can also be concluded that the prevalence was zero percent in males and 24.56% in females.

## References

- [1] Gillespie S. Major Issues in the control of Iron deficiency. The micronutrient initiative. UNICEF, 1998.
- [2] Mehta, M.N. : Effectiveness of daily iron and weekly iron and folic acid supplementation in anaemic adolescent girls. UNICEF Funded final report of the research project, Bombay urban ICDS project, 1998.
- [3] Nelson, M. Bakalion, F. Trivedi, A. : Iron Deficiency anaemia and physical performance in adolescent girls from different ethnic background. British Journal of Nutrition. 4:433-437, 1994.
- [4] NHANES-II : National Centre for Health Statistics: Hematological and Nutritional Biochemistry reference data for persons six months to seventy four years of age United States, 1976-1980. Vital and Health Statistics Series II – No 232. 1982.
- [5] Raina, N. Gupta, M. : Operational study on nutritional anaemia in pregnant lactating women and adolescent girls in a rural community in India. Improving the quality of iron supplementation programme. The mother care experience. Mother care. John Snow incorporated. 1996
- [6] Rajaratham, J. Rajaratham, A. : Prevalence of anaemia among rural girls of Tamil Naidu. Indian Paediatrics. 37:532-536,2000.
- [7] V. Anuradha, and Sangeetha, K. : Development and impact ofsoyamalt product in improving the iron status of anaemic adolescent girls (16-18 years). The Indian Journal of Nutrition and Dietetics. 38:141-145,2001.
- [8] World health Organization. Iron deficiency anaemia. Assessment, prevention and control: A guide for programme managers. Geneva, Switzerland: World Health Organization;2001.
- [9] Shah BK, Gupta P. Weekly v/s Daily Iron and Folic acid supplementation in adolescent Nepalese girls. Arch Paediatric Adolescence Med.2002;156:131-5.
- [10] Stoltzfus RJ, L Mullany, RE Black. Iron deficiency anaemia. In; Ezzati M, Lopez AD, Odgers A, Murray JL, editors. Comparative quantification of health risks: global and regional burden of diseases attributable to selected major risk factors. Geneva, World Health Organization;2004:163-209.
- [11] Abalkhail B, Shawky S. Prevalence of daily breakfast intake, iron deficiency anaemia and awareness of being anaemic among Saudi School Students. Int Journal Food Science Nutrition.2002;53(6):519-28.
- [12] Ayoub AI. Iron deficiency anaemia in Dubai Medical College for Girls: A preliminary Study. J Egypt Public Health Assoc. 1995;70(1-2):213-28
- [13] Iron deficiency anaemia: Assessment, Prevention & Control. A guide for programme managers. Geneva, World Health Organization, 2001 (WHO/NHD/01.3).
- [14] Shill KB, Karmakar P, Kibria MG, Das A, Rehman MA, Hossain MS et al. Prevalence of Iron deficiency anaemia among University students in Noakhali region, Bangladesh. Journal Health Population Nutrition 2014;32(1):103-10.
- [15] Bhimrao NJ. A critical review on iron deficiency anaemia in female population of developing India. International Journal Fauna Biologic Studies. 2014;3(5):116-9.
- [16] Ghorpade VV, Shinde PP, Madhekar NS, Pol VS. Prevalence of anaemia in female students of pharmacy college and its association with various socio-demographic variables: A study conducted in rural teaching institute, Kasegaon. International Journal Contemp Med Res.2016;3(8):2295-7.
- [17] Kurubaran G, Sami AR, Ahmed MQ, Al-abed AA. Social and Psychological factors affecting eating habits among university students in a Malaysian Medica School: A cross sectional study. Nutrition J. 2012,11:48.
- [18] Consensus group. Consensus statement for diagnosis of obesity, abdominal obesity and metabolic syndrome for Asian Indians and recommendations for physical activity, medical and surgical management. JAPI. 2009;57:163-70.
- [19] Roy RN, Saha I, Mahajan, Gupta. Textbook of preventive and social medicine. Fourth edition, Jaypee Brothers Medical Publishers (P) Ltd. New Delhi. 2013:134.
- [20] WHO Report, World Prevalence of Anaemia 1993-2005. WHO Global Database on Anaemia, World Health Organization, Geneva, Switzerland, 2008.
- [21] Khan HM, Sohail M, Ali A, Akhtar N, Khan H, Rasool F. Symptoms based on evaluation of iron deficiency anaemia in students of Bahawalpur correlated with their eating habits. Tropical Journal Pharm Res.2014;13(5);769-72.
- [22] Al Hassan NN. The prevalence of iron deficiency anaemia in a Saudi University female students. Journal Microscopy Ultrastructure. 2015;(1):25-8.

- [23] Subramaniyam K, George M, Seshadri D, Jena A, Chandrababha N. Prevalence of anaemia among health science students of a university in South India. *Int J Res Med Sci.* 2016;4(10):4598-601.
- [24] Manjula VD, Parameshwari P, Pothen L, Sobha A. Prevalence of anaemia among female undergraduate students of Government Medical College Kottayam, Kerala. *Int J Med Health Sci.* 2014;3(2):133-8.
- [25] Sultan A H. Anaemia among female college students attending the University of Sharjah, UAE: Prevalence and Classification. *J Egypt Public Health Assoc.* 2007;82(3-4):261-71.
- [26] Al-Sayes F, Gari M, Qusti S, Bagatian N, Abuzenadah A. Prevalence of iron deficiency anaemia among females at university stage. *J Med Lab Diag.* 2011;2(1):5-11.
- [27] Salameh P, Jomaa L, Issa C, Farhat G, Salame J, Zeidan N et al. Lebanese National Conference for Health in University Research Group. Assessment of dietary intake patterns and their correlates among University students in Lebanon. *Frontiers Public Health.* 2014;2:185.
- [28] Waseem SM, Bano R, Ahmad N, Kumar J, Khan A. Study of Haematological Profile and Body Mass Index in Undergraduate Medical Students in Lucknow, Uttar Pradesh. *IJHSR.* 2015;5(6):257-62.

