

# Assessment of Awareness on Cashew Insect Pests, Diseases and Management Practices in Tanga Region, Tanzania

W. A. Nene<sup>1</sup>, A. R. Makale<sup>2</sup>, M. William<sup>3</sup>

<sup>1,2,3</sup>Naliendele Agricultural Research Institute, Box 509, Mtwara, Tanzania

**Abstract:** Cashew production in Tanzania suffers from numerous biotic constraints particularly insect pests and fungal pathogens. Different pest management options are available to cashew growers; however, the most devastating diseases and insect pests that attack cashew and management options against these diseases and pests particularly in Tanga region are not well known. We conducted a study to find out the level of knowledge on cashew nuts pests, diseases and the management practices. The results showed that the cashew nut bugs and stem borers were the major (53 %) insect pests attacking cashew crop, whereas Powdery Mildew and Cashew Leaf and Nut Blight were the most (86 %) devastating diseases. Majority of the respondents (45%) controlled these pests and diseases mainly using pesticides. More than 60 % of the respondents did not use protective gears. Most (49 %) of them lacked the knowledge on the importance of wearing protective gears during the application of the pesticides. We recommend that farmers' training is needed to improve awareness among them on insect pests and diseases management, particularly the use of Integrated Pest Management program. Also protective gears should be subsidized and be available to users.

**Keywords:** Cashew, diseases, insect pests, pesticides, Tanzania

## 1. Introduction

The majority (about 80 percent) of the Tanzanian population lives in the rural areas and largely depends on agriculture for their livelihood [1]. The rural Tanzanians have been engaged in the cultivation of both cash and food crops. In the case of cash crops, the Cashew, *Anacardium occidentale* L. (Sapindales: Anacardiaceae) is an important foreign-exchange earner for Tanzanian and ranked the first in the year 2016 [2]. Globally, the crop has been reported for its multiple advantages which include being a source of food and income [3, 4, 5], its use in the reforestation and prevention of erosion [3, 6], its medicinal value [7, 8] and the utilization of the cashew apple in the production of various products such as syrups, juice, wine alcohol, jam, vinegar and candy [9]. Thus, any interventions that aim at improving the agricultural sector and the cashew crop production in particularly can cause a significant impact in improving the livelihood of the majority of people in the country. In this respect, the Government of Tanzania has continued to support cashew farmers in various ways. Among these include; the reduction of unnecessary taxes which negatively affect farmers' motivation, free provision of agricultural inputs such as pesticides (fungicides and insecticides), and planting materials such as seeds and seedlings. The Government through Cashew Board of Tanzania (CBT) established a Warehouse Receipt System (WRS) which seems to benefit majority of cashew farmers [10]. Despite these measures, cashew productivity in Tanzania has not been optimal mainly due to insect pests and diseases [11, 12, 13, 14]. Thus, in order to sustain farmers' morale in engaging in cashew production resulting from Government incentives to cashew farmers, there is a need to reduce and ultimately eliminate huge cashew nut losses which are caused by major insect pests and diseases.

Several insect pests and diseases that attack cashew crop have been reported elsewhere causing significant losses in

yields [11, 15, 16, 13, 17]. Fungal diseases such as Powdery mildew which is caused by *Oidiumanacardii* and Leaf and Nut blight which is caused by *Cryptosporiopsis* sp. cause yield losses of between 70 and 100 % if no control measures are taken [13]. Similarly, insect pests such as cashew bugs (*Helopeltis spp.*) and coconut bug (*Pseudotheraptuswayi*), have been the greatest threat in cashew crop production [18, 19, 20].

Different management strategies have been developed to control insect pests and diseases of cashew. These include the use of resistant cashew variety [13], biological control [21, 22, 23, 24], synthetic chemicals [13], botanical pesticides [11], and ecological management strategies [13]. There is, however scant information on the major insect pests, diseases that affect cash crop and management practices of cashew farming community in Tanga region. This paper presents insect pests and diseases that attack cashew crop in Tanga region. It also looks at the management strategies which are practiced by farmers and reasons for adopting such strategies. The information to be obtained would help in developing appropriate control options which may result to improved cashew productivity in Tanga region.

## 2. Materials and Methods

### Description of the study area

The current study was conducted in Tanga region which is located in the Eastern zone of Tanzania. The region lies between latitudes 4° and 6° south of the Equator, and between longitudes 37° and 39° east of Greenwich. Eighty eight (98) Cashew stakeholders (Farmers = 57, Extension officer = 19 and blower operators = 22) from Muheza, Korogwe, Pangani, Mkinga and Tanga districts in Tanga region were involved in the study. The districts were selected based on their potential in cashew production. The respondents were purposively selected by the District

Agriculture, Irrigation and Cooperative Officer, on the basis of engagement in cashew farming (farmer), working as agricultural field officer or working as a blower operator who was applying pesticides against cashew insect pests and diseases using motorized blower machine. A semi structured questionnaire was administered to the respondents between June and July 2017. The respondents were allowed to give their opinions freely and without being interrupted. Samples of the common insect pests and disease symptoms attacking cashew crops were displayed in order to remind the respondents and help them to explain the real pests and diseases attacking cashew crop in their area. In this case, the respondents were brought together but interviewed individually. The information gathered included major insect pests and diseases attacking cashew in their area, and management tactics which were used against insect pests and diseases.

### Data analysis

The data which were collected were analysed using Statistical Package for Social Sciences (SPSS 16.0 for windows) computer software. Descriptive statistics which were obtained were summarized and presented in the form of tables and graphs. Secondary data were collected from various sources including NARI annual reports, proceedings, review of published papers and official reports, and online information.

### 3. Results

The study included cashew farmers, blower operators and agricultural field officer from Muheza, Mkinga, Pangani, Korogwe and Tanga districts. As presented in Table 1, 81 % of the respondents were males while 19 % were females. The majority of the respondents were aged between 31 and 45 years with the minimum age of 25 years.

**Table 1:** Socio economic characteristics of respondents

Characteristics	Frequency	Percent
<b>Gender</b>		
Male	79	80.6
Female	19	19.4
<b>Age</b>		
25-30 years old	18	18.4
31-45 years old	39	39.8
> 46 years old	38	38.8
Don't know their age	3	3.1
<b>Occupation</b>		
Farmers	57	58.2
Blower Operators	22	22.4
Agriculture Field Officer (AFOs)	19	19.4
<b>Districts</b>		
Muheza	9	9.2
Mkinga	26	26.5
Pangani	18	18.4
Korogwe	32	32.7
Tanga	13	13.3

Table 2 (a) and (b) present the respondents' knowledge on diseases and insect pests attacking cashew farms in the selected districts of Tanga region. Majority of the respondents had knowledge on cashew insect pests and diseases. However, the respondents from Korogwe region

had the least knowledge (ranges from 16 and 40%) on cashew diseases and insect pests attacking cashew crop.

**Table 2a:** Knowledge on diseases attacking cashew in Tanga region

Cashew diseases	District (%)				
	Muheza	Mkinga	Pangani	Korogwe	Tanga
Powdery mildew and Leaf and nut blight	8.3	22.9	22.9	35.4	10.4
Only Powdery mildew	12.5	41.7	20.8	16.7	8.3
Powdery mildew, Leaf and nut blight, die back and Anthracnose	8.3	25	0	33.3	33.3
Don't know	0	20	20	40	20
Powdery mildew and Anthracnose	25	25	25	25	0
Leaf and nut blight	0	0	0	75	25
Anthracnose	0	0	0	100	0

**Table 2b:** Knowledge of insect pests attacking cashew in Tanga region

Cashew insect pests	District (%)				
	Muheza	Mkinga	Pangani	Korogwe	Tanga
Cashew bugs and Stem borers	9	24.4	17.9	32.1	16.7
Cashew and coconut bugs, Stem borers	0	50	25	25	0
Cashew and coconut bugs	50	50	0	0	0
Only Stem borers	0	37.5	25	37.5	0
Don't know	16.6	16.7	16.7	50	0

The majority (53 %) of the respondents indicated that, the major insect pests attacking cashew were cashew nut bugs (*Helopeltisanacardii* and *H. schoutedeni*) and stem borer (*Mecocorynusloripes*); while 2% of the respondents mentioned coconut bug (*Pseudothertapswayi*) as the least insect pest problem attacking cashew. On the other hand, Powdery Mildew Disease (PMD) and Cashew Leaf and Nut Blight (CLNB) were mentioned by 86 % of the respondents as the major diseases attacking cashew crop (Table 3).


**Table 3:** Major insect pests and diseases attacking cashew in Tanga region

Insect pests	Frequency	Percent
Cashew nut bugs and stem borers	52	53.1
Only Stem borer	22	22.4
Only cashew nut bugs	12	12.2
Only coconut bugs	2	2.0
Cashew nut and Coconut bugs	5	5.1
<b>Diseases</b>		
PMD and CLNB	85	86.7
Anthracnose and CLNB	4	4.1


The study revealed that 45% of the respondents controlled these pests and diseases mainly through the application of pesticides. Other methods of pest control used included sanitation only (reported by 24 %), sanitation and pesticide application (11 %), sanitation and biological control (8%), and the use of pesticides, improved cashew planting materials and sanitation (5%). About 7% of the respondents

did not use any insect pests and diseases control methods (Figure 1). The results in Figure 2 indicate that both Ninja and Duduall were the most pesticides used to control insect pests. Other insecticides were Karate, Selecron, Subachlo and Fastac. Majority of the respondents control PMD using

either Sulphur dust or a mixture of spraying regimes of Sulphur dusts and other water based fungicides such as Bayfidan, Anvil, Transmute, Acanto, Subatex, and Nativo.



**Figure 1:** Knowledge on different methods used to control cashew insect pests and diseases



**Figure 2:** Common/trade names and level of pesticides use in cashew diseases and insect pests control in Tanga region

The results in Table 4 indicate that, more than 60 % of the respondents do not use protective gears. Most of them (49 %) lack knowledge on the importance of wearing protective gears during the application of pesticides (Table 5a). More than 50 % of farmers being interviewed (Table 5b) show that, high cost of protective gear, lack of knowledge or a combination of both are the major constraints of using protective gears during the application of pesticides.

**Table 4:** Knowledge on the use of protective gears

Level of protective gears use among the operators	Gender		Frequency	Percent
	Male	Female		
Protective gears are used	75.8	24.2	33	33.6
No use of protective gears	83.9	16.1	62	63.3
Don't know	66.7	33.3	3	3.1

**Table 5a:** Reasons for not using protective gears

Reasons for not using protective gears	Frequency	Percent
High cost of protective gears	17	17.3
Lack of knowledge	48	49
Lack of knowledge, poor cooperation and high cost	33	33.7

**Table 5b:** Reasons for not using protective gears among Cashew stakeholder

Stake holders	High cost of protective gears	Lack of knowledge	Lack of knowledge, poor cooperation and high cost
Farmers	50.0	87.5	54.5
Operators	16.7	12.5	24.2
AFOs	33.3	0.0	21.3

#### 4. Discussion

The study revealed the presence of insect pests and diseases attacking cashew crop. The major insect pests which were revealed include cashew nut bugs (*H. anacardii* and *H. schoutedeni*) and stem borer (*M. loripes*) while PMD which is caused by *O. anacardii*Noack and CLNB diseases which is caused by a fungal pathogen known as *Cryptosporiopsis* spp were the major diseases in cashew crop. Cashew nut bugs and stem borers are serious pests reported to cause significant losses of cashew nuts [13]. Cashew bugs feed on tender leaves and shoots. These pests may cause death of the leaves or shoots under intense feeding [25, 26]. Adult weevils of the stem borers (*M. loripes*) lay their eggs beneath the bark of cashew tree branches. When the larvae hatch they burrow into the cambium making large tunnels, eating the sapwood of the tree and this may cause death of the tree. *Mecocorynus* sp. was also reported as a new serious pest in Ghana [27]. Similarly, PMD [28] and CLNB [29] were cited as the leading devastating diseases of cashew crop in the country. These diseases can cause cashew nut losses of between 70 and 100 % if no any control measures have been taken [13].

This study found that majority of cashew farmers control insect pests and diseases using pesticides. It was mentioned that, Ninja and Duduall are the insecticides which are mostly used to control insect pests. Ninja 5EC is a synthetic pyrethroid containing Lambda cyhalothrin as an active ingredient, whereas Duduall contains Cypermethrin 150g/L+Chlorpyrifos 300g/L. These insecticides have been registered by Tropical Pesticides Research Institute (TPRI) to control cashew sucking pests such as cashew nut and coconut bugs [30]. Other insecticides which are used by cashew farmers in Tanga region and registered by TPRI are Karate (Lambda cyhalothrin 50g/L, Subachlo (Chlorpyrifos 500g/L+Cypermethrin 55g/L), Fastac (Alphacypermethrin 100g/L) and Selecron (Profenofos 720g/L). Fungicides which are used by most of farmers in the study area to control diseases include Sulphur or Sulphur with other fungicides. Other water based fungicides used contain Triadimenol (Bayfidan and Transimute), Hexaconazole (Anvil), Tebuconazole and Triadimenol (Subatex), Trifloxystrobin and Tebuconazole (Nativo) and Picoxystrobin (Acanto). The Naliendele Agricultural Research institute (NARI) has recommended several options

of controlling insect pests and diseases, and these include the use of pesticides, cultural practices including sanitation, the use of resistant varieties, and biological control such as the use of weaver ant (*Oecophyllalolonginoda*Latreille) [13].

Our findings established that majority of the respondents had knowledge on insect pests, diseases and management practices. However, their level of knowledge varied across districts. Respondents from Korogwe District had the least level of knowledge (16 to 40 %) compared to other districts. This implies that training is needed particularly in Korogwe district.

The results of the present study also revealed that, majority of the respondents use a single technique; in this case, it is the application of chemicals to control insect pests and diseases. If only chemical are used for a long time, this can result to negative impacts such as pest resurgence, secondary pest outbreak, and pest resistance to chemicals [31]. Thus, Integrated Pest Management (IPM) strategies which involve a combination of more than one technique in pests and diseases management are recommended. Few of the interviewees were observed using biological methods such as weaver ant, *Oecophyllalolonginoda*Latreille to control insect pests. Weaver ants have been utilized as potential bio agents for protecting crops from attack by insect pests [23, 24,32, 33]. Thus, optimizing the use of weaver ant in an IPM program could not only reduce yield losses caused by insect pests but also it could also reduce the costs incurred by farmers to buy chemicals [2015].

We also found that, most of the respondents do not use protective gears during the use of pesticides. This can cause negative impact to their health. The Government is subsidizing motorized blowers; however, the protective gears are not included by the suppliers in the package for the subsidy. Thus, it is suggested that subsidized motorized blowers should be supplied with protective gears. Furthermore, farmers should be trained to create awareness among them on the importance of wearing protective gears while using the chemicals.

#### 5. Conclusion

This study looked at insect pests and diseases that attack cashew crop in Tanga region. It also looked at the management strategies which are practiced by farmers and reasons for adopting such strategies. The study found that cashew nut bugs and stem borers were the major insect pests attacking cashew crop; and PMD and CLNB diseases were the major diseases causing significant loses of cashew. The application of chemicals was the main technique used to control cashew pests and disease, and majority of respondents do not use protective gears during chemical usage. Awareness creation is needed to improve farmers' knowledge on insect pests, diseases and their management strategies particularly the use of IPM program. Protective gears should be available and subsidized to enable users access them. Furthermore, farmers' training on the importance of protective gears usage is essential so as to reduce health risks which might occur from neglecting their usage particularly during the application of these synthetic chemicals.

## 6. Acknowledgement

This study was financially supported by the Government of Tanzania through the Cashew Board of Tanzania (CBT). Extension officers, blower operators and farmers are acknowledged for their willingness to provide information regarding this study.

## References

- [1] ESRF (2013). Agricultural Trade Policies Tanzania. [<http://www.fao.org/docrep/013/al668e/al668e03.pdf>]. Site visited on 24/10/2017.
- [2] BOT (2017). Bank of Tanzania monthly economic review march, 2017 <http://www.bot.go.tz/Publications/MonthlyEconomicReviews/MER170300%20MER%20MAR%202017.pdf>
- [3] Adeigbe O.O., Olasupo F.O., Adewale B.D. and Muyiwa A.A., (2015). A review on cashew research and production in Nigeria in the last four decades Scientific Research and Essays, 10(5): 196-209.
- [4] International Nuts and Dried Fruit Council, (2015). [www.nutfruit.org](http://www.nutfruit.org). Site visited on 15/10/2017
- [5] Jøker D., (2003). Information about Cashew Nut (*Anacardium occidentale*). [[http://www.hubrural.org/IMG/pdf/anacarde\\_danida.pdf](http://www.hubrural.org/IMG/pdf/anacarde_danida.pdf)]. Site visited on 12/11/2017.
- [6] Davis K., (1999). Cashew. Echo Technical Note. <http://www.echotech.org/network/modules.php?name=News&file=article&sid=597>. Site visited on 10/11/2017.
- [7] Morais S.M., Silva K.A., Araujo H., Vieira I.G.P., Alves D.R., Fontenelle R.O.S. and Silva A.M.S. (2017). Anacardic Acid Constituents from Cashew Nut Shell Liquid: NMR Characterization and the Effect of Unsaturation on Its Biological Activities. Pharmaceuticals
- [8] Aderiyi B.I., David O.M. and Atere V. A., (2014). Administration of cashew extracts in the treatment of some infections and diseases. Advancement in Medicinal Plant Research 3(3): 75-86.
- [9] Nwosu C., Adejumo O.A. and Udoha W.N., (2016). Cashew apple utilization in Nigeria: Challenges and prospects. Journal of Stored Products and Postharvest Research, 7(2): 29-31. DOI: 10.5897/JSPPR2015.0190.
- [10] Mkonya N., and Cameron A., (2015). FAO Report: Analysis of price incentives for cashew nuts in the United Republic of Tanzania, 2005-2013, 54pg.
- [11] Nene W.A., Shomari, S.H., and Assenga, B.B., (2017). The efficacy of botanical pesticides for managing powdery mildew, *Oidium anacardii* Noack disease in cashew, *Anacardium occidentale* L. plantations in Tanzania. Research Journal of Agriculture and Forestry Sciences, 5(10): 1-6.
- [12] Malegesi, M. (2015). Status of Cashew nut Industry in Tanzania, Cashew Board of Tanzania. Proceedings of the Third International Cashew Conference, Tanzania, 16-19<sup>th</sup>, November 2015.
- [13] Sijaona M.E.R., (2013). Important diseases and Insect Pests of Cashew in Tanzania. Naliendele Agricultural Research Institute, Tanzania, 44pp.
- [14] Kasuga L.J. (2010). Status of the Cashew nut Industry in Tanzania, Second International Conference, 26-29 April, Kampala Uganda.
- [15] Wonni I., Sereme D., Ouedraogo I., Kassankagno A.I., Dao I., Ouedraogo L., and Nacro S., (2017). Diseases of Cashew Nut Plants (*Anacardium occidentale* L.) in Burkina Faso. Advances in Plants & Agriculture Research, 6(3): 00216
- [16] Agboton, B.V., Salifu, D., Seguni Z., Sijaona M.E., Shomari S., Ekesi, S., and Maniania N.K., (2013). Bioecology of some key cashew insect pests and diseases in diverse habitats and landscapes in Tanzania. Journal of Applied Entomology, 137 (10): 782-789.
- [17] Dwomoh E.A., Ackonor J.B., and Afun J.V.K., (2008). Survey of insect species associated with cashew (*Anacardium occidentale* Linn.) and their distribution in Ghana. African Journal of Agricultural Research 3 (3): 205-214.
- [18] Chipojola, F.M. and Kondowe, E.M. (2015). The Status of the Cashew Industry in Malawi. Proceedings of the Third International Cashew Conference, Tanzania, 16-19<sup>th</sup>, November 2015.
- [19] Shomari, S.H., Menge, D. and W. Nene (2015). Evaluation of Five Selected Potential Botanicals against Cashew Powdery Mildew Disease. Cashew Research Programme, Naliendele Agriculture Research Institute, Mtwara-Tanzania and Masai Mara University, Narock-Kenya. Proceedings of the third international cashew conference, Tanzania, 16-19<sup>th</sup>, November 2015.
- [20] Nene, W., Mwakanyamale, D. E., Shomari, S.H. and Kidunda, B., (2015). Assessing Farmers' Awareness on the Utilization of the Weaver Ants, *Oecophylla longinoda* Latreille for the Control of Cashew Insect Pests in the Eastern Zone of Tanzania. Proceedings of the Third International Cashew Conference, Tanzania, 16-19<sup>th</sup>, November 2015.
- [21] Nene, W., Rwegasira, G.M, Offenber, J. Mwatawala, M. Nielsen, M.G., (2015). Mating Behavior of the African Weaver Ant, *Oecophylla longinoda* (Latreille) (Hymenoptera: Formicidae). *Sociobiology*, doi: 10.13102/sociobiology.v62i3.
- [22] Dwomoh, E.A., Afun, J., V., K., Ackonor, J., B., & Agene, V., N., (2009). Investigations on *Oecophylla longinoda* (Latreille) (Hymenoptera: Formicidae) as bio control agents in the protection of cashew plantations. Pest Management Sciences, 65: 41-46.
- [23] Olotu M.I., du Plessis H., Seguni Z.S., Maniania N.K., (2013a). Efficacy of the African weaver ant *Oecophylla longinoda* (Hymenoptera: Formicidae) in the control of *Helopeltis* spp. (Hemiptera: Miridae) and *Pseudotheraptus wayi* (Hemiptera: Coreidae) in cashew crop in Tanzania. Pest Management Sciences 69: 911-918. doi: 10.1002/ps.3451
- [24] Abdulla N.R., Rwegasira G, Jensen KV, Mwatawala MW, Offenber J (2015) Effect of supplementary feeding of *Oecophylla longinoda* on their abundance and predatory activities against cashew insect pests. Biocontrol Science and Technology 25(11): 1333-1345. doi: 10.1080/09583157.2015.1057476
- [25] Boma, F., Topper, C.P. & Stathers, T. (1998). Population dynamics of *Helopeltis* spp on cashew in Southern Tanzania. In Topper *et al* eds. (1998e),

- Proceedings of the International Cashew and Coconut conference, Dar es Salaam, Tanzania, 17<sup>th</sup> to 20<sup>th</sup> February, 1997.
- [26] Topper, C.P., Grunshaw, J., Pearce, M., Boma, F. Stathers, T. and Anthony, J. (1998). Preliminary observations on *Helopeltis* and *Pseudotheraptus* damage to cashew leaves and panicles. In: Topper *et al* eds.(1998e), Proceedings of the International Cashew and Coconut conference, Dar es Salaam, Tanzania, 17<sup>th</sup> to 20<sup>th</sup> February, 1997.
- [27] Dwomoh E.A., Ahadzie, S.K. , Somuah, G.A. and Amenga A.D. (2011). Preliminary studies on the damage symptoms and the spatial distribution of an emerging insect pest, *Mecocorynus* sp. (Coleoptera: Anthribidae) on cashew in Ghana. Journal of Cell and Animal Biology, 5(7): 144-151.
- [28] Sijaona, M.E.R. and Mansfield J.W., (2001). Variation in the response of cashew genotypes to the targeted application of fungicide flower panicles for control of powdery mildew disease. Plant Pathology, 50: 244-248
- [29] Sijaona M.E.R., Reeder R.H., and Waller J.M. (2006). Cashew leaf and nut blight. A new disease of cashew in Tanzania caused by *Cryptosporiopsis* spp. Plant Pathology 55:576
- [30] TPRI (2011). Wizaraya Kilimo Chakulana Ushirika: Orodhayaviatifu vilivyosajiliwa Tanzania. [http://www.tpri.or.tz/news/Pesticides\_Gazette\_2011] site visited on 03/11/2017
- [31] Hajek, A. (2004). Natural Enemies. An Introduction to Biological Control. Cambridge University press, Department of Entomology, Cornell University.
- [32] Peng R.K., Christian K. (2007). The effect of the weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae), on the mango seed weevil, *Sternochetus mangiferae* (Coleoptera: Curculionidae) in mango orchards in the Northern Territory of Australia. International Journal of Pest Management 53(1): 13–24.
- [33] Peng R.K., Christian K., Lan L.P., Binh N.T., (2008) Integrated cashew improvement program using weaver ants as a major component. Manual for ICI program trainers and extension officers in Vietnam. Charles Darwin University and Institute of Agricultural Science for South Vietnam, 93 pp.
- [34] William, J.G., Hella, J., Lars, E., Offenber, J., Mwatawala, M., & Rwegasira, G. (2015). Benefit–cost Analysis of alternative insect pests’ management in cashew and mango orchards in Tanzania. Quarterly Journal of Econometrics Research, 1: 32–44.