

Phytochemical Determination and *In Vitro* Antimicrobial Activity of Crude Ethanolic Extract of Stem Bark of *Boswellia dalzielii*

Baha'uddeen Salisu Dandashire¹, Abdulkadir Magaji Magashi², Bashir Abdulkadir³

^{1,3}Department of Microbiology, Faculty of Natural and Applied Sciences, Umaru Musa Yar'adua University, Katsina, Katsina State, Nigeria

²Department of Microbiology, Faculty of Science, Bayero University, Kano, Nigeria

Abstract: **Background:** *Boswellia dalzielii* is a medicinal plant widely used in the traditional treatments of Rheumatism, Septic sores, venereal diseases and gastrointestinal ailments. However, the scientific evaluations of the ethanopharmacological claims of the plant have not been adequately resolved. **Aim:** To carryout automated phytochemical screening and determination of the in vitro antimicrobial activity of the crude ethanolic stem bark extract of *B. dalzielii* on some common pathogenic microorganisms which included *Staphylococcus aureus*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Salmonella typhi* and *Candida albicans*. **Methods:** Fifty grams (50g) of the dried powder of the stem bark of the plant was extracted exhaustively in 500ml of 95% ethanol by percolation method for two weeks. Phytochemical screening of the extract was performed using High Performance Liquid Chromatography (HPLC), Fourier Transformed Infrared Spectroscopy (FTIR) and Gas Chromatography-Mass Spectrometry (GCMS) after which its antimicrobial activity was evaluated by agar well diffusion method. The minimum inhibitory concentrations (MICs) and minimum bactericidal concentration (MBC)/ Minimum Fungicidal concentration (MFC) were also determined using the broth micro-tube dilution technique. **Results:** The HPLC analysis revealed the presence of 8 components with major ones at peaks 3, 4, 5 and 2 with peak areas of 31.52%, 21.96%, 17.23% and 16.21% respectively; the FT-IR spectroscopy revealed 11 functional groups which included 1°, 2° amines and amides, alkanes, alkenes, alkynes, alkyl halides, aromatics and aliphatic amines; while the GCMS analysis revealed 13 compounds and the major ones were *n*-Hexadecanoic acid (23.54%), Oxacyclotetradecan-2-one (20.33%), Pelargic acid (16.83%), *n*-Eicosanol (13.51%) and Ethyl striate (10.05%). Several studies reported that these compounds have antimicrobial, anticancer and anti-oxidant activities except *n*-Eicosanol and Ethyl steriate which are highly toxic. The susceptibility test showed that the extract was active against all the test isolates with higher zones of inhibition of 21±0.00mm for *C. albicans* and *S. typhi*, 20±0.80mm for *S. aureus*, and 19±0.50mm for *K. Pneumoniae*, 18±0.80mm for *E. coli* and 18±0.80mm for *P. aeruginosa* at 50mg/ml concentration each. Resistance was only observed at 2.5mg/ml concentration for *S. pyogenes*. Similarly, lowest MIC values of 3.12mg/ml were obtained for *C. albicans* and *S. typhi*, 6.25mg/ml for *S. aureus*, 12.5mg/ml for *K. Pneumoniae*, *E. faecalis* and *P.mirabilis*, 25mg/ml for *E.coli* and *S. pyogenes* and the highest MIC of 50mg/ml for *P. aeruginosa*. The MBC/MFC values did not exceed the corresponding MIC values by more than a factor of 2. **Conclusion:** Although the stem bark of *B. dalzielii* showed strong antibacterial and antifungal activity and contained many antimicrobially active compounds, it also contained potentially toxic components and hence adequate toxicological data is needed to validate the safety of the stem bark of this plant for consumption.

Keywords: Antimicrobial, Chromatography, Ethanolic, Extract, Fungicidal and Phytochemical

1. Introduction

The uncontrolled use of antibiotics is thought to have spurred evolutionary adaptations that enable microorganisms to survive most of the powerful drugs, thereby encouraging the proliferation of drug resistance among pathogens and normal flora (Schwarz *et al.*, 2001; Rantala *et al.*, 2004; Cabello, 2006; Sande-Bruinsma *et al.*, 2008; Franco, *et al.*, 2009; Susanne and Karen, 2011; Da Costa, *et al.*, 2013; MFMER, 2014). Transfer of R-factor to other bacteria increases the probability of therapeutic failure (Susanne and Karen, 2011; Händel *et al.*, 2015; Hsiao-Han, *et al.*, 2015). Infections from resistant bacteria are now a worldwide problem as some pathogens have even become resistant to multiple types or classes of antimicrobials (CDC, 2013). Consequently, as many existing antimicrobials are becoming less effective, the rapid spread of multi-drug resistant (MDR) bacteria will results to reaching a point where by common everyday infections or diseases could not be treated or prevented (Nemesisbio, 2017). This loss of effective antibiotics will definitely devitalize the means of fighting infectious diseases and management of infectious

complications common in patients with impaired immune system, for which the ability to treat secondary infections is pivotal (CDC, 2013).

It has been shown that the choice of drugs in the treatment of many infections is quite narrow today due to the wide scale resistance that the common pathogens show to drugs which have been used previously (Ahmad *et al.*, 2012). In fact, many drugs which are considered as effective against many pathogens are now rarely prescribed as empirical therapy in areas where resistance rate to those antibiotics is high (Rawat and Umesh, 2010; Shalini *et al.*, 2011). Thus, the process of developing new antimicrobials and new technologies to allow quicker diagnosis and facilitate targeted treatment must be accelerated (BBC, 2014).

The above problems necessitate the urgent need to carry out more research works with a view to finding some newer, safer, and more effective agents that can be used for effective management of infections. Medicinal plants have been the nature's major source of medicines and the search for them have led to the discovery of many compounds with

Volume 6 Issue 12, December 2017

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

antimicrobial potential (Sermakkani *et al.*, 2012). The choice of *B. dalzielii* in this research was due to the high traditional use of the plant, especially the stem bark, in treating many ailments. Although some research works were previously conducted on the stem bark of the plant, little or no work has been done so far using advanced automated techniques to determine the identity of the phytochemical components present in the ethanolic extract of the stem bark of the plant, hence the need to embark on the research and its contribution to knowledge.

2. Materials and Methods

Collection, Identification and Authentication of the Plant Material

The stem bark of the plant, *B. dalzielii*, was collected from the bushes around Kafur Local Government area of Katsina State, Nigeria in the month of February, 2015. It was identified and authenticated at the Herbarium of the Department of Plant Biology, Bayero University, Kano where a voucher specimen was deposited. The stem bark was washed under running tap water, air-dried at room temperature in the laboratory until brittle, and then pulverized to fine powder and stored in airtight glass containers at room temperature in the dark until use as according to technique of Sunday *et al.* (2010) and Salisu *et al.*, (2017).

Extraction of the Crude Extract

The powdered sample of the stem bark was extracted following the method of Salisu *et al.*, (2017).

Fifty grams (50g) of the dried powder of the stem bark of the plant was weighed into a glass container and extracted exhaustively in 500ml of 95% ethanol by percolation method for two weeks during which the sealed bottle was undergoing vigorous shaking at regular intervals. The mixture obtained was filtered through muslin cloth and then re-filtered by passing through Number 1 Whatman's filter paper. The filtrate was concentrated by complete evaporation of the solvent at room temperature. The extract was subsequently transferred into clean sterile airtight glass container and stored in the refrigerator at 4°C before downstream application.

Automated Phytochemical Screening of the Extract

High Performance Liquid Chromatography Analysis (HPLC)

The High Performance Liquid Chromatography Analysis (HPLC) was carried out to determine the number of phytochemicals and their percentages in the extract as follows.

Sample Preparation: Stock solution of the extract was prepared in HPLC grade methanol at a concentration of 100µg mL⁻¹ and stored in a refrigerator until use. The sample was filtered through Whatman No.1 filter paper before undertaking HPLC analysis.

Analysis: HPLC was carried out on the extract as according to Salisu *et al.*, (2017). The analysis was performed using an Agilent Zorbax column (Xdb-C18 Type MG 5 µm, 4.6 × 250

mm). The detection wavelength was 254nm. The analysis was carried out at a constant flow rate of 1.2 mL/ min all throughout, with an average pressure of about 2100 psi. Prior to injection of the sample into the system, the mobile phase was pumped through the column for at least 30 minutes in order to warm up the instrument and achieved equilibration of the column. Elution was carried out with CH₃CN-H₂O. The injection volume was 100 µl. All chromatographic data were recorded and processed using autochro-3000 software.

Fourier Transform Infrared Spectrophotometer (FTIR) Analysis

Fourier Transform Infrared Spectrophotometer (FTIR) Analysis was carried out on the extract using Shimadzu, Japan, FT-IR spectrometer as described by Salisu *et al.*, (2017). A 0.1ml of the sample was dropped into an Arsenic selenide (As₂Se₃) aperture plate of 0.1mm thick and loaded in the FTIR spectroscope. The sample was scanned from 650 to 4000 cm⁻¹ for 16 times to increase the signal to noise ratio. FT-IR spectra were recorded in the absorption range between 650 and 4000 wave number per centimetre at a resolution of 4cm⁻¹. The types of chemical bonds/functional groups present in the extract were identified by interpreting the infrared absorption spectrum using IR spectra Table.

Gas Chromatography- Mass Spectrometry Analysis (GC-MS)

GC-MS of the extract was carried out to further confirm the identity of the phytocomponents following the technique of Salisu *et al.*, (2017). The analysis was carried out using GC-MS (Model: QP 2010 series, Shimadzu, Japan) equipped with a VF-5ms fused silica capillary column of 30m length, 0.25mm diameter and 0.25µm film thickness. The column oven temperature was programmed from 70°C to 200°C for 2°C min⁻¹. Ionization of the sample components was performed in electron impact mode (EI, 70 eV). The temperature of the injector was fixed to 200.00°C and one of the detectors to 250°C. Helium (99.9995% purity) was the carrier gas fixed with a flow rate of 1.80ml min⁻¹. The mass ranges from 40-1000m/z was scanned at a rate of 3.0 scans/s. 1.0µl of the extract was injected with a Hamilton syringe to the GM-MS manually for total ion chromatographic analysis in split injection technique. Total running time of GC-MS was 35min.

Identification of the Constituents: Identity of the various compounds detected by the GC-MS from the extract was carried out based on the comparison of their retention indices and mass spectra fragmentation patterns with those stored in the computer library (i.e the spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST Library) and the interpretation of the mass spectrum GC-MS was conducted using data base of National Institute of Standard Technology (NIST08s), Wiley Registry of Mass Spectral data New York (Wiley 8) and Fatty Acid Methyl Esters Library, version 1.0 (FAME Library) sources were used for matching the detected compounds in the Extract. The molecular weights, molecular formulae and the number of hits used to identify the name of the compounds from NIST and Wiley spectra were also recorded.

Quantification of the Constituents: The relative percentage of each constituent in the extract was expressed as percentage with peak area normalization.

3. Bioassay Studies

Test Organisms: The clinical isolates of bacteria and the fungus used for this study were obtained from Aminu Kano Teaching Hospital, Kano (AKTH). These organisms are as follows: *Staphylococcus aureus*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella typhi* and *Candida albicans*.

Storage and Maintenance of Cultures: The pure cultures of the bacteria were streaked on to Nutrient Agar slants and then incubated at 37°C for 24 hours, while the fungus was streaked on to Sabroud Dextrose Agar slant and incubated at room temperature of 25°C – 28°C for 48 hours. The cultures were stored in the refrigerator at 4°C until needed for subsequent use.

Standardization of Inocula

The inocula of the test organisms were standardized using 0.5 McFarland's turbidity standard. Approximately 99.5ml of 1% BaCl₂ was added to 0.5ml of 1% H₂SO₄ in order to obtain 100ml of BaSO₄ which corresponded to 0.5 McFarland's turbidity standard equivalent to 1.0 X 10⁸ cfu/ml population for bacterial isolates. Bacterial colonies from the 18hours overnight cultures were used to make direct suspension of each bacterial isolate in separate test tubes containing 1ml of sterile physiological saline. The suspensions were adjusted to match the 0.5 McFarland turbidity standard, using saline and a vortex mixer. Similarly, the fungal culture was standardized according to the methods of Murugan *et al.* (2007) by dissolving a loopful of the spore suspension into 10ml of 20% tween 80 solution to obtain 6.0 X 10⁵cfu/ml.

Antimicrobial Susceptibility Test

The antimicrobial activity of the extract was tested using agar well diffusion method adopted from CILS (1998). Müeller Hinton agar plates were prepared for *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa* and *Salmonella typhi*; Müeller Hinton Blood agar plates for *Streptococcus pyogenes*, *Enterococcus faecalis*; and SDA plates for *Candida albicans*.

The plates were separately inoculated with the suspension of the standardized cultures using sterile swabs. In each case, the sterile swab was submerged in the suspension, lifted out, and the excess fluid was removed by pressing and rotating the swab against the wall of the test tube. The swab was then used to inoculate the entire surface of the plate three times, rotating the plate 60 degrees between each inoculation. Each inoculum was allowed to dry for few minutes.

Five wells of 6mm diameter were made in each plate using 6mm sterile cork borer. The wells were filled with 0.1ml of diluted concentrations (50mg/ml, 25mg/ml, 10mg/ml, 5mg/ml and 2.5mg/ml) of the extract with the aid of sterile pipettes per well. Sterile distilled water and 10µg/ml of

Ciprofloxacin were used in separate plates for each organism to serve as negative and positive controls respectively. Diameters of the zones of inhibition were measured with a transparent ruler and the result was recorded in millimetres after incubating the plates at 37°C for 24 hours (bacteria) and 25°C for 48 hours (fungus). The plates were replicated in triplicates and the means and standard errors of the zones of inhibitions for each organism at each concentration of each extract were calculated and recorded accordingly as described by Baha'uddeen *et al.* (2017).

Determination of Minimum Inhibitory Concentration (MIC)

The MIC of the extract was determined by standard two-fold micro dilution method adopted from CILS, (1998). A stock solution of the extract was serially diluted in nine test tubes containing 5mls double strength Mueller Hinton broth each (for bacteria) or SDA broth (for fungus) and labelled 1, 2, 3, 4, 5, 6, 7, 8, and 9; to obtain a concentration of 100, 50, 25, 12.5, 6.25, 3.12, 1.56 and 0.78 mg/ml. Tube 9 was the control and had no drug or extract. 0.1ml of the standardized inocula was inoculated in the separate tubes 1-9 for each organism. The various test tube racks were kept at 37°C for an overnight incubation (bacteria) and 25°C for 48hours (fungus). Then, the MICs were recorded as the lowest concentration of the extract inhibiting the visible growth of test organisms. This was determined by comparing the tubes with the control against a source of light with white back ground and some contrasting black lines.

Determination of Minimum Bactericidal Concentration (MBC) and the Minimum Fungicidal Concentration (MFC)

The MBCs and the MFCs were determined using the methods of Hugo and Russel (1994). The tubes in each set, which did not show any growth during the MIC and MFCs determination, were used. A loop full of the content from each of the required MIC and MFC tubes was streaked unto new nutrient agar plates for the bacteria and SDA plates for the fungus. The plates were incubated for another 24 hours at 37°C (bacteria) and for 48 hours at room temperature of 25°C (fungus). After the incubation period, the plates were examined for growth or otherwise. The MBCs and the MFCs of the extract were recorded as the smallest concentration of the extract that is capable of killing the entire organism present in the specific inoculum size of test organism.

4. Results

A. Physical Characteristics and the Percentage Yield of the Extract

The physical characteristics and the percentage yield of the extract are shown in (Table 1). Both the filtrate and residue appeared reddish in colour, whereas the extract appeared light red in colour with gummy and solid textures. The percentage yield of the extract was 13.4% of the total sample extracted. The extract was found to be soluble in 10% dimethyl sulphoxide (DMSO) and partially soluble in distilled water.

Table 1: Physical Characteristics and the Percentage Yield of the Extract

S/N	Parameters	Observation/Result
1	Weight of the stem bark Powder Extracted	500g
2	Colour of the Filtrate	Red
3	Colour and Texture of the Extract	Red and Gummy
4	Weight of the extract	67.0g
5	% Yield	13.4%

B. Phytochemical Screening

Medicinal plants are generally indispensable at the present age because, being the major sources of many pharmacologically active compounds; they continue to play a dominant role in the maintenance of human health.

The HPLC chromatogram of the crude Ethanolic extract of *B. dalzielii* stem bark showing the HPLC finger prints of the various compounds present are given in (Figure 1). The result showed that the extract contained 8 phytocomponents at different peaks of the chromatogram. Table 2 on the other

hand, showed the detailed tabulation of the HPLC analysis of the extract. From the Table, the major constituents were at peaks 3 (peak area 31.52%), Peak 4 (peak area 21.96%), peak 5 (peak area 17.23%) and peak 2 (peak area 16.21%). While the rest constitute less than 10% by peak area of the extract.

Table 2: Phytochemical Components of the Ethanolic Extract of the Stem Bark of *Boswellia dalzielii* Obtained by HPLC Analysis

Peak Number	Retention Time (min)	Area (%)
1	2.144	4.92
2	2.413	16.21
3	5.868	31.52
4	6.073	21.96
5	6.266	17.23
6	6.593	6.25
7	6.824	0.65
8	12.728	1.23




Figure 1: HPLC Chromatogram of the Crude Ethanolic Extract of *Boswellia dalzielii*

Similarly, the FTIR spectra of the extract showing the various types of chemical bonds/functional groups were given in (Figure 2). On the other hand, Table 3, showed the

interpretation of the FTIR spectral compounds in the extract. From the table, 11 functional groups were identified in the extract including 1°, 2° amines and amides, alkanes, alkenes,

alkynes, alkyl halides, aromatics and aliphatic amines (Table 3).

Table 3: Various Functional Groups/Chemical Bonds Present in the Ethanolic Extract of Stem Bark of *B. dalzielii* Identified by FTIR Analysis.

Peak No.	Wave Length (cm ⁻¹)	Transmittance (t)	Peak Shape	Type of Bond Identified	Functional Group
1	3319	66.919	Curve	N-H stretch	1°, 2° amines, amides
2	2944	76.826	Sharp	C-H stretch	Alkanes
3	2833	77.555	Sharp	C-H stretch	Alkanes
4	2123	99.941	Weak	-C≡C- stretch	Alkynes
5	1655	95.626	Narrow	-C=C- stretch	Alkenes
6	1451	82.221	Sharp	C-C stretch (in-ring)	Aromatics
7	1413	83.512	Narrow	C-C stretch (in-ring)	Aromatics
8	1117	86.879	Sharp	C-N stretch	Aliphatic amines
9	1024	20.675	Very sharp	C-N stretch	Aliphatic amines
10	672	66.003	Weak	C-Br stretch	Alkyl halides
11	657	63.189	Weak	C-Br stretch	Alkyl halides




Figure 2: FTIR Spectra of the Crude Ethanolic Extract of *Boswellia dalzielii*

Also, the total ion chromatograms (TIC) of the extract, showing the GC-MS profile of the compounds identified were given in (Figure 3). The peaks in the chromatograms were integrated and were compared with the database of spectrum of known components stored in the GC-MS NIST library. On the other hand, Table 4 gave the detailed tabulation of GC-MS analysis of the extract in which 13 different constituents have been elucidated and effectively

matched and identified. The major constituents observed were at peaks 4 (n-Hexadecanoic acid with peak area 23.54%), Peak 7 (Oxacyclotetradecan-2-one with peak area 20.33%), peak 8 (Pelargic acid with peak area 16.83%), peak 12 (Methyl (Arachidic alcohol with peak area 13.51%) and peak 9 (Ethyl steriate with peak area 10.05%). While the rest constitute less than 16% composition by peak area.




Figure 3: GC-MS Showing Total Ion Chromatogram (TIC) of Ethanollic Extract of the Stem Bark of *Boswellia dalzielii*

Table 4: Phytochemical Components Identified form the Crude Ethanollic Extract of the Stem Bark of *Boswellia dalzielii* by GC-MS Analysis

Peak Number	Retention Time (minutes)	% Composition by Area	Matched compound IUPAC Name	Chemical Structure
1	9.624	0.20	1-Dodecene	
2	11.721	0.93	3,5-Di-tert-butylphenol	
3	12.30	0.25	1-Hexadecene	
4	18.069	23.54	n-Hexadecanoic acid (Palmitic acid)	
5	18.370	2.43	Ethyl docosanoate	
6	20.100	0.77	9,12-Octadecadienoyl chloride(Linoleic acid Chloride)	
7	21.148	20.33	Oxacyclotetradecan-2-one	
8	21.362	16.83	n-Nonanoic acid (Pelargic acid)	
9	21.850	10.05	Ethyl- octadecanoate (Ethyl stearate)	
10	23.979	3.97	1,E-11,Z-13-Octadecatriene	
11	24.358	4.13	3,5,6,7,8,8a-hexahydro-4,8a-dimethyl-6-(1-methylethenyl)- 2(1H)Naphthalenone	
12	25.574	13.51	n-Eicosanol (Arachidic alcohol)	
13	25.982	3.03	Sulfurous acid, 2-propyl tetradecyl ester	

B. Antimicrobial activity

The result of the antimicrobial activity of the extract showing the means and standard error means of the diameters of zones of inhibition produced by the various concentrations of the extract as well as the MIC and MBC/MFC values of the extract for each of the organisms are shown in Table 5. The result showed that the extract is active against all the test isolates with higher zones of inhibition of 21±0.00mm for *C. albicans* and *S. typhi*, 20±0.80mm for *S. aureus*, and 19±0.50mm for *K.*

pneumoniae, 18±0.80mm for *E. coli* and 18±0.80mm for *P. aeruginosa* at 50mg/ml concentration each. Resistance was only observed at 2.5mg/ml concentration for *S. pyogenes*. Similarly, lowest MIC values of 3.12mg/ml were obtained for *C. albicans* and *S. typhi*, 6.25mg/ml for *S. aureus*, 12.5mg/ml for *K. Pneumoniae*, *E. faecalis* and *P. mirabilis*, 25mg/ml for *E. coli* and *S. pyogenes* and the highest MIC of 50mg/ml for *P. aeruginosa*. The MBC/MFC values did not exceed the corresponding MIC values by more than a factor of 2 (Table 5).

Table 5: Antimicrobial Activity of the Crude Petroleum Ether Extract of the Stem Bark of *Boswellia dalzielii*

S/N	Test Organisms	Means and Standard Error Means of the Diameters of the Zones of Inhibition Produced by the Various Concentrations of the Extract against the test Organisms (mm)					MIC (mg/ml)	MBC/MFC (mg/ml)
		50 mg/ml	25 mg/ml	10 mg/ml	5 mg/ml	2.5 mg/ml		
1	<i>S. aureus</i>	20 ± 0.80	16 ± 0.20	11 ± 0.40	7 ± 0.80	7 ± 0.10	6.25	12.5
2	<i>E. faecalis</i>	15 ± 0.30	11 ± 0.20	9 ± 0.72	9 ± 0.03	8 ± 0.11	12.5	25
3	<i>S. pyogenes</i>	15 ± 0.20	13 ± 0.22	10 ± 0.50	8 ± 0.60	6± 0.00*	25	50
4	<i>P. aeruginosa</i>	17 ± 0.60	9 ± 0.78	9 ± 0.30	7 ± 0.09	7 ± 0.10	50	100
5	<i>K. pneumoniae</i>	19 ± 0.50	14 ± 0.30	10 ± 0.04	9 ± 0.41	8 ± 0.70	12.5	25
6	<i>E. coli</i>	18 ± 0.80	17± 0.03	11 ± 0.60	10 ± 0.35	9 ± 0.10	25	50
7	<i>S. typhi</i>	21 ± 0.00	18 ± 0.40	15 ± 0.06	9 ± 0.20	8 ± 0.70	3.12	12.5
8	<i>P. mirabilis</i>	16 ± 0.20	14 ± 0.80	10 ± 0.21	7 ± 0.50	7 ± 0.80	12.5	25
9	<i>C. albicans</i>	21 ± 0.00	17 ± 0.03	11 ± 0.20	8 ± 0.10	8 ± 0.53	3.12	12.5

Values are means plus or minus standard error means obtained from 3 replicas

Key: * Indicates resistance.,

MIC = Minimum Inhibitory Concentration

MBC = Minimum Bactericidal Concentration,

MFC = Minimum Fungicidal Concentration

5. Discussions

Medicinal plants have been one of the major sources of natural chemotherapeutic agents due to their ability to synthesise and accumulate pharmacologically important secondary metabolites. According to Nandagopalan *et al.*, (2015), knowledge of the chemical constituents of plants is desirable not only for the discovery of therapeutic agents, but also because such information may be of great value in disclosing new sources of economic phytochemical compounds for the synthesis of complex chemical substances and for discovering the actual significance of folkloric remedies. This study was conducted using advanced automated techniques (GC-MS, HPLC and FTIR) to determine the full identity of the phytochemical compounds present in the ethanolic stem bark extract of *B. dalzielii*. In addition, the *in vitro* antimicrobial activity of the extract on some common human pathogenic microorganisms was also determined.

Findings from this study showed that the stem bark of *B. dalzielii* is very rich in pharmacologically active phytochemicals among which are n-Hexadecanoic acid - palmitic acid (peak area 23.5%) which is reported to have an antioxidant, hypocholesterolemic, nematocidal, pesticide, lubricant activities and hemolytic 5-alpha is a reductase inhibitors (Sermakkani *et al.* 2012); 9,12-Octadecadienoyl chloride (Linoleic acid Chloride) which possesses anti-inflammatory, insectifuge, hypocholesterolemic, cancer preventive, nematocidal, hepatoprotective, antihistaminic, antieczemic, antiacne, 5-alpha reductase inhibitor, antiandrogenic, antiarthritic and anticoronary properties (Sermakkani *et al.* 2012) and n-Eicosanol -Arachidic alcohol (peak area 13.51%) which has antimicrobial activity. This finding explains the observed antimicrobial activities of the extract in this study and also corroborates the findings of

reported of Alemika and Oluwole, (1991), Adelakun *et al.*, (2001) and Etuk *et al.*, (2006) who reported that the stem bark of *B. dalzielii* has strong anti-oxidant and antimicrobial activities and contain many pharmacologically active secondary metabolites such as saponins, tannins, flavonoids, cardiac glycosides, steroids and terpenes.

In addition, the extract also contain Oxacyclotetradecan-2-one (peak area 20.33%) which is the second most abundant phytochemicals identified in the extract. This compound is an important component of Erythromycin and Roxithromycin antibiotics. This further supports the strong antimicrobial activity exhibited by the stem bark of *B. dalzielii*.

However, it should be noted that, although the extract contained many medicinally important constituents, it contain some toxic compounds such as n-nonanoic acid-Pelargic acid (peak area 16.83%) which was reported as being highly corrosive disinfectant and algacide that is commonly used in conjunction with glyphosate, a non-selective herbicide, for a quick burn-down effect in the control of weeds in turf grass and in the preparation of plasticizers and lacquers (Regulation (EU) n°528, 2012). It also contained Ethyl steriate (peak area 10.05%) which was reported to cause skin, eye, respiratory and digestive tract irritations (MSDS, 1998). The presence of these toxic compounds corroborates the findings of Abdalazeez *et al.*, (2013) that reported mild toxicity of aqueous stem bark extract of *B. dalzielii*. Further research is therefore needed to validate the safety of the stem bark of *B. dalzielii* for consumption.

On the other hand, the results of the antimicrobial activity, in this study, showed that the extract has broad spectrum of

activity against the gram negative and gram positive bacteria as well as the fungal isolates used in the study. The result therefore, corroborates the findings of Alemika and Oluwole, (1991); Duwiejua *et al.*, (1993); Adelakun *et al.*, (2001) and Abdulazeez *et al.*, (2013) who reported that *B. dalzielii* stem bark has broad spectrum of antimicrobial activity.

However, *Pseudomonas aeruginosa* appeared to have the highest MIC (50mg/ml) and MBC (100mg/ml). This might not be unconnected with the fact that *P. aeruginosa* is one of the notable drug resistant organisms due to combination of many factors such as low permeability of its cell wall, its genetic capacity to express a wide repertoire of resistance mechanisms, mutation in chromosomal genes which regulate resistance genes, or acquisition of additional resistance genes from other organisms via plasmids, transposons and bacteriophages as reported by Lambert, (2002).

Furthermore, the fact that none of the organisms used in this study was completely resistant to the extract implies that, the stem bark of *B. dalzielii* can be used in the treatment of many infections caused by the organisms especially those caused by *S. typhi* and *C. albicans* as they appeared to be the most susceptible organisms with MIC of 3.12mg/ml each.

6. Conclusion

The crude ethanolic extract of stem bark of *Boswellia dalzielii* has strong antibacterial and antifungal activity and contained many antimicrobially active compounds. However, further research is needed to separate these bioactive compounds from the potentially toxic components of the stem bark identified in this study. In addition, adequate toxicological data is needed to validate the safety of the stem bark of this plant for consumption purpose. Furthermore, metabolomic study of the extract is also recommended in future to identify the metabolites present in the active compounds identified in this research as well as their systemic functions and effects on human consumption.

7. Acknowledgement

We acknowledge the management of Umaru Musa Yar'adua University for their support and TETFUND Nigeria for funding the research.

8. Conflict of Interest

We declare none

References

- [1] Abdulazeez A. T, Kabele-Toge B., Lawal M. and Abubakar M. G. (2013). Phytochemical, Antibacterial and Toxicological Studies of Aqueous Stem Bark Extract of *Boswellia dalzielii*. African Journal of Pharmaceutical Research & Development, Vol. 5 No.1 pp.61-67.
- [2] Adelakun E.A., Finbar E.A.V., Agina S.E. and Makinde A.A. (2001): Antimicrobial activity of *Boswellia dalzielii* stem bark; *Fitoterapia* 72(7), 822-824.
- [3] Ahmed Salih Sahib, Imad Hashim Mohammed, Saba Jasim Hamdan. (2012). Use of aqueous extract of corn silk in the treatment of urinary tract infection. *Journal of Intercultural Ethnopharmacology*;1(2):93-96. Available at www.scopemed.org
- [4] Alemika T.O.E. and Oluwole F.S. (1991): An investigation of the potentials of *Boswellia dalzielii* and *Commiphora kerstingii* in the treatment of peptic ulcer; *West African Journal of Pharmacology & Drug Research.*, 9/10, 91-94.
- [5] BBC News. Antibiotic resistance rise continues. 2014. Available at: <http://www.bbc.co.uk/news/health-29553435> (accessed 19.03.2015).
- [6] Cabello F. Heavy use of prophylactic antibiotics in aquaculture: A growing problem for human and animal health and for the environment. *Environ. Microbiol.* 2006;8:1137–1144. doi: 10.1111/j.1462-2920.2006.01054.x.
- [7] Centers for Disease Control and Prevention (2013). Meeting the Challenges of Drug-Resistant Diseases in Developing Countries Committee on Foreign Affairs Subcommittee on Africa, Global Health, Human Rights, and International Organizations, United States House of Representatives. April 23, 2013.
- [8] CILS and Antimicrobial Susceptibility Testing. (1998). Performance Standards for Antimicrobial Susceptibility Testing. In Blood Safety and Clinical Technology Guidelines on Standard Operating Procedures for Microbiology. Eight Informational Supplement. M100-S8, 18 (1), CILS Pennsylvania, USA.
- [9] Da Costa, P. M., Loureiro, L., & Matos, A. J. F. (2013). Transfer of Multidrug-Resistant Bacteria between Intermingled Ecological Niches: The Interface between Humans, Animals and the Environment. *International Journal of Environmental Research and Public Health*, 10(1), 278–294. <http://doi.org/10.3390/ijerph10010278>
- [10] Duwiejua M., Zeitlin I.J., Waterman P.G., Chapman J., Mhango G.J. and Provan G.J. (1993): Anti-inflammatory activity of resins from some of the plant family Burseraceae; *Plant Medicine.*, 59(1), 12-16.
- [11] Etuk E U, Agaie B M, Onyeyili P A, Ottah C U. (2006). Toxicological studies of aqueous stem bark extract of *Boswellia dalzielii* in albino rats. *Indian J Pharmacology*;38:359-60. Available from: <http://www.ijp-online.com/text.asp?2006/38/5/359/27707>
- [12] Franco, B. E., Altagracia Martínez, M., Sánchez Rodríguez, M. A., & Wertheimer, A. I. (2009). The determinants of the antibiotic resistance process. *Infection and Drug Resistance*, 2, 1–11.
- [13] Hugo and Russell (2004). *Pharmaceutical microbiology*, 8th edi, edited by Stephen P. D., Norman A. H and Sean P. G., 11(187-201).
- [14] Händel N, Otte S, Jonker M, Brul S, ter Kuile BH (2015) Factors That Affect Transfer of the Inc11 β -Lactam Resistance Plasmid pESBL-283 between *E. coli* Strains. *PLoS ONE* 10(4): e0123039. <https://doi.org/10.1371/journal.pone.0123039>
- [15] Hsiao-Han Chang, Ted Cohen, Yonatan H. Grad, William P. Hanage, Thomas F. O'Brien and Marc Lipsitch. (2015). Origin and Proliferation of Multiple-Drug Resistance in Bacterial Pathogens. *Microbiology and Molecular Biology Rev.* vol. 79 no. 1 101-116

- [16] Lambert P.A. (2002). Mechanisms of antibiotic resistance in *Pseudomonas aeruginosa*; Section of Paediatrics & Child Health. *Journal of research in soc medicine* ;95(suppl. 41):22–26
- [17] Material Safety Data Sheet (1998). Ethyl stearate, 99%, ACC# 05986. Revised on 9/26/2007, assessed on 17/7/2016.
- [18] MFMER (Mayo Foundation for Medical Education and Research). (2014). Antibiotics: Misuse puts you and others at risk. Available at <https://www.mayoclinic.org/healthy-lifestyle/consumer-health/in-depth/antibiotics/art-20045720>. (Accessed 16/12/2017).
- [19] Murugan, K., Saravanababu, S., Arnachalam, M. (2007). Screening of tannin acyl hydrolase (E. C. 3.1.1.20) producing tannery effluent fungal isolates using simple agar plate and smF process. *Bioresources and Technology*, 98: 946–949.
- [20] Nandagopalan, V., Gritto, M. J., & Doss, A. (2015). GC-MS analysis of bioactive components of the methanol extract of *Hibiscus tiliaceus* Linn. *Asian Journal of Plant Science and Research*, 5(3), 6–10. Retrieved from www.pelagiaresearchlibrary.com
- [21] Nemesis Bioscience News (2017). Available at <http://www.nemesisbio.com/the-problem/> (accessed 16-12-2017)
- [22] Rantala M., Hölsö K., Lillas A., Huovinen P., Kaartinen L. Survey of condition-based prescribing of antimicrobial drugs for dogs at a veterinary teaching hospital. *Vet. Rec.* 2004;155:259–262. doi: 10.1136/vr.155.9.259.
- [23] Rawat, Vinita and Umesh Paul, Priyanka. (2010). Antibiotic resistance pattern of urinary tract isolates of *Escherichia coli* from Kumaun region. *The Journal of communicable diseases*. 42. 63-6
- [24] Regulation (EU) n°528/2012 concerning the making available on the market and use of biocidal products. Nonanoic acid Product-type 2 *Assessment report finalised on 27/09/2013*, assessed on 27/7/2016
- [25] Salisu, B., Magashi, A. M., & Mohammed, A. (2017). Determination of Phytochemicals and Antimicrobial Activity of Aqueous Stem Bark Extract of *Boswellia dalzielii* against Some Common Pathogenic Microorganisms. *UMYU Journal of Microbiology Research*, 2(1), 238–246. Retrieved from www.umyu.edu.ng
- [26] Sande-Bruinsma N., Grundmann H., Verloo D., Tiemersma E., Monen J., Goossens H., Ferech M. Antimicrobial drug use and resistance in Europe. *Emerg. Infect. Dis.* 2008;11:1722–1730.
- [27] Schwarz S., Kehrenberg C., Walsh T. Use of antimicrobial agents in veterinary medicine and food animal production. *Int. J. Antimicrob. Agents.* 2001;17:431–437. doi: 10.1016/S0924-8579(01)00297-7.
- [28] Sermakkani M & Thangapandian V. (2012). GCMS analysis of *Cassia italic* leaf methanol extract. *Asian journal of pharmaceutical and clinical research* .vol 5.
- [29] Shalini., Joshi, MC., Rashid, MK., and Joshi, HS. 2011. Study of Antibiotic Sensitivity Pattern In Urinary Tract Infection At A Tertiary Hospital. *NJIRM*; Vol. 2(3): 43-46.
- [30] Sunday Ene-Ojo, Atawodi Gboyega and Suleiman Onaolapo (2010). Comparative in vitro antioxidant potential of different parts of *Ipomoea asarifolia*, Roemer & Schultes, *Guiera senegalensis*, J. F. Gmel and *Anisopus mannii* N. E. Brown. *Braz. J. Pharm. Sci.* vol.46 no.2 São Paulo Apr./June 2010 ,<http://dx.doi.org/10.1590/S1984-82502010000200011>
- [31] Susanne Schjørring and Karen A. Krogfelt, “Assessment of Bacterial Antibiotic Resistance Transfer in the Gut,” *International Journal of Microbiology*, vol. 2011, Article ID 312956, 10 pages, 2011. doi:10.1155/2011/312956