

k-Super Root Square Mean Labeling of Some Graphs

Akilandeswari. K

¹Research Scholar, Department of Mathematics, Seethalakshmi Ramaswami College, Tiruchirappalli – 620002, India

Abstract: Let G be a (p, q) graph and $f: V(G) \rightarrow \{k, k+1, k+2, \dots, p+q+k-1\}$ be an injection. For each edge $e = uv$, let $f^*(e) = \left\lfloor \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rfloor$ or $\left\lceil \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rceil$, then f is called k -Super root square mean labeling if $f(V) \cup \{f^*(e) : e \in E(G)\} = \{k, k+1, k+2, \dots, p+q+k-1\}$. A graph that admits a k -Super root square mean labeling is called k -Super root square mean graph. In this paper, we investigate k -Super root square mean labeling of some path related graphs.

Keywords: Super root square mean labeling, Super root square mean graph, k -Super root square mean labeling, k -Super root square mean graph, $P_n \odot K_{1,2}$, $P_n \odot K_{1,3}$, $D(T_n)$, $Q_n \odot K_1$.

1. Introduction

All graphs in this paper are finite, simple and undirected. Terms not defined here are used in the sense of Harary [5]. The symbols $V(G)$ and $E(G)$ will denote the vertex set and edge set of a graph G .

Graph labeling was first introduced in the late 1960's. Many studies in graph labeling refer to Rosa's research in 1967 [6]. Labeled graphs serve as useful models for a broad range of applications such as X-ray, crystallography, radar, coding theory, astronomy, circuit design and communication network addressing. Particularly interesting applications of graph labeling can be found in [1-4].

The concept of mean labeling was introduced and studied by S. Somasundaram and R. Ponraj [7].

Root square mean labeling was introduced by S.S. Sandhya, R. Ponraj and S. Anusa [8].

The concept of super root square mean labeling was introduced and studied by K. Thirugnanasambandam et al. [9].

In this paper, I introduce k -Super root square mean labeling and investigate k -Super root square mean labeling of $P_n \odot K_{1,2}$, $P_n \odot K_{1,3}$, $D(T_n)$, $Q_n \odot K_1$. Throughout this paper, k denote any positive integer greater than or equal to 1.

For brevity, we use k -SRSML for k -Super root square mean labeling.

Definition 1.1

Let G be a (p, q) graph and $f: V(G) \rightarrow \{1, 2, \dots, p+q\}$ be an injection. For each edge $e = uv$, let $f^*(e) = \left\lfloor \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rfloor$ or $\left\lceil \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rceil$, then f is called **Super root square mean labeling** if $f(V) \cup \{f^*(e) : e \in E(G)\} = \{1, 2, \dots, p+q\}$. A

graph that admits a Super root square mean labeling is called **Super root square mean graph**.

Definition 1.2

Let G be a (p, q) graph and $f: V(G) \rightarrow \{k, k+1, k+2, \dots, p+q+k-1\}$ be an injection. For each edge $e = uv$, let $f^*(e) = \left\lfloor \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rfloor$ or $\left\lceil \sqrt{\frac{f(u)^2 + f(v)^2}{2}} \right\rceil$, then f is called **k -Super root square mean labeling** if $f(V) \cup \{f^*(e) : e \in E(G)\} = \{k, k+1, k+2, \dots, p+q+k-1\}$.

A graph that admits a k -Super root square mean labeling is called **k -Super root square mean graph**.

Definition 1.3:

If G has order n , the corona of G with H , $G \odot H$ is the graph obtained by taking one copy of G and n copies of H and joining the i th vertex of G with an edge to every vertex in the i th copy of H .

Definition 1.4:

A Double Triangular Snake $D(T_n)$ consists of two triangular snakes that have a common path.

2. Main Results

Theorem 2.1:

The graph $P_n \odot K_{1,2}$ is a k -Super root square mean graph for $n \geq 2$.

Proof:

Let $V(P_n \odot K_{1,2}) = \{u_i, v_i, w_i; 1 \leq i \leq n\}$ and $E(P_n \odot K_{1,2}) = \{e_i = (u_i, u_{i+1}); 1 \leq i \leq n-1\} \cup \{e'_i = (u_i, v_i); 1 \leq i \leq n\} \cup \{e''_i = (u_i, w_i); 1 \leq i \leq n\}$

be the vertices and edges of $P_n \odot K_{1,2}$ respectively.

Define

$$f: V(P_n \odot K_{1,2}) \rightarrow \{k, k+1, k+2, \dots, k+6n-2\}$$

$$f(u_i) = k+6i-4; 1 \leq i \leq n,$$

$$f(v_i) = k+6i-6; 1 \leq i \leq n,$$

$$f(w_i) = k+6i-2; 1 \leq i \leq n.$$

Now the induced edge labels are as follows:

$$f^*(e_i) = k+6i-1; 1 \leq i \leq n-1,$$

$$f^*(e'_i) = k+6i-5; 1 \leq i \leq n,$$

$$f^*(e''_i) = k+6i-3; 1 \leq i \leq n.$$

Here $p = 3n$ and $q = 3n-1$.

Clearly, $f(V) \cup \{f^*(e) : e \in E(P_n \odot K_{1,2})\} = \{k, k+1, \dots, k+6n-2\}$.

So f is a k -Super root square mean labeling.

Hence $P_n \odot K_{1,2}$ is a k -Super root square mean graph.

Example 2.2:

100- SRSML of $P_3 \odot K_{1,2}$ is given in figure 2.1:




Figure 2.1: 100-SRSML of $P_3 \odot K_{1,2}$

Theorem 2.3:

The graph $P_n \odot K_{1,2}$ is a k -Super root square mean graph for $n \geq 2$.

Proof:

Let $V(P_n \odot K_{1,2}) = \{u_i, v_i, w_i, s_i; 1 \leq i \leq n\}$ and $E(P_n \odot K_{1,2}) = \{e_i = (u_i, u_{i+1}); 1 \leq i \leq n-1\} \cup \{e'_i = (u_i, v_i); 1 \leq i \leq n\} \cup \{e''_i = (u_i, w_i); 1 \leq i \leq n\} \cup \{e'''_i = (u_i, s_i); 1 \leq i \leq n\}$ be the vertices and edges of $P_n \odot K_{1,2}$ respectively.

Define

$$f: V(P_n \odot K_{1,2}) \rightarrow \{k, k+1, k+2, \dots, k+8n-2\}$$

$$f(u_i) = k+8i-6; 1 \leq i \leq n,$$

$$f(v_i) = k+8i-8; 1 \leq i \leq n,$$

$$f(w_i) = k+8i-4; 1 \leq i \leq n,$$

$$f(s_i) = k+8i-2; 1 \leq i \leq n$$

Now the induced edge labels are as follows:

$$f^*(e_i) = k+8i-1; 1 \leq i \leq n-1,$$

$$f^*(e'_i) = k+8i-7; 1 \leq i \leq n,$$

$$f^*(e''_i) = k+8i-5; 1 \leq i \leq n$$

$$f^*(e'''_i) = k+8i-3; 1 \leq i \leq n.$$

Here $p = 4n$ and $q = 4n-1$.

Clearly, $f(V) \cup \{f^*(e) : e \in E(P_n \odot K_{1,2})\} =$

$$\{k, k+1, \dots, k+8n-2\}.$$

So f is a k -Super root square mean labeling.

Hence $P_n \odot K_{1,2}$ is a k -Super root square mean graph.

Example 2.4:

200- SRSML of $P_3 \odot K_{1,3}$ is given in figure 2.2:




Figure 2.2: 200-SRSML of $P_3 \odot K_{1,3}$

Theorem 2.5:

The graph $Q_n \odot K_1$ is a super root square mean graph for $n \geq 2$.

Proof:

Let $V(Q_n \odot K_1) = \{u_i, u'_i; 1 \leq i \leq n\} \cup \{v_i, v'_i, w_i, w'_i; 1 \leq i \leq n-1\}$ and $E(Q_n \odot K_1) = \{e_i = (u_i, u_{i+1}); 1 \leq i \leq n-1\} \cup \{e'_i = (u_i, u'_i); 1 \leq i \leq n\} \cup \{e''_i = (u_i, v_i); 1 \leq i \leq n-1\} \cup \{e'''_i = (u_{i+1}, w_i); 1 \leq i \leq n-1\} \cup \{e^{iv}_i = (u_i, w_i); 1 \leq i \leq n-1\} \cup \{e^{iv}_i = (v_i, v'_i); 1 \leq i \leq n-1\} \cup \{e^{vi}_i = (w_i, w'_i); 1 \leq i \leq n-1\}$

be the vertices and edges of $Q_n \odot K_1$ respectively.

Define

$$f: V(Q_n \odot K_1) \rightarrow \{k, k+1, k+2, \dots, k+13n-11\}$$

$$f(u_i) = k+13i-11; 1 \leq i \leq n,$$

$$f(v_i) = k+13i-9; 1 \leq i \leq n-1,$$

$$f(w_i) = k+10,$$

$$f(w'_i) = k+13i-2; 2 \leq i \leq n-1,$$

$$f(u'_i) = k,$$

$$f(u''_i) = k+13i-14; 2 \leq i \leq n,$$

$$f(v'_i) = k+13i-7; 1 \leq i \leq n-1,$$

$$f(w'_i) = k+13i-5; 1 \leq i \leq n-1.$$

Now the induced edge labels are as follows:

$$f^*(e_1) = k+11,$$

$$f^*(e_i) = k+13i-3; 2 \leq i \leq n-1,$$

$$f^*(e'_i) = k+13i-12; 1 \leq i \leq n,$$

$$f^*(e''_i) = k+13i-10; 1 \leq i \leq n-1,$$

$$f^*(e'''_i) = k+13i; 1 \leq i \leq n-1,$$

$$f^*(e^{iv}_i) = k+13i-6; 1 \leq i \leq n-1,$$

$$f^*(e^{iv}_i) = k+13i-8; 1 \leq i \leq n-1,$$

$$f^*(e^{vi}_i) = k+13i-4; 1 \leq i \leq n-1.$$

Here $p = 6n-4$ and $q = 7n-6$.

Clearly, $f(V) \cup \{f^*(e) : e \in E(Q_n \odot K_1)\} = \{k, k+1, k+2, \dots, k+13n-11\}$.

So f is a k -Super root square mean labeling.

Hence $Q_n \odot K_1$ is a k-Super root square mean graph.

Example 2.6:

250- SRSML of $Q_4 \odot K_1$ is given in figure 2.3:




Figure 2.3: 250-SRSML of $Q_4 \odot K_1$

Theorem 2.7:

Any Double Triangular Snake $D(T_n)$ is a k-Super root square mean labeling.

Proof:

Let $V(D(T_n)) = \{u_i; 1 \leq i \leq n\} \cup \{v_i, w_i; 1 \leq i \leq n-1\}$

and

$$E(D(T_n)) = \{e_i = (u_i, u_{i+1}); 1 \leq i \leq n-1\} \cup \\ \{e'_i = (u_i, v_i); 1 \leq i \leq n-1\} \cup \\ \{e''_i = (u_{i+1}, v_i); 1 \leq i \leq n-1\} \cup \\ \{e'''_i = (u_i, w_i); 1 \leq i \leq n-1\} \cup \\ \{e^{iv}_i = (u_{i+1}, w_i); 1 \leq i \leq n-1\}$$

be the vertices and edges of $D(T_n)$ respectively.

Define $f: V(D(T_n)) \rightarrow \{k, k+1, k+2, \dots, k+8n-8\}$ by

$$f(u_i) = k + 8i - 8; 1 \leq i \leq n, \\ f(v_i) = k + 8i - 6; 1 \leq i \leq n-1, \\ f(w_i) = k + 8i - 4; 1 \leq i \leq n-1$$

Now the induced edge labels are

$$f^*(e_i) = k + 8i - 3; 1 \leq i \leq n-1, \\ f^*(e'_i) = k + 8i - 7; 1 \leq i \leq n-1, \\ f^*(e''_i) = k + 8i - 2; 1 \leq i \leq n-1, \\ f^*(e'''_i) = k + 8i - 5; 1 \leq i \leq n-1, \\ f^*(e^{iv}_i) = k + 8i - 1; 1 \leq i \leq n-1$$

Here $p = 3n-2$ and $q = 5n-5$.

Clearly, $f(V) \cup \{f^*(e); e \in E(D(T_n))\} =$

$$\{k, k+1, k+2, \dots, k+8n-8\}.$$

So f is a k-Super root square mean labeling.

Hence $D(T_n)$ is a k-Super root square mean graph.

Example 2.8:

10- SRSML of $D(T_5)$ is given in figure 2.4:




Figure 2.4: 10-SRSML of $D(T_5)$

References

- [1] G.S. Bloom, S.W. Golomb, Applications of numbered undirected graphs, Proc. IEEE, 65 (1977), 562-570.
- [2] G.S. Bloom, S.W. Golomb, Numbered complete graphs unusual rulers and assorted applications, Theory and Applications of Graphs-Lecture notes in Math., Springer Verlag, New York, 642 (1978), 53-65.
- [3] G.S. Bloom, D.F. Hsu, On graceful digraphs and a problem in network addressing, Congressus Numerantium, 35 (1982) 91-103.
- [4] J.A. Gallian, A dynamic survey of graph labeling, Electronic Journal of Combinatorics, 18 (2015) # DS6.
- [5] F. Harary, Graph Theory, Addison Wesley, Massachusetts (1972).
- [6] Rosa, On certain valuations of the vertices of a graph Theory of Graphs (Internet Symposium, Rome, July (1966), Gordon and Breach, N.Y. and Duhod, Paris (1967) 349-355.
- [7] S. Somasundaram and R. Ponraj, Mean labeling of graphs, National Academy Science Letter, 26 (2003), 210-213.
- [8] S. Somasundaram, S.S. Sandhya and S. Anusa, Root Square Mean Labeling of Graphs, Int. J. Comtemp. Math. Sciences, Vol. 9, 14 (2014), 667-676.
- [9] K. Thirugnanasambandam and K. Venkatesan, Super Root Square Mean Labeling of Graphs, Int. J. of Mat. and Soft Computing, Vol 5, No. 2 (2015), 189-195.