

A Study on Avifaunal Species Diversity of Purbasthali Oxbow Lake, West Bengal, India

Santanu Chowdhury

Department of Environmental Science, Nabadwip Vidyasagar College, Nabadwip, Nadia-741302, West Bengal, India

Abstract: *The present study deals with the diversity, abundance and variations of avian species at Purbasthali Oxbow Lake (23°27'5"N 88°20'35"E), West Bengal, India. A total number of 89 bird species belonging to 15 orders and 36 families were recorded during this study. Among the recorded species 47 were resident, 14 were local migrant and 28 were winter migratory. All the winter migrants such as Northern Pintail, Lesser Whistling Duck, Red crested Pochard, Common Teal, Spot Billed Duck, Garganey, Gadwal etc. were observed in large numbers during winter season. It was found that one near threatened (Black headed Ibis) and one vulnerable (common pochard) species also regularly visited this lake in the winter season. The results of relative diversity (RD) index showed that anatidae (RD value = 16.85) was the dominant family in the area.*

Keywords: Avian, diversity, wetland, Purbasthali oxbow lake, migratory

1. Introduction

Avifaunal species are one of the main indicators which determine the health of the wetlands [1-3]. Bird species play a significant role in many food webs of aquatic system through nutrient cycling and as a part of food web, as potential pollinators and bio-indicators [4-5]. Wetlands are important bird habitats for feeding, nesting and breeding of aquatic birds [6-8]. Now-a-days, avifaunal diversity has been decreasing due to the destruction of natural habitats and anthropogenic interference. The wetlands are facing tremendous anthropogenic pressure, which can greatly influence the population structure of the bird community [9-12].

Purbasthali Oxbow Lake also known as Chupi Char is created by the Ganges River on its Western bank, in Burdwan district of West Bengal, India. This lake harbors a number of aquatic plants in the submerged as well as floating state, on which thrive a large number of organisms. Due to abundant food available throughout the year in the form of aquatic crustaceans, insects, mollusks, fishes etc. the lake attracts a number of birds throughout year. This beautiful lake harbour large populations of migratory water birds during the winter season. Information on distribution and abundance of water birds is essential to provide guidelines for the management and conservation of wetlands

[13-15]. The present study was conducted to analyze the diversity and richness of wetland birds and to identify the consequences of direct and indirect human interferences.

2. Material and Method

Study area

The study area is Purbasthali Oxbow lake (88°19'45" to 88°22' E longitude 23° 26' to 23°26'45" N latitude) also known as Chupi Char (**Fig 1**) created by the Ganges river on its Western bank, in Burdwan district of West Bengal, India. This area is only 8 km from the old and holy town of Nabadwip. The lake was formed by the meandering river, over last 40 years, the area has transformed into a closed loop, allowing emergence of the oxbow lake. This channel of water course feeds the oxbow lake with thin connectivity with the main river with shoals forming at the river mouth. Remote sensing images of the wetland clearly establishes the differences of turbidity between the main river and the wetland which has sandy clay sediment and crystal clear waters because of sedimentation of suspended solid particles in the stagnant stretches. The ongoing sedimentation process threatens to cut off the channel in near future. The oxbow lake of Purbasthali sprawls over an area 3.50 km². The water depth of this lake varies between is 1.0 m to 4.5m.



Figure 1: Map of the study area

Methodology

Avifauna in and around Purbasthali Oxbow Lake was recorded during January 2013 to December 2016. Sampling was carried out for three years to record variation in avifaunal diversity. Regular field trips were made throughout this period. Two different methods were adapted to study avifaunal diversity. The first method was Line transects method and second method was Point count method. Following these methods checklist was prepared. Olympus Binocular 10x50, was used for close observation of birds and for photography Cannon-EOS 550 D camera, with sigma Lens 150-500 mm. The check list of species was prepared following Ali [16], Grimmett and Inskipp [17]. The other most important aspect kept in consideration was to make the observations during the peak activity of birds, which is 1 or 2 hours after sunrise or before sunset. The following formula was used for determining percentage of occurrence or relative diversity (RD) of Families [18].

$$\text{Relative diversity (RD)} = \frac{\text{No of species of each family}}{\text{Total no of species}} \times 100$$

3. Result and Discussion

The checklist of observed avian species in and around Purbasthali oxbow lake along with their order, family, common name, scientific name and residential status are given in **Table 1**. In total, 89 species of birds belonging to 15 orders and 36 families were observed. Out of total 89 species of birds, 52.8% were Resident, 31.4% were winter Migrant and 15.7% Local Migrant (**Fig 2**). Order Passeriformes is dominant in the study area, including 13 families and 23 species (25.8%), followed by order Charadriiformes (17.9%) and Anseriformes (16.8%). Family Anatidae (16.8%) are dominant with 15 no of species, followed by Scolopacidae (7.8%) and Ardeidae (6.7%). Avian diversity in terms of different order is given in **Fig.3**. Anatidae was found to be the most dominant family in the area (RD Index value = 16.85) followed by Scolopacidae (RD Index value = 8.98), Rallidae and Sturnidae (RD Index value = 5.61). One near threatened (Black headed ibis) and one vulnerable (Common Pochard) species according to IUCN red data book were found in this area during winter season. Remaining all other species found during this survey are categorized as least concerned according to IUCN red data book. The bird's population changes in this lake in different seasons due to local environmentally dependant factors, agricultural activities, water availability, local & regional habitat changes and climatic conditions [19-20].

Table 1: A classified chart of various bird species in the study area

No	Family	Common Name	Scientific Name	Abundance	Residential Status	IUCN Status
Order : Charadriiformes						
1	Charadriidae	Red-Wattled Lapwing	<i>Vanellu sindicus</i>	C	LM	LC
2		Yellow-Walted Lapwing	<i>Vanellus malabaricus</i>	C	LM	LC
3		Grey-headed Lapwing	<i>Vanellus cinereus</i>	C	WM	LC
4		Little Ringed Plover	<i>Charadrius dubius</i>	R	WM	LC
5	Jacaniidae	Bronze winged jacana	<i>Metopidius indicus</i>	A	RE	LC
6		Pheasant tailed jacana	<i>Hydrophasianus chirurgus</i>	A	RE	LC
7	Scolopacidae	Common redshank	<i>Tringa totanus</i>	C	LM	LC
8		Common greenshank	<i>Tringa nebularia</i>	C	LM	LC
9		wood sandpiper	<i>Tringa glareola</i>	C	LM	LC
10		marsh sandpiper	<i>Tringa stagnatilis</i>	C	LM	LC

11		Green sandpiper	<i>Tringa ochropus</i>	C	WM	LC
12		Common sandpiper	<i>Actitis hypoleucos</i>	A	LM	LC
13		little stint	<i>Calidris minuta</i>	R	WM	LC
14		Common Snipe	<i>Gallinago gallinago</i>	C	WM	LC
15	Recurvirostridae	Black-winged stilt	<i>Himantopus himantopus</i>	O	WM	LC
16	Glareolidae	Small pratincole	<i>Glareola lactea</i>	R	WM	LC
Order : Suliformes						
17	Phalacrocoracidae	Little Cormorant	<i>Phalacrocorax niger</i>	A	RE	LC
18		Indian Cormorant	<i>Phalacrocorax carbo</i>	R	LM	LC
Order: Ciconiiformes						
19	Halcyonidae	White-Breasted Kingfisher	<i>Halcyon smyrnensis</i>	A	RE	LC
20		Stork-billed Kingfisher	<i>Plargopsis capensis</i>	C	RE	LC
21	Cerylidae	Pied Kingfisher	<i>Ceryle rudis</i>	C	RE	LC
22		Common Kingfisher	<i>Alcedo atthis</i>	C	RE	LC
23	Meropidae	Green Bee eater	<i>Merops orientalis</i>	C	RE	LC
24	Coraciidae	Indian roller	<i>Coracias benghalensis</i>	C	RE	LC
Order : Gruiformes						
25	Rallidae	White Breasted Waterhen	<i>Amaurornis phoenicurus</i>	C	RE	LC
26		Purple moorhen	<i>Porphyrio porphyrio</i>	A	RE	LC
27		Common Moorhen	<i>Gallinula chloropus</i>	A	RE	LC
28		Eurasian coot	<i>Fulica atra</i>	A	WM	LC
29		Baillon's crane	<i>Porzana pusilla</i>	R	WM	LC
Order : Anseriformes						
30	Anatidae	Ruddy Shelduck	<i>Tadorna ferruginea</i>	R	WM	LC
31		Eurasian Teal	<i>Anas crecca</i>	R	WM	LC
32		Cotton pigmy goose	<i>Nettion coromandelianus</i>	A	WM	LC
33		Gadwall	<i>Anas strepera</i>	C	WM	LC
34		Garganey	<i>Anas querquedula</i>	C	WM	LC
35		Ferruginous duck	<i>Aythya nyroca</i>	A	WM	LC
36		Tufted duck	<i>Aythya fuligula</i>	C	WM	LC
37		Common Pochard	<i>Aythya ferina</i>	C	WM	VU
38		Northern Shoveler	<i>Anas clypeata</i>	R	WM	LC
39		Northern Pintail	<i>Anas acuta</i>	C	WM	LC
40		Spot billed duck	<i>Anas poecilorhyncha</i>	C	WM	LC
41		Red crested pochard	<i>Nettion rufina</i>	A	WM	LC
42		Lesser Whistling Duck	<i>Dendrocygna javanica</i>	A	WM	LC
43		Knob-billed duck	<i>Sarkidiornis melanotos</i>	O	WM	LC
44		Eurasian wigeon	<i>Anas penelope</i>	O	WM	LC
Order : Pelecaniformes						
45	Threskiornithidae	Black headed ibis	<i>Threskiornis melanocephalus</i>	C	WM	NT
46		Glossy Ibis	<i>Plegadis falcinellus</i>	O	WM	LC
47	Ardeidae	Purple Heron	<i>Ardea purpurea</i>	C	RE	LC
48		yellow bittern	<i>Ixobrychus sinensis</i>	R	WM	LC
49		Cattle Egret	<i>Bubulcus ibis</i>	A	RE	LC
50		Little Egret	<i>Egretta garzetta</i>	A	RE	LC
51		Great Egret	<i>Casmerodius albus</i>	C	RE	LC
52		Indian Pond Heron	<i>Ardeola grayii</i>	A	RE	LC
Order : Podicipediformes						
53	Podicipedidae	Little grebe	<i>Tachybaptus ruficollis</i>	A	WM	LC
54		Great crested grebe	<i>Podiceps cristatus</i>	O	WM	LC
Order : Passeriformes						
55	Motacillidae	White Wagtail	<i>Motacilla alba</i>	C	RE	LC
56		Citrine Wagtail	<i>Motacilla citreola</i>	C	LM	LC
57		yellow wagtail	<i>Motacilla flava</i>	C	LM	LC
58		Paddyfield pipit	<i>Anthus rufulus</i>	C	RE	LC
59	Ploceidae	Baya Weaver	<i>Ploceus philippinus</i>	C	RE	LC
60	Dicruridae	Black drongo	<i>Dicrurus macrocerus</i>	A	RE	LC
61	Hirundinidae	Barn Swallow	<i>Hirundo rustica</i>	A	RE	LC
62		Wire-tailed Swallow	<i>Hirundo smithii</i>	C	RE	LC
63	Sturnidae	Bank myna	<i>Acridotheres ginginianus</i>	C	LM	LC
64		Jungle myna	<i>Acridotheres fuscus</i>	A	RE	LC
65		piebald myna	<i>Gracupica contra</i>	C	RE	LC
66		Brambling starling	<i>Sturnus roseus</i>	A	RE	LC
67		Common Myna	<i>Acridotheres tristis</i>	A	RE	LC
68	Nectarinidae	Purple Sunbird	<i>Cinnyris asiaticus</i>	C	RE	LC
69	Corvidae	House crow	<i>Corvus splendens</i>	C	RE	LC
70	Oriolidae	Black hooded oriole	<i>Oriolus xanthornus</i>	C	RE	LC

71	Muscicapidae	Oriental Magpie Robin	<i>Copsychus saularis</i>	C	RE	LC
72	Leiothrichidae	Jungle Babbler	<i>Turdoides striata</i>	A	RE	LC
73	Cisticolidae	Ashy Prinia	<i>Priniasocialis</i>	C	RE	LC
74		Zitting cisticola	<i>Cisticola juncidis</i>	C	RE	LC
75		Common Tailorbird	<i>Orthotomus sutorius</i>	A	RE	LC
76	Pycnonotidae	Red-vented Bulbul	<i>Pycnonotus cafer</i>	C	RE	LC
77	Laniidae	Brown shrike	<i>Lanius cristatus</i>	C	RE	LC
Order : Falconiformes						
78	Falconidae	common kestrel	<i>Falco tinnunculus</i>	R	LM	LC
Order : Accipitriformes						
79	Pandionidae	Osprey	<i>Pandion haliaetus</i>	R	LM	LC
80	Accipitridae	black-shouldered kite	<i>Elanus axillaris</i>	C	RE	LC
81		Black kite	<i>Milvus migrans</i>	C	RE	LC
Order : Bucerotiformes						
82	Upupidae	Common Hoopoe	<i>Upupa epops</i>	C	RE	LC
Order: Piciformes						
83	Megalaimidae	Copper smith Barbet	<i>Megalaima haemacephala</i>	C	RE	LC
84		Blue throated Barbet	<i>Psilopogon asiaticus</i>	C	RE	LC
Order : Cuculiformes						
85	Cuculidae	Asian Koel	<i>Eudynamis scolopacea</i>	A	RE	LC
86		Greater Coucal	<i>Centropus sinensis</i>	C	RE	LC
87		Indian Cuckoo	<i>Cuculus micropterus</i>	C	RE	LC
Order: Psittaciformes						
88	Psittaculidae	rose-ringed parakeet	<i>Psittacula krameri</i>	C	RE	LC
Order: Columbiformes						
89	Columbidae	Spotted dove	<i>Spilopelia chinensis</i>	C	RE	LC

Note: C = Common, A = Abundant, O = Occasional, R = Rare, RE- Resident, LM - Local Migrant, WM - Winter migrant, (IUCN) Categories: LC: Least Concern; VU: Vulnerable; NT : Near threatened

Maximum bird species were recorded during winter season due to presence of various types of migratory birds. The density of water bird was lowest during summer season followed by monsoon. A winter migratory bird generally arrives in the month of november and stayed up to March in every year. Overall Status of bird recorded at Purbasthali Oxbow Lake is given in Table 2. Among the winter migratory birds Lesser whistling duck, Cotton pygmy goose, Red crested pochard and Ferruginous duck are abundant in this lake. Birds were categorized as Common (C) 49 species, Abundant (A) 24 species, Rare (R) 10 species and Occasional (O) 6 species.




Figure 2: Residential Status of Birds in Purbasthali Oxbow Lake

Table 2: Status of bird recorded at Purbasthali Oxbow Lake

Sl.No	Order	No of Families	No of Species	Occurrence				Residential Status		
				C	A	O	R	RE	LM	WM
1	Charadriiformes	5	16	9	3	1	3	2	7	7
2	Suliformes	1	2	0	1	0	1	1	1	0
3	Ciconiiformes	4	6	5	1	0	0	6	0	0
4	Gruiformes	1	5	1	3	1	0	4	0	1
5	Anseriformes	1	15	6	4	2	3	0	0	15
6	Pelecaniformes	2	8	3	3	1	1	4	1	3
7	Podicipediformes	1	2	0	1	1	0	0	0	2
8	Passeriformes	13	23	16	7	0	0	20	3	0
9	Falconiformes	1	1	0	0	0	1	0	1	0
10	Accipitriformes	2	3	2	0	0	1	2	1	0
11	Bucerotiformes	1	1	1	0	0	0	1	0	0
12	Piciformes	1	2	2	0	0	0	2	0	0
13	Cuculiformes	1	3	2	1	0	0	3	0	0
14	Psittaciformes	1	1	1	0	0	0	1	0	0
15	Columbiformes	1	1	1	0	0	0	1	0	0
Total		36	89	49	24	6	10	47	14	28




Figure 3: Avian diversity of Purbasthali Oxbow Lake in terms of order

The lake thrives lots of residential bird species like Bronze winged jacana, Pheasant tailed jacana, Purple moorhen, Common Moorhen, Purple Heron, Indian Pond Heron etc. Local migratory species like Red-Wattled Lapwing, Yellow-Wattled Lapwing, Common redshank, Green sandpiper, common crestol were also very important to the lake ecosystem. Among the winter migratory birds, Lesser Whistling Duck were the most abundant species. Cotton pigmy goose, Ferruginous Pochard, Red crested Pochard was also found in good number in this area. The species like Knob-billed duck, Glossy Ibis, Great Crested Grebe, Black winged stilt were occasionally visited this area. Some species like Ruddy shelduck, Eurasian teal, little stint etc. are rarely found in the Purbasthali oxbow lake.

4. Conclusion

The present study reveals that the Purbasthali Oxbow Lake is an important habitat for various avifaunal species including winter migratory birds. Most of the avian species found during the winter season due to presence of twenty eight migratory birds species. Now-a-days the lake is undergoing unwanted change in biodiversity due to unplanned management strategies. Expansion of agricultural activities, use of pesticides, livestock grazing and use of submerges fishing net were the main threats to the survival of the birds in and around Purbasthali Oxbow Lake. Activities of the tourists are also causing disturbances to the lake ecosystem. There should be strict management rules for the tourists for the conservation of biodiversity in the lake area. The ecological health of the lake and habitat quality for birds can be improved by initiation of vegetation control program, regulation on the use of pesticides, pollution control strategies, strict check on land encroachment in this area, regulated tourism practices, scientific research and monitoring practices etc. A sustainable and holistic management planning is necessary for conservation of Purbasthali oxbow lake.

5. Acknowledgement

The author is thankfully acknowledged the financial support received from UGC as minor research project (sanction no PSW-199/13-14 ERO). Author is also grateful to Nabadwip Vidyasagar College for providing infrastructural facilities.

References

- [1] Amat J.A., Green A.J. (2010) Waterbirds as Bioindicators of Environmental Conditions. Conservation Monitoring in Freshwater Habitats, *Springer Netherlands*, 45-52
- [2] Anon (2000) Wetland values and functions. The Ramsar Bureau. Gland, Switzerland. 20-25.
- [3] Gregory, R. (2006) Birds as biodiversity indicators for Europe. *Significance* (3) 106-110
- [4] Jorvinen, O., Vaisenen, R.A. (1978) Recent changes in forest bird population in Northern Finland. *Anim. Zool.*(15) 279-289.
- [5] Bowden, C.G.R. (1990) Selection of foraging habitat by woodlarks nesting in pin plantation. *Applied Ecology*. (27) 410-419.
- [6] Rundle, W.D., (1982) A case for esophageal analysis in sho-rebird food studies. *Journal of Field Ornithology*, (55) 249-257.
- [7] Mcparland, C.E., Paszkowski, C.A. (2007) Water bird assemblages in the Aspen Parkland of western Canada: the influence of fishes, invertebrates, and the environment on species composition. *Ornithol. Sci.*(6) 53-65..
- [8] Schwartz, C.W., Schwartz, E.R. (1951) An ecological reconnaissance of the pheasants of Hawaii. *Auk*. (68) 281-314
- [9] Gupta. A.K, Singh.S.K, (2003) Changing wetlands due to discharge of effluents from small scale industries around varuna river corridor, *Ecol.Env. & Cons.* 9 (2) 209-212.

- [10] Rottenborn, S.C. (1999) Predicting the impacts of urbanization on riparian bird communities. *Biological Conservation*, **88**, 289-299.
- [11] Mckinney, M.L. (2002) Urbanization, biodiversity, Conservation. *Bioscience*, (52) 883-890.
- [12] Czech, H.A., K.C. Parsons, (2002) Agricultural wetlands and water birds: A Review. *Water birds* (25) 56-65.
- [13] R. Manakadan and A. Pittie, (2001) Standardized common and common names of the birds of the Indian sub continent, *Buceros*, 6 (1) 1-38.
- [14] Buckland, S.T., Marsden, S.J. & Green, R.E. (2008) Estimating bird abundance: Making methods work. *Bird Conservation International* 18 (S1) 91-108.
- [15] A. Kumar., J.P. Sati., P.C. Tak., J.R.B. Alfred. (2005) Handbook of Indian wetland birds and their conservation. *Director Zoological Survey India*, 167-170.
- [16] Ali, S. (2002) The Book of Indian Birds (13th Edition) Published by Bombay Natural History Society, Oxford University press.
- [17] Grimmett R, Inskipp C, Inskipp T. Helm field guides. (2011) birds of the Indian subcontinent (2nd edition). *Oxford University Press*,.
- [18] Basavarajappa S. (2006) Avifauna of Agro ecosystem of Maidan area of Karnataka. *Zoos' Print J.* 21 (4) 2217-2219.

