

# On the Non-Homogeneous Quintic with Five Unknowns $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^3$

Dr.P.Jayakumar<sup>1</sup>, R.Venkatraman<sup>2</sup>

<sup>1</sup>Professor of Mathematics, Annai Vailankanni Arts and Science College, Thanjavur-613007

<sup>2</sup>Assistant Professor, Department of Mathematics, SRM University, Vadapalani Campus, Chennai- 600026

**Abstract:** Five different methods of the non-zero integral solutions of the Non homogeneous Quintic Diophantine equation with five unknowns  $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^3$  are determined. Some interesting relations among the special numbers and the solutions are exposed.

**Keywords:** The non homogeneous Diophantine equation, Quintic equation with five unknowns. Integral solutions, special numbers, a few interesting relation, 2010 Mathematics subject Classification: 11D09

## Notations

- $T_{m,n}$  - Polygonal number of rank n with size m
- $P_n^m$  - Pyramidal number of rank n with size m
- $g_n$  - Gnomonic number of rank n
- $Pr_n$  - Pronic number of rank n
- $Ct_{16,n}$  - Centered hexadecagonal pyramidal number of rank n
- $OH_n$  - Octahedral number of rank n
- $SO_n$  - Stella octangular number of rank n
- $RD_n$  - Rhombic dodecagonal number of rank n
- $Hon$  - Haury Octahedral number of rank n

## 1. Introduction

The number theory is the Queen of Mathematics. In particular, the Diophantine equations have a blend of attracted interesting problems. For an extensive review of variety of problems, one may refer to [1,9] the problem of finding all integer solutions of a Diophantine equation with three or more Variables and degree at least three, in general presents a good deal of difficulties. There is vast general theory of homogeneous quadratic equations with three variables [1-4,9]. Cubic equations in two variables fall in to the theory of elliptic curves which is a very developed theory but still an important topic of current research [5-7]. A lot is known about equations in two variables in higher degrees. For equations with more than three variables and degree at least three very little is known. It is worth to note that undesirability appears in equations, even perhaps at degree four with fairly small coefficients. In [1,8 and 9] a few higher order equations are considered for integral solutions. In this communication a seventh degree non-homogeneous equation with five variables represented by  $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^3$  is considered and in particular a few interesting relations among the solutions are presented.

## 2. Description of Method

Consider the Quintic Diophantine equation

$$(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^3 \quad (1)$$

Introduce of the linear transformation

$$x = u + v, y = u - v, X = 2u + v, Y = 2u - v \quad (2)$$

In (1) leads to

$$u^2 + 17v^2 = 21z^3 \quad (3)$$

Now we solve (3) through various choices and the different methods of solutions of (1) are obtained as follows.

### 2.1 Method: I

Assume

$$z = z(a, b) = a^2 + 17b^2 \quad (4)$$

Where a & b are non zero distinct integers

Write 13 as

$$21 = (2 + i\sqrt{17})(2 - i\sqrt{17}) \quad (5)$$

Using, (4) and (5) in (3) and applying the method of factorization. Define

$$(u + i\sqrt{17}v) = (2 + i\sqrt{17})(a + i\sqrt{17}b)^2$$

Equating real and imaginary parts, we get

$$u = u(a, b) = 2a^3 - 102ab^2 - 51a^2b + 289b^3$$

$$v = v(a, b) = a^3 - 51ab^2 + 6a^2b - 34b^3$$

Hence in view of (2) the corresponding solutions of (1) are given by

$$x = x(a, b) = 3a^3 - 153ab^2 - 45a^2b + 255b^3$$

$$y = y(a, b) = a^3 - 51ab^2 - 57a^2b + 323b^3$$

$$X = X(a, b) = 5a^3 - 255ab^2 - 96a^2b + 544b^3$$

$$Y = Y(a, b) = 3a^3 - 153ab^2 - 108a^2b + 612b^3$$

$$z = z(a, b) = a^2 + 17b^2$$

### Observations:-

1.  $3X(a,1) - 5Y(a,1) - 252T_{4,a} \equiv 7 \pmod{204}$
2.  $x(a,1) - 3y(a,1) - 126T_{4,a} \equiv 0 \pmod{2}$
3.  $3z(3a,3a)$  a Nasty number
4.  $x(a+1,1) + y(a+1,1) - RD_a - 84T_{4,a} - G_{200a} \equiv 0 \pmod{2}$
5.  $3X[a(a+1),1] - 5Y[a(a+1),1] - 252(P_a)^2 \equiv 0 \pmod{2}$
6.  $2[x(a+1,1) - y(a+1,1)] - 3HO_a - 42T_{4,a} + G_{76a} \equiv 0 \pmod{2}$

### 2.2 Method: II

Instead of (5) Write 21 as

$$21 = \frac{1}{25} (10 + i5\sqrt{17})(10 - i5\sqrt{17}) \quad (6)$$

Following the procedure similar to Method-I, the corresponding non-zero distinct integral solutions of (1) are found to be

$$\begin{aligned} x &= x(a,b) = 375a^3 - 19125ab^2 - 5625a^2b + 31875b^3 \\ y &= y(a,b) = 125a^3 - 6375ab^2 - 7125a^2b + 40375b^3 \\ X &= X(a,b) = 875a^3 - 44625ab^2 - 18375a^2b + 104125b^3 \\ Y &= Y(a,b) = 625a^3 - 31875ab^2 - 4125a^2b + 23375b^3 \\ z &= z(a,b) = 25a^2 + 425b^2 \end{aligned}$$

### Observations:-

1.  $\frac{1}{10}z(2a,2a)$  is a Nasty number.
2.  $x(a,1) - 3y(a,1) - 15750T_{4,a} \equiv 0 \pmod{2}$
3.  $625X[a(2a^2-1),1] - 875Y[a(2a^2-1),1] + 7875000[SO_a]^2 \equiv 0 \pmod{2}$
4.  $z[a+1,a+1] - 450T_{4,a} - G_{45a} \equiv 11 \pmod{41}$
5.  $625X[a(a+1),1] - 875Y[a(a+1),1] + (P_a)^2 \equiv 7 \pmod{408}$
6.  $x[(2a-1),1] - 3y[(2a-1),1] - 15750(G_a)^2 \equiv 0 \pmod{2}$

### 2.3 Method: III

Rewrite (3) as  $u^2 + 17v^2 = 21z^3 * 1$

Write 1 as

$$1 = \frac{1}{81} (8 + i\sqrt{17})(8 - i\sqrt{17})$$

Following the procedure similar to Method-I, the corresponding non-zero distinct integral solutions of (1) are found to be

$$\begin{aligned} x &= x(a,b) = 15309a^3 - 780759ab^2 - 45927a^2b + 260253b^3 \\ y &= y(a,b) = 11907a^3 - 607257ab^2 - 127575a^2b + 722925b^3 \\ X &= X(a,b) = 28917a^3 - 1474767ab^2 - 132678a^2b + 751842b^3 \\ Y &= Y(a,b) = 25515a^3 - 1301265ab^2 - 214326a^2b + 1214514b^3 \\ z &= z(a,b) = 81a^2 + 1377b^2 \end{aligned}$$

### Observations:-

1.  $\frac{1}{72}Z(2a,2a)$  is a Perfect Square
2.  $\frac{1}{18}Z(1,1)$  is a Cubic integer
3.  $\frac{1}{183709}[x(1,1) - y(1,1)]$  is a Cubic integer

### 2.4 Method: IV

Instead of (5) Write 21 as

$$21 = \frac{1}{49} (14 + i7\sqrt{17})(14 - i7\sqrt{17})$$

Following the procedure similar to Method-I, the corresponding non-zero distinct integral solutions of (1) are found to be

$$\begin{aligned} x &= x(a,b) = 1029a^3 - 52479ab^2 - 15435a^2b + 87465b^3 \\ y &= y(a,b) = 343a^3 - 17493ab^2 - 19551a^2b + 110789b^3 \\ X &= X(a,b) = 1715a^3 - 87465ab^2 - 32928a^2b + 18659b^3 \\ Y &= Y(a,b) = 1029a^3 - 52479ab^2 - 37044a^2b + 209916b^3 \\ z &= z(a,b) = 25a^2 + 425b^2 \end{aligned}$$

### Observations:

1.  $343x(a,1) - 1029y(a,1) - 14823774T_{4,a} \equiv 17 \pmod{4941258}$
2.  $\frac{1}{5713}[X(1,1) + 2Y(1,1)]$  is a Perfect Square
3.  $1029X[a(a+1),1] - 1715Y[a(a+1),1] - 29647548(P_a)^2 \equiv 349 \pmod{976521}$
4.  $\frac{1}{4116}[x(1,1) + y(1,1)]$  is a woodall number.
5.  $\frac{1}{162}z[9a,9a]$  is a perfect square

### 2.5 Method: V

Instead of (5) Write 21 as

$$21 = \frac{1}{64} (16 + i8\sqrt{17})(16 - i8\sqrt{17})$$

Following the procedure similar to Method-I, the corresponding non-zero distinct integral solutions of (1) are found to be

$$\begin{aligned} x &= x(a,b) = 1536a^3 - 78336ab^2 - 23040a^2b + 130560b^3 \\ y &= y(a,b) = 512a^3 - 26112ab^2 - 29184a^2b + 165376b^3 \\ X &= X(a,b) = 2560a^3 - 130560ab^2 - 49152a^2b + 278528b^3 \\ Y &= Y(a,b) = 1536a^3 - 78336ab^2 - 55296a^2b + 313344b^3 \end{aligned}$$

### Observations:

1.  $\frac{1}{524288}[512x(a,1) - 1536y(a,1) - 33030144T_{4,a}] - 21$  is a nasty number
2.  $1536X[a(a+1),1] - 2560Y[a(a+1),1] - 66060288(P_a)^2 \equiv 0 \pmod{2}$
3.  $\frac{1}{18}z[2a,2a]$  is a Perfect square
4.  $1536X(a,1) - 2560Y(a,1) \equiv 0 \pmod{2}$

## 3. Conclusion

In linear transformation (2), the variables X and Y may also be represented by  $X=2uv+1$ ,  $Y=2uv-1$ . Applying the procedure similar to that of Method I-V choices of integral solutions to (1) are obtained. To conclude, one may search

for other method of solutions and their corresponding properties.

## References

- [1] Dickson, L.E., History of theory of numbers, Vol.11, Chelsea publishing company, New –York (1952)
- [2] Mordell, L.J., Diophantine equation, Academic press, London (1969) Journal of Science and Research, Vol (3) Issue 12, 20-22 (December 14)
- [3] Jayakumar. P, Sangeetha, K “Lattice points on the cone  $X^2 + Y^2 = 50Z^2$  ” International Journal of science and Research, Vol(3), Issue 12, 20-22 (December -2014 )
- [4] Jayakumar P, Kanaga Dhurga. C. “On Quadratic Diophantine equation  $X^2 + 16Y^2 = 20Z^2$  ” Galois J. Maths, 1(1) (2014), 17-23.
- [5] Jayakumar P, Kanaga Dhurga. C. “Lattice points on the cone  $X^2 + 9Y^2 = 50Z^2$  ” Diophantus J. Math3(2) 2014.
- [6] Jayakumar. P, Prabha. S “On Ternary Quadratic Diophantine equation  $X^2 + 15Y^2 = 14Z^2$ ” Archimedes J. Math., 4(3) (2014), 159-164
- [7] Jayakumar, P, Meena, J “Integral solutions of the Ternary Quadratic Diophantine equation :  $X^2 + 7Y^2 = 16Z^2$  ” International Journal of Science and Technology, Vol.4, Issue 4, 1-4, Dec 2014.
- [8] Jayakumar. P, Shankarakalidoss, G “Lattice points on Homogenous cone  $X^2 + 9Y^2 = 50Z^2$  ” International journal of Science and Research, Vol (4), Issue 1, 2053-2055, January -2015
- [9] Jayakumar. P, Shankarakalidoss. G “Integral points on the Homogenous cone  $X^2 + Y^2 = 10Z^2$ ” International Journal for Scientific Research and Development, Vol (2), Issue 11, 234-235, January -2015
- [10] Jayakumar.P & Sangeetha.k “On the Non Homogeneous Heptic Equation With Five Unknowns  $(X^2 - Y^2)(9X^2 + 9Y^2 - 16XY) = 15Y^2 = 21(X^2 - Y^2)Z^5$  ” International journal of Science and Research volume 3 issue 11, November 2014
- [11] Jayakumar.P & Venkatraman.R “On The Homogeneous Biquadratic Equation With 5 Unknowns:  $X^4 - Y^4 = 65(Z^2 - W^2)R^2$  ” International Journal of Science and Research (IJSR) Volume 5 Issue 3, March 2016.
- [12] Jayakumar.P & Venkatraman.R “Lattice Points on the Homogeneous Cone  $X^2 + Y^2 = 26Z^2$  ” International Journal of Science and Research (IJSR) Volume 5 Issue 3, March 2016.
- [13] Jayakumar.P, Pandian.V & Venkatraman.R “On Non-Homogeneous Bi-Quadratic Diophantine Equation  $4(X^2 + Y^2) - 7XY = 19Z^4$  ” International Journal of Science and Research (IJSR), Volume 5 Issue 4, April 2016.
- [14] Jayakumar.P, Pandian.V & Venkatraman.R “On Non-Homogeneous Cubic Diophantine Equation  $4X^2 + 4Y^2 - 7XY = 19Z^3$  ” International Journal

of Science and Research (IJSR) , Volume 5 Issue 4, April 2016.

- [15] Jayakumar.P & Venkatraman.R “On Non-Homogeneous Biquadratic Diophantine Equation  $7(X^2 + Y^2) - 13XY = 31Z^4$  ” International Journal of Science and Research (IJSR), Volume 5 Issue 11, November 2016.

## Author Profile

**Dr. P. Jayakumar** received the B. Sc, M.Sc degrees in Mathematics from Madras University in 1980 and 1983 and the M. Phil, Ph.D degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 1988 and 2010. Who is now working as Professor of Mathematics, Annai velankanni Arts and Science college, Thanajvur-613 007, Tamil Nadu, India.

**R. Venkatraman** received the B.Sc, M.Sc, and MPhil degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 2002, 2004 and 2006. Who is now working as Assistant Professor of Mathematics, SRM University Vadapalani Campus, Chennai – 600026, Tamil Nadu, India.