Criteria for Success in Dental Implants: A Systematic Review

Nurul Syamimi binti Mohd Azlan Sunil¹, Dr. Ashok², Dr. Dhanraj³

¹Bachelor of Dental Surgery (Year 2), Saveetha Dental College and Hospitals, No. 162, PH Road, Chennai-600077

²Saveetha Dental College and Hospitals, No.162, PH road, Chennai-600077

³Saveetha Dental College and Hospitals, No.162, PH road, Chennai-600077

Abstract: <u>Aim and Objective</u>: The purpose of this study was to examine the most frequently used criteria to define treatment success in implant dentistry. <u>Materials and Method</u>: An electronic search in the MEDLINE/PubMED database was performed for studies published in English from January 1980 until October 2010. The search strategy included the following key word combinations: 'success criteria AND 'implant', 'success rates' AND 'implant', 'survival rates' AND 'implant', and 'outcomes' AND 'implant dentistry'. <u>Results</u>: The electronic and manual search yielded 2 randomized clinical trials and 15 prospective. In total, 17 publications were suitable for analysis. Some of the included articles had mixed data reporting on more than one type of prosthesis. Most of the articles included four most used parameters which are taken into consideration are the implant fixture, the peri-implant soft tissue, prosthetic level and also patient's satisfaction. <u>Conclusion</u>: Each criteria has its own consideration to be evaluated in order to determine success. Pain, mobility and bone loss are the most commonly used criteria to report success at the implant level. Suppuration and probing depth are the frequently used criteria at peri-implant soft tissue level. As for prosthetic level, the commonly used criteria include function and esthetics while patient's satisfaction is commonly considered a success if comfort and appearance satisfaction are achieved. The reported success rate consistently decreased when the number of parameters included for the assessment of success was increased.

Keywords: implant success criteria, implant survival rates, implant success rate, implant dentistry outcome, implant dentistry

1. Introduction

Research and technology of implants has made a revolutionary change during the last two decades in thereplacement of missing teeth with endosseous implants the standard care and an implant supported prosthesis as the first line of treatment and long lasting rehabilitation. Over the pass thirty years, success and survival in implant dentistry has been evaluated by the survival rate, radiographic crestal bone loss, prosthesis stability and also the presence of peri-implant diseases.[1-6] Albrektsson and colleagues[1] proposed criteria for the assessment of implant success, to determine implant survival and clinical evidence of successful osseointegration, which is the most commonly accepted criteria.

There are many new parameters that have been introduced to assess success in implant restorations, which includes health status and natural-looking peri-implant soft tissues, aesthetics and patient fulfilment. However, osseointegration remains as an important parameter in implant dentistry. It appears to be coherent that the current meaning of success criteria ought to be comprehensive, to incorporate these additional factors.[6-10]

This systematic review is done to find out the most frequently used criteria to determine success of treatment in implant dentistry.

2. Materials and Methods

The search strategy involves the MEDLINE/PubMED database electronic search. The key word which was searched include: *'implant success criteria', 'implant*

survival rates', and *'implant dentistry outcome'*. The search for studies in English which were published from January 1980 until October 2010.

This systematic review focuses on specific studies of success and survival rate of implants, therefore, a few inclusion criteria were used to conduct the studies selection. The inclusion criteria include studies with five-year follow up (minimum), studies which involves at least 10 patients, studies reporting on success or survival criteria used to assess the outcomes of a treatment. The success rate of implant of the selected articles are the primary outcome variable. The studies were grouped according to four differentiated clinical situations which are implant fixed complete dental prostheses (FCDP), implant overdentures (IOD), fixed partial dentures (FPD), and single crowns (SC). This is done so to ensure the homogeneity of the result. The parameters used to define success are the secondary outcome variables: implant fixtures, peri-implant soft tissue, prosthesis, and patient's evaluation.

3. Results

Identification of Studies

The initial search according to the key words yielded 231 citations. Out of the 231 studies, 214 studies were excluded as the studies did not meet the inclusion criteria. A majority of the studies did not have either a minimum follow-up period of 5 years, less than 10 patients in the study and/or did not have a detailed report on the survival and success rate of implant and the treatment outcome. After two rounds of inclusion and exclusion, a total of 17 studies were eligible for analysis.

Volume 6 Issue 4, April 2017 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

Included studies

A total of 17 articles satisfy the inclusion criteria and were suitable for analysis. The included studies are of 2 randomized clinical trials and 15 perspective studies.Some of the studies reported more than one type of prosthesis.

Clinical situation	Study	Patients	Follow up (years)	Criteria	Success/Survival Rate	
Implant Fixed Complete	Degidi et al., 2009 [11]	38	5	Based on Albrektsson et al., 1986	100% for implants; 100 % for prostheses	
Dental Prostheses	Fischer <i>et al.</i> , 2008 [12]	24	5	Self-defined criteria	95.2% for implants; 100% for prostheses	
(FCDP)	Rasmussan et. al., 2004[13]	36	10	Self-defined criteria	96.9% for implants; 100% for prostheses	
	Astrand et al., 2004 [14]	66	5	Based on Albrektsson et al., 1986	96.30%	
	Buser et al., 1997[15]	1003	Up to 8 yrs	Based on Buser et al., 1990	97.3% for 5 yrs; 93.3 % for 8 yrs	
	Romeo et al., 2004[16]	255	1.3~7 (Mean: 3.85)	Based on Albrektsson <i>et al.</i> , 1986 & Buser et al., 1990	63.8 % for prostheses	
Implant Overdentures	Ma et al., 2010[17]	79	10	Based on Albrektsson et al., 1986	85.9% for 5 yrs; 74.5 % for 10 yrs	
(IO)	Al Fadda et al., 2009[18]	77	5	Based on Albrektsson & Zarb, 1998	98%	
	Zinsli et al., 2004[19]	149	10	Based on Buser et al., 1990	98.70%	
	Buser et al., 1997	1003	Up to 8 yrs	Based on Buser et al., 1990	97.3% for 5 yrs; 93.3 % for 8 yrs	
	Romeo et al., 2004	255	1.3~7 (Mean: 3.85)	Based on Albrektsson <i>et al.</i> , 1986 & Buser et al., 1990	63.8 % for prostheses	
Implant Fixed Partial Denture	Payer et al., 2010[20]	24	5	Self-defined criteria	92.50%	
(FPD)	Degidi et al., 2009	72	5	Based on Albrektsson et al., 1986	99.4% for implants, 100% for prostheses	
	Glauser et al., 2007[21]	38	5	Based on Albrektsson et al., 1986	97.10%	
	Bornstein et al., 2005[22]	51	5	Based on Buser et al., 1990	99%	
	Zinsli et al., 2004	149	10	Based on Buser et al., 1990	98.70%	
	Romeo et al., 2004	255	1.3~7 (Mean: 3.85)	Self-defined criteria	63.8 % for prostheses	
	Buser et al., 1997	1003	Up to 8 yrs tients	Based on Buser et al., 1990	97.3% for 5 yrs; 93.3 % for 8 yrs	
	Weber <i>et al.</i> , 2000[23]	46	5	Self-defined criteria	99.10%	
Implant Single	Payer et al., 2010	24	5	Self-defined criteria	92.50%	
Crown (SC)	Degidi et al., 2009	38	5	Based on Albrektsson et al., 1986	100% for implants; 100% for prostheses	
	Glauser et al., 2007	38	5	Based on Albrektsson et al., 1986	97.10%	
	Bornstein et al., 2005	51	5	Based on Buser et al., 1990	99%	
	Romeo et al., 2004	255	1.3~7 (Mean: 3.85)	Based on Albrektsson <i>et al.</i> , 1986 & Buser et al., 1990	75.6 % for prostheses	
	Zinsli et al., 2004	149	10	Based on Buser et al., 1990	98.70%	
	Buser et al., 1997	1003	Up to 8 yrs tients	Based on Buser et al., 1990	97.3% for 5 yrs; 93.3 % for 8 yrs	
	Weber et al., 2000	46	5	Self-defined criteria	99.10%	
	Roccuzzo et al., 2008[24]	27	5	Self-defined criteria	100%	
	Wennstrom et al., 2005[25]	36	5	Self-defined criteria	97.70%	

Six articles reported on Implant Fixed Complete Dental Prosthesis, five articles were on Implant Over Dentures, eight articles reported on Implant Fixed Partial Denture and Single Crown was reported by ten articles.

To determine the success of implant, most of the articles included four most used parameters which are taken into consideration are the implant fixture, the peri-implant soft tissue, prosthetic level and also patient's satisfaction.Each criteria has its own consideration to be evaluated in order to determine success. Pain, mobility and bone loss are the most commonly used criteria to report success at the implant level. Suppuration and probing depth are the frequently used criteria at peri-implant soft tissue level. As for prosthetic level, the commonly used criteria include function and esthetics while patient's satisfaction is commonly considered a success if comfort and appearance satisfaction are achieved.

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

Success Criteria		FCDP	IOD	FPD	SC
		(6 articles)	(5 articles)	(8 articles)	(10 articles)
Implant level	Pain	4	5	5	7
	Annual bone $loss < 0.2$ mm thereafter	2	3	4	3
	Radiolucency	3	3	5	7
	Mobility	5	5	6	8
Peri-implant soft tissue	Probing depth $> 3 \text{ mm}$	2	1	2	2
	Suppuration	3	3	5	6
	Bleeding	2	0	0	0
	Swelling	1	0	0	0
	Recession	1	0	0	0
Prosthetic level	Minor complications (chairside approach)	2	0	0	0
	Major complications/failures	2	0	1	0
	Esthetics	1	0	0	0
	Functional	1	2	3	3
Patient satisfaction	Discomfort/paresthesia	4	4	1	4
	Satisfaction with appearance	1	0	4	1
	Ability to chew	1	0	0	0
	Ability to taste	1	0	0	0

Table 2: Reported Success Criteria as Described in All Selected Articles

Table 3 shows the range of success rate of implant for each of the number of parameter included. As the number of parameter adds up, the range of success rate decreases.

Table 3: Number of Included Parameters and Success Rat
--

No. of Parameters	FCPD (6 articles)		FPD (8 articles)		IOD (5 articles)		SC (10 articles)	
	Reported	Range of	Reported	Range of	Reported	Range of	Reported	Range of
	Articles	Success Rate	Articles	Success Rate	Articles	Success Rate	Articles	Success Rate
1	6 (100%)	95.2%-100%	8 (100%)	92.5%-100%	5 (100%)	63.8%-98.7%	10 (100%)	75.6%-100 %
2	-	-	2 (25%)	98.7%-99.1%	1 (20%)	74.5%-85.9%	4 (40%)	97.7 %-99.1%
3	4 (66.7%)	92.7%-100%	4 (50%)	93.3%-100%	2 (40%)	93.3%-98%	4 (40%)	97.1 %-100%
4	2 (33.3%)	63.8%	1 (12.5%)	63.8%	1 (20%)	63.80%	1 (10%)	75.60%

4. Discussion

As shown in Table 3, there is a constant decrease in the rate of success when there is an increase in the number of parameters included. This indicated that as the higher the number of parameter are considered in an implant, the chances of a successful implant thins down as many aspects need to be taken into consideration and evaluated to measure the outcome.The four most used parameters in this systemic review included the implant level, peri-implant soft-tissue level, prosthesis level, and the patient's subjective assessment. Besides that, aesthetics is also as important in dentistry and sets a benchmark for the inclusion of other various factors to measure the success of implant prostheses.

The success or failure of dental implants is influenced by many factors. Well-defined criteria that report and evaluate outcomes and also complications should ideally be used.[26]Many modifications have been made throughout the years that involves other parameters, for example; aesthetics and patient-centered outcome as a criteria to measure success on dental implants.

References

Albrektsson, T., et al., *The long-term efficacy of currently used dental implants: a review and proposed criteria of success.* Int J Oral Maxillofac Implants, 1986. 1(1): p. 11-25.

- [2] Smith, D.E. and G.A. Zarb, *Criteria for success of osseointegrated endosseous implants*. The Journal of prosthetic dentistry, 1989. 62(5): p. 567-572.
- [3] Buser, D., H.P. Weber, and N. Lang, *Tissue integration* of non-submerged implants. *l-year results of a* prospective study with 100 ITI hollow-cylinder and hollow-screw implants. Clinical oral implants research, 1990. **1**(1): p. 33-40.
- [4] Albrektsson, T. and G. Zarb, *Determinants of correct clinical reporting*. The International journal of prosthodontics, 1998. **11**(5): p. 517.
- [5] Misch, C.E., et al., Implant success, survival, and failure: the International Congress of Oral Implantologists (ICOI) pisa consensus conference. Implant dentistry, 2008. 17(1): p. 5-15.
- [6] Annibali, S., et al., Usefulness of the aesthetic result as a success criterion for implant therapy: a review. Clinical implant dentistry and related research, 2012. 14(1): p. 3-40.
- [7] Fürhauser, R., et al., Evaluation of soft tissue around single-tooth implant crowns: the pink esthetic score. Clinical oral implants research, 2005. 16(6): p. 639-644.
- [8] Meijer, H.J., et al., A new index for rating aesthetics of implant-supported single crowns and adjacent soft tissues-the Implant Crown Aesthetic Index. Clinical oral implants research, 2005. 16(6): p. 645-649.
- [9] Belser, U.C., et al., Outcome evaluation of early placed maxillary anterior single-tooth implants using objective esthetic criteria: a cross-sectional, retrospective study in 45 patients with a 2-to 4-year follow-up using pink

Volume 6 Issue 4, April 2017

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

and white esthetic scores. Journal of periodontology, 2009. **80**(1): p. 140-151.

- [10] Pjetursson, B.E., et al., A systematic review of the survival and complication rates of all-ceramic and metal-ceramic reconstructions after an observation period of at least 3 years. Part I: single crowns. Clinical Oral Implants Research, 2007. 18(s3): p. 73-85.
- [11] Degidi, M., et al., Comparative Analysis of Immediate Functional Loading and Immediate Nonfunctional Loading to Traditional Healing Periods: A 5-Year Follow-Up of 550 Dental Implants. Clinical implant dentistry and related research, 2009. 11(4): p. 257-266.
- [12] Fischer, K., et al., Five-year results from a randomized, controlled trial on early and delayed loading of implants supporting full-arch prosthesis in the edentulous maxilla. Clinical oral implants research, 2008. 19(5): p. 433-441.
- [13] Rasmusson, L., J. Roos, and H. Bystedt, A 10-Year Follow-Up Study of Titanium Dioxide–Blasted Implants. Clinical implant dentistry and related research, 2005. 7(1): p. 36-42.
- [14] Åstrand, P., et al., Astra Tech and Brånemark system implants: a 5-year prospective study of marginal bone reactions. Clinical oral implants research, 2004. 15(4): p. 413-420.
- [15] Buser, D., et al., Long-term evaluation of non-submerged ITI implants. Part 1: 8-year life table analysis of a prospective multi-center study with 2359 implants. Clinical oral implants research, 1997. 8(3): p. 161-172.
- [16] Romeo, E., et al., Long-term survival and success of oral implants in the treatment of full and partial arches: a 7-year prospective study with the ITI dental implant system. International Journal of Oral and Maxillofacial Implants, 2004. 19(2): p. 247-259.
- [17] Ma, S., et al., Marginal bone loss with mandibular twoimplant overdentures using different loading protocols and attachment systems: 10-year outcomes. International Journal of Prosthodontics, 2010. 23(4).
- [18] Alfadda, S.A., N.J. Attard, and L.A. David, *Five-year clinical results of immediately loaded dental implants using mandibular overdentures*. International Journal of Prosthodontics, 2009. 22(4).
- [19] Zinsli, B., et al., Clinical evaluation of small-diameter ITI implants: a prospective study. International Journal of Oral and Maxillofacial Implants, 2004. 19(1): p. 92-99.
- [20] Payer, M., et al., Immediate provisional restoration of screw-type implants in the posterior mandible: results after 5 years of clinical function. Clinical oral implants research, 2010. 21(8): p. 815-821.
- [21] Glauser, R., et al., Five-year results of implants with an oxidized surface placed predominantly in soft quality bone and subjected to immediate occlusal loading. The Journal of prosthetic dentistry, 2007. 97(6): p. S59-S68.
- [22] Bornstein, M.M., et al., Early loading of non-submerged titanium implants with a sandblasted and acid-etched surface. Clinical oral implants research, 2005. 16(6): p. 631-638.
- [23] Weber, H.P., C.C. Crohin, and J.P. Fiorellini, A 5-year prospective clinical and radiographic study of non-submerged dental implants. Clinical oral implants research, 2000. 11(2): p. 144-153.

- [24] Roccuzzo, M., et al., Early loading of sandblasted and acid-etched implants: a randomized-controlled double-blind split-mouth study. Five-year results. Clinical oral implants research, 2008. 19(2): p. 148-152.
- [25] Wennström, J.L., et al., *Implant-supported single-tooth restorations: a 5-year prospective study*. Journal of clinical periodontology, 2005. **32**(6): p. 567-574.
- [26] Eckert, S.E., et al., Comparison of dental implant systems: quality of clinical evidence and prediction of 5-year survival. International Journal of Oral & Maxillofacial Implants, 2005. 20(3).