Geomorphology and Soil Properties at the Middle and Lower Eastern Clay Plains - Nuba Mountains. South Kordofan, Sudan

El-Abbas Doka M. Ali¹, Omima Omer A. Showgi², Hashim Ali Dawoud³

¹Land Evaluation and Remote Sensing, College of Agricultural Studies, Sudan University of Science and Technology, Shambat

²Properties and Genesis of Soils, College of Science and Technology, Department of Botany, Shendi University, Nile State

³Soil Pedology and land Use Planning, Department of Horticulture Science, College of Agriculture, Al Azhari University, Khartoum North, Sudan

Abstract: This research study aimed to describe the geomorphology and characterize the soil properties at the middle and lower clay plains located at the eastern parts of Nuba Mountains, Sudan. Soil samples were collected from nineprofile pits alonga transect stretching from Abbasyia to Megenis through Terter (64 Km). The area is a broad, slightly sloping to undulating upland clay plains. The plains are dissected by Khors and Wadis (Khor and Wadi are water course in Arabic language) draining the Nuba Mountains hilly area, numoures low ridges (viens and dykes) separated by shallow depressions running through the area. The majority of the soils identified belong to the dark cracking clays. The soils are formed <u>insitu</u> and on colluviums and alluvium derived from basement complex rocks. The origin of these clays is thought to belong to Quarternary alluvial deposits probably similar in time to other dark cracking clays of central Sudan but under different forming processes (Showgi et al 2016). These deposits form well developed Vertisols (Pacheco and Dawoud, 1976; Ahmed 1983; Abedine and Robibson 1970). Wide cracking, gilgae microrelief and very dark grayish colours dominate the surfaces. Topsoil structure has loose granular mulch, the subsoil has subangular blocky structure, below that the structure is massive. Subsoil and substrata layers have parallelepiped and wedge shaped structure aggregates. Slickensides was also observed in all profiles, reflecting expansion and contraction upon wetting and drying of the soil. Fine and medium tubular pores, small and large whitish irregular CaCO₃ concretions were noted. Fine and medium sand grains and polished quartz pebble arecommon. The soil reaction is slightly alkaline in all profiles increasing with depth. The sois are non saline and nonesodicon middle plains but mildly sodic and sodic at lower plains. Cation exchange capacity is high at top horizons and very high below indicating presence of illiticand montmorillonitic clay minerals. Sodium Adsorption Ratio (SAR) is lowand the soils are non - calcareous with low amount of $CaCO_3Organic$ matter is low decreasing with depth and the total nitrogen is also low. Nutrient status expressed as extractable cations meq/100 g soil is generally high: Soil texture is clay throughout the profile and clay content increases with depth. The bulk density (BD) is high increasing with depth, and porosity results coincidewith bulk density trend. Although most of the cracking clay soils in Sudan are sharing similar origin (basaltic igneous materials) but their subsequent depositional processes, landscape geomorphology and local climate have affected their mineralogical, physical and chemical properties. This applies to the clay plains of Nuba Mountains as most soils are characterized by lower CEC, mild alkalinity and low calcium carbonate content.

Keywords: Nuba Mountains, Middle and Lower Clay Plains, Soil Properties

1. Introduction

In the Sudan, Vertisols occur on large tract of land, totaling perhaps 50 million ha in area, is divided naturally into four separated areas: the central clay plain (Gezira), the eastern clay plain (El Gadarif), the Nuba Mountains region and the southern clay plain (Abyei-Bahr Al Arab). Vertisols in the Nuba Mountains region occur on gently undulating plains and have better surface drained than those in extremely flat landscape of the other three clay plains (Blockhuis et al 1964; Eswaran et al 1999; FAO 1970). Dark cracking clay soils "Vertisols" cover not only large areas of the low lying plains but also they occur on undulating higher plains and in many intermountain valleys throughout the Nuba Mountains area(Vail, J.R. 1973). Some of these clays, notably those in eastern and southern lower clay plains, Khor Abu Habil, Wadi Al Ghalla and the clays of Bahr el Arab valley, can only be alluvial. But some other clay, for instance, on undulating high plains in the Nuba Mountains uplands area indicate a non-alluvial origin(Showgi 2011; Doka et al 2016).

The Nuba Mountains Vertisols are considered to be derived from the underlying bedrock. Therefore, it is expected that the parent materials at upper clay plains have undergone through a shorter pedogenetic processes compared to those on low alluvial plains (Khoddarry 1978). These non-alluvial clays are thought to be moved slowly down slope by combined process of wash and mass movement. Bases are thought to be washed from higher areas to accumulate on the plains where they help in the formation of the higher base status clays. Because the lower horizons are subject to pressure when the soils swell, they are compact and very slowly permeable to water. They are usually neutral to slightly in reaction (Pacheco and Dawoud1976). This research study attempts to identify the chemical and physical properties of the Vertisols at Middle and Lower clay plains at Nuba Mountains to provide information about soil inherent and dynamic properties. Many research studies were carried to characterize the chemical and physical properties of Vertisols at different parts the alluvial lowland clay plains of Sudan which are mainly used for irrigated farming (Blockhuis et al 1964; Kevie and Buraymah; 1987 Khalil 1986). The clay soils at the rainfed lands found less detailed research attention and particularly the upper, middle and

lower clay plains at Nuba Mountains region which need more understanding of their formation and behavior as related to their use and management.

2. The Study Area

The study area is located in the eastern part of the Nuba Mountains in South Kordofan State. The sampling transect (65 Km), starts from Al Abbasyia town (height 561 m ASL) in the northeast and ending towards southeast at Megenis (height 406 m ASL), Figure1 and Figure 2. The study area has tropical semi – arid zone of summer rain (Van der Kevie 1973). The rainfall is comparatively high as the survey area lies within isohyets 600 - 800 mm. The amount of rainfall increases from northeast to southwest (Table 1). The rainy season starts from late June and extends to early October month. Maximum temperatures range between 31C° and 40 C° with March to May as the hottest months and monthly minimum temperatures range between 17 C° and 24 (Abbasyia) C° coldest months being from early December to the end February (ELTom 1972 and Dawoud 1974).



Figure 1: The Location of the Study Area at the Eastern Clay Plains of Nuba Mountains - South Kordofan State

Table 1: Rainfall at the Study Area

Season	Al Abbasyia	AL Terter	Abu Jubayhah	Season	Al Abbasyia	AL Terter	Abu Jubayhah
1997 – 98	482.4	667.4	573.5	2002 - 03	510	509	526
1998 – 99	668	-	472	2003 - 04	554.5	499.6	812
1999 - 00	555.6	506	568	2004 - 05	532.5	414	661.5
2000 -01	477	483	614	2005 - 06	555.7	427	662
2001 - 02	589	711	638	2006-07	723.3	749.4	671.5

Source: Sudan Meteorological Authority

3. Materials and Methods

Transect stretches in eastward direction from Al Abbasyiato Megenis. The area under study is situated between longitudes 31°00" and 32°35" East and latitudes 11°56" and 12°09" north. Eight soil samples were collected from eight sites shown in Table 2 and represented in Figure 2. Subscenes of Landsat TM Satellite image 2010 covering the study area were used for locating the sampling sites. The elevation of the study sites ranges between 561 and 381 m ASL (Figure 3).

 Table 2: Sample sites, location and elevation along the Toposequence of Middle and lower slopes

	reposeduence of thirdale and to wer propes								
Area	Sample site	North	East	Elevation					
				(m)					
	AT009	12°08'32.00''	31°18'05.00''	561					
Al Abbasyia	AT008	12°09'21.00''	31°27'41.00''	541					
	AT007	12°05'09.00"	31°39'35.00''	523					
	AT006	12°03'51.00''	31°44'42.00''	503					
Tenten	AT005	11°55'52.00"	31°45'20.00''	473					
Terter	AT004	11°55'57.00''	31°52'28.00''	451					
Magania	ME003	11°56'02.00''	32°15'43.00''	437					
Megenis	ME002	11°56'22.00''	32°33'55.00''	406					
White Nile	ME001	11°56'52 00''	32°40'43 00''	381					
Terrace	1112001	11 50 52.00	52 10 15:00	501					



International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

	AT009	AT008	AT007	AT006	AT005	AT004	ME003	ME002		
ĺ	Abbasyia 4	Abbasyia 3	Abbasyia 2	Abbasyia 1	Al Terter 2	Al-Terter 1	Megenis 3	Megenis 2		
5	imme 2: Londont scope showing the compling sites starting from Abbesvie (AT000) on the North Fast to Maganic (ME00									

Figure 2: Landsat scene showing the sampling sites starting from Abbasyia (AT009) on the North East to Megenis (ME002) on the South East

Eight profiles AT009, AT008, AT007, AT006, AT005 and AT004, ME002 and ME003 were selected along the transect in the study area AL - Abbasiyia, AL - Terter and Meginis (Figure 2 and Table 2).A pit of about 1×2 m in length with 1.5 m depth was dug for sampling at each site. The face of the pit to be used for observation, photographing and sampling should face the sun. Notes on morphology, horizons sequence and soil classification were recorded following the standard procedures(FAO 2006 and USDA 1999). Soil samples, one from each soil horizon were taken for laboratory analyses. The soil samples were air dried and crushed using a wooden mortar and pestle and sieved to pass the 2 mm sieve and laboratory analyses were performed on the less than 2 mm fraction. All the physical and chemical determinations were carried out according to the international procedure of soil analysis (Showgi 2011).

4. Results and Discussion

The study area is a broad, slightly sloping to undulating upland clay plains. The plains are dissected by Khors draining the Nuba Mountains hilly area, numoures low ridges (viens and dykes) separated by shallow depressions running through the area. The soils are formed <u>insitu</u> and on colluviums and alluvium derived from basement complex rocks. The origin of these clays is thought to belong to Quarternary alluvial deposits probably similar to other dark cracking clays of central Sudan (Pacheco and Dawoud, 1976).They form well developed Vertisols (Plates 1, 2, 3 and 40.

Field study showed, common surface vertical cracks 1 - 4 cm wide and 50 -100 cm deep, polygons and cracks of variable shape and dimensions were measured, surface gilgai microrelief were observed, soil color is dominantly very dark grayish brown (10YR 3/2) when moist throughout the profiles. The soils are of nearly uniform clay texture. Topsoil structure has loose granular mulch of thickness ranging from 3-5 cm, the subsoil has weak to moderate coarse and very

coarse subangular blocky structure, below that the structure is massive. subsoil and substrata layers have parallelepiped and wedge shaped structure aggregates. Slickensides was also observed in all profiles, reflecting expansion and contraction upon wetting and drying of the soil. It was also noted the presence of fine and medium tubular pores, small and large whitish irregular $CaCO_3$ concretions, fine and medium sand grains and polished quartz pebblse arecommon.

Chemical and physical analyses of soil samples are represented by two profiles; AT004 and ME002. The results at profile AT004 (Table 3) show that the soil reaction is slightly alkaline in allprofiles, pH ranging between 7.4 to 8.3 increasing with depth. The soil is non - saline, ECe values ranges between 0.4 to 0.83 dS/m. It is also none sodic as indicated by exchangeable sodium percentage (ESP) range is between 0.56 and 10.18. Cation exchange capacity is higher and it coincides with the clay content, increasing with depth, it ranges between 51 to 79.3 mel/100 g. Sodium Adsorption Ratio (SAR) range is 3.04 - 7.81. The soil is slightly - calcareous with low amount of CaCO₃ (5.2 - 8.9 %). Organic matter is low decreasing with depth (0.554 -1.46 %). Total nitrogen is also low (0.02 - 0.059%). Nutrient status expressed as extractable cations meq/100 g soil is generally high: Ca^{++} range is 11.55 - 3.4; $Mg^{++} 2.2 - 3.4$ 17.4; $K^+ 0.175 - 0.782$; Na⁺ 0.41 - 5.45. Soil texture is clay throughout the profile. clay content increases with depth, then it decreases at the lowest horizon. range is (55.2 - 78 %), silt increases with depth, it ranges between 13.44 - 30.8%, the sand fraction is higher in the top and lower horizons than in the two middle horizons, (8.26 - 24.66%). The textue of the lower most horizon 94 - 125 cm is relatively lighter as it has the highest sand content, a high silt content and the lowest clay content. it looks similar to the top - most horizon. The bulk density (BD) is relatively and high increasing with depth, (1.29 - 1.72 g\ cc). Porosity is relatively high at the topsoil decreasing with depth, ranges (33 - 51%) - (Tables 3).



Plate 1: Grass and Wood Land around Gadoom Al –Gattar Hill.



Plate 2: Wood and Grass land vegetation (Acacia Seyal) on Flat Vertisols at AL - Terter Area

Depth Ci	n Horizon	Particle	e size distribut	ion %	pH paste	ECe dS\m	CaCO ₃ %	OC %
.1.		Sand 2000-50	Silt 50-2	Clay <0.2	I I			
0 - 22	A11	6.44	33.29	60.27	7.80	0.58	4.6	1.40
22-68	A12	3.83	25.62	70.55	7.77	0.42	4.5	0.936
68 – 94	A13	2.73	25.92	71.35	7.75	0.6	6.72	1.24
94-125	AC	10.63	31.00	58.37	7.82	0.65	6.5	1.08
N%	C/N ratio	CECmeq/100 g	ESP	SAR	Sat%	P%	B.D g\cc	Porosit%
0.030	46.6	63.94	0.569	5.21	54.4	0.030	1.42	47
0.025	37.4	64.17	1.175	4,41	61.49	0.030	1.63	39
0.027	45.9	73.36	7.306	5.3	53.5	0.019	1.68	37
0.020	54	72.82	7.333	5.76	59.3	0.020	1.74	35

Table 3: Laboratory data for profile No. AT004

Extractable ca	Extractable cations, meq/100 g Exch. NaSaturation extract, soluble (meq/l) meq/100 g										
Ca ⁺⁺	Mg^{++}	Na ⁺	\mathbf{K}^+		Ča ⁺⁺	Mg ⁺⁺	Na^+	\mathbf{K}^+			
1.55	1.45	0.55	0.41	0.364	3	1.0	7.34	0.870			
2.45	1.90	0.37	0.86	0.754	2	1.0	5.38	0.580			
1.80	1.55	0.34	5.43	5.36	1	1.0	5.30	0.580			
1.90	2,02	0.34	5.43	5.34	2	1.5	7.60	0.058			

	Saturatio	on extract	soluble	(meq	/ 100 g)	
cation	S					anions
Ca ⁺⁺	Mg ⁺⁺	K ⁺	Na ⁺	CO3	HCO3.	CI-
0.163	0.054	0.040	0.046	0.0	0.122	0.144
0.123	0.061	0.036	0.106	0.0	0.122	0.153
0.053	0.053	0.031	0.069	0.0	0.093	0.080
0.110	0.089	0.034	0.083	0.0	0.110	0.133

Soil Taxonomy: Typic Haplusterts, isohyperthermic, fine clayey, montmorillonitic, calcareous (USDA 1999)

World Reference Base (WRB): Haplic Chromic Vertisols (Grumic/Mazic) – (FAO 2014)

The Chemical and physical analyses results forprofileME002 at Megeinis (Table 4) show that the soil reaction is slightly alkaline, pH ranging between 7.49 to 8.53 increasing with depth and the soil is none saline with ECe values ranges between 0.45 to 0.68 dS/m.The soils are slightly and highly sodic; ESP range is between 10.61 and 36.07. Cation exchange capacity is high and it coincides with the clay content, increasing with depth, it ranges between 44.4 to 72.28 mel/100 g. Sodium.

Adsorption Ratio (SAR) range is 3.44 - 6.83. The soil is slightly calcareous with low amount of CaCO₃ (4.18 - 6.66%).Organic matter is low decreasing with depth (0.52 - 1.34%). Total nitrogen is also low (0.02 - 0.08%). Nutrient status expressed as extractable cations meq/100 g soil is

Volume 6 Issue 5, May 2017 www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

generally high: Ca⁺⁺ range is 38.74 - 69.99; Mg⁺⁺ 13.74 - 19.99; K⁺ 0.481 - 0.613;Na⁺ 5.17 - 21.04.Soil texture is clay throughout the profile except in topsoil's it is clay loam. Clay content increases with depth. range is between (33.2 - 65.24 %), silt decreases with depth, it ranges between 20.12 - 34.72 %, the sand fraction is higher in the top and decreases with depth, (4.10 - 32.08%). So texture – wise the

higher horizon 0 – 25 cm is relatively lighter - texture, has the highest sand content, a high silt content and the lowest clay content. The bulk density almost (BD) is high increasing with depth, (1.44 - 1.71 g\ cc). Porosity is relatively high at the topsoil decreasing with depth, ranges (35.5 - 45.7%) - (Table 4).



Plate 3: Abandoned Cultivated Flat Land atMegeinis Area



Plate4: Regenerated Vegetation on Cultivated Flat Land (Vertisols) atMegenis Area

Depth	Horizon	Particle size distribution %			pH paste	ECe dS\m	CaCO ₃ %	O C%				
Cm		Sand 2000-50	Silt	Clay <0.2								
			50-2	-								
0 - 26	A11	28.26	28.66	43.08	7.67	0.63	4.94	1.34				
26 - 43	A12	16.04	25.92	58.04	8.03	0.62	5.58	0.72				
43 – 87	A13	18.05	20.12	61.83	8.53	0.63	5.3	0.52				
87 -125	AC	18.84	20.95	60.21	7.71	0.64	6.66	0.93				

Table 4:	Laboratory	data for	profile No.	ME002
----------	------------	----------	-------------	--------------

N%	C/N	CEC	ESP	SAR	Sat%	P%	B.D	Porosity
	ratio						g∖cc	%
0.08	3.00	44.40	13.08	4.35	53.06	0.861	1.54	41.9
0.05	24.20	55.55	20.42	5.29	62.26	0.846	1.58	40.4
0.03	26.00	59.83	36.07	5.22	68.48	0.846	1.60	39.7
0.04	23.25	72.85	16.00	6.13	63.49	1.504	1.71	35.5

		Extract	able <u>catio</u>	ns meq	į/100g	g Exch	-			
	Na Saturation extract soluble(meq/l) meq/l00g									
Ca ⁺⁺ Mg ⁺⁺ K ⁺ Na ⁺⁺ Ca ⁺⁺ Mg ⁺⁺ Na ⁺ K ⁺										
6.75	1.37	0.481	5.912	5.80	1.3	0.75	5.420	0.40		
6.75	1.87	0.542	12.37	12.22	1.0	0.50	4.558	0.39		
6.50	1.62	0.613	21.04	20.83	1.0	0.50	4.490	0.66		
3.87	1.99	.0.553	11.776	11.66	1.0	0.50	5.280	0.56		

	Saturation extract soluble (meq/ 100 g)										
<u>c</u>	cations anions										
Ca ⁺⁺	Mg ⁺⁺	K+	Na ⁺	CO3 ⁻	HCO ₃ -	Cl-					
0.042	0.024	0.013	0.112	0.0	0.165	0.231					
0.53	0.530	0.020	0.15	0.0	0.266	0.266					
0.048	0.024	0.081	0.217	0.0	0.290	0.290					
0.063	0.031	0.035	0.116	0.0	0.380	0.444					

Soil Taxonomy: Typic Haplusterts, fine clayey, montmorillonitic, isohyperthermic, calcareous(USDA 1999)

World Reference Base (WRB):HaplicChromic Vertisols (Grumic/Mazic, sodic) –(FAO 2014)

5. Conclusions

This research was conducted to characterize and investigated the chemical, physical and morphological properties of cracking clays at Middle and Lower clay plains of Eastern Nuba Mountains. Ninesite locations were selected for this study; these sites were along a transect starting at the Middle clay plains (Al-Abbasyia and Terter area) and ending at Lower clay Plains (Megenis area) to compare the different types of clay soils in these plains.

Themorphological and surface features observations showed that the soils are dominantly cracking clay soils formed on sloping, undulating and flat clay plains. The surface features are characterized by common surface vertical cracks 1 - 4 cm wide and 50 -100 cm deep and forming polygons of variable shape and dimensions.Surface gilgai microrelief were observed at islated virgin areas. Soil color is dominantly by very dark grayish brown (10YR 3/2) when moist throughout the profiles. The soils are of nearly uniform clay texture. Topsoil structure has loose granular mulch of thickness ranging from 3-5 cm, the subsoil has weak to moderate coarse and very coarse subangular blocky structure, below that the structure is massive. subsoil and substrata layers have parallelepiped and wedge shaped structure aggregates. Slickensides was also observed in all profiles, reflecting expansion and contraction upon wetting and drying of the soil. fine and medium tubular pores, small and large whitish irregular CaCO₃ concretions and fine and medium sand grains, polish quartz pebble arecommon.

The analytical data for soils from middle clay plains (represented by profile AT004) reflected that the soil textures are dominantly very fine clay particles (> 60%). The soil reaction is slightly alkaline in allprofiles increasing with depth. The soil is non – saline and it is also none sodic as indicated by exchangeable sodium percentage (ESP) Cation exchange capacity is higher and it coincides with the clay content, increasing with depth. Sodium Adsorption Ratio (SAR) is low to moderate. The soilsare slightly calcareous with low amount of CaCO₃.Organic matter is low decreasing

with depth and the total nitrogen is also low.Nutrient status expressed as extractable cations meq/100 g soil is generally high. Soil texture is clay throughout the profile, clay content increases with depth, silt increases with depth and the sand fraction is higher in the top and lower horizons than in the two middle horizons. The bulk density almost (BD) is high increasing with depth and porosity is relatively high at the topsoil decreasing with depth, ranges.

The analytical data for soils from lower clay plains (represented by profile ME002 reflected that the soil textures are dominantly very fine clay particles (> 60%) but slightly lower at topsoil (<45%). The soil reaction is slightly alkaline in all profiles, pH ranging between increasing with depth. The soils are none saline but slightly sodic and sodic as indicated by exchangeable sodium percentage (ESP. Cation exchange capacity is lower and it coincides with the clay content, increasing with depth. Sodium Adsorption Ratio (SAR) is low to moderate. The soil is slightly calcareous with low amount of CaCO₃. Organic matter is low decreasing with depth and the total nitrogen is also low. Nutrient status expressed as extractable cations meq/100 g soil is generally high.Soil texture is clay throughout the profile except in topsoil's it is clay loam. Clay content increases with depth. range is between, silt decreases with depth and the sand fraction is higher in the top and decreases with depth. The bulk density almost (BD) is high increasing with depth and porosity is relatively high at the topsoil decreasing with depth.

Although most of the cracking clay soils in Sudan are sharing similar origin (basaltic igneous materials) but their subsequent depositional processes, landscape geomorphology and local climate have affected their mineralogical, physical and chemical properties. This applies to the clay plains of Nuba Mountains as most soils are characterized by lower CEC, mild alkalinity and low calcium carbonate content. More research is required to investigate the behaviour of these soils under different land use types and management.

References

- [1] Abedine, G., H. Robinson. 1970. A study on Cracking in Some Vertisols of the Sudan. Geoderma (3) 223-241.
- [2] Ahmed, N., 1983. Vertisols There Genesis Morphology and Properties In Wilding, L.P. Smeck and G. Hall, Ed's vol.11 Soil orders Chpt.3. Elsevier. Ci, Amsterdam.
- [3] Blockhuis, W.A., L.H.S. Ochtman, K.H. Peters. 1964. Vertisols in the Gezira and Khashm El - Girba clay plains. Sudan, Paper for the V111th International Congress ofSoil Science, Bucharest. Trans 8th. Int. Congr. Soil. Sci No 5, 592 - 603.
- [4] **Dawoud, H.A. 1974**. Reconnaissance Soil Survey and Land Capability Classification of Habila South Extension – South Kordofan Province Soil Survey Administration Wad Medani,
- [5] **Dawoud, H.A. 1976**. Exploratory, Soil Survey of North and South Kordofan Ministry of Agriculture Fisheries and Forestry. Soil Survey Administration Wad Medani, Sudan.

- [6] Doka M. Ali, El-Abbas, O. O. A., E. Showgi, H. A. ,Dawoud, and M. S., Mohamedain (2016). Mineralogical properties of cracking clay Soils along toposequence at Eastern Middle and Lower Clay Plains of Nuba Mountains South Kordofan, Sudan. Int. J. Curr. Microbiol. App. Sci. (2016) 5(10): 1049-1065
- [7] **Eltom, O.A. 1972**. Reconnaissance Soil Survey of Habila Extension, Kordofan Province, Soil Survey Administration Wad Medani
- [8] Elias, E. A., F.M. Salih, A.A. Salih, F. Alaily. 2001. Selected Morphological Characteristics of Soils From Gezira Vertisols With Particular Reference To Cracking. Int Agro - physics, 15, 79-86.
- [9] Eswaran, H., F.H. Beinroth, P.F. Reich, L.A. Quandt. 1999. Vertisols: their properties, classification, distribution and management. Washington, DC: USDA Natural Resources Conservation Service. Online version:
- [10] FAO. 1970. Semi-Detailed Soil Survey of Parts of The Central Clay Plain. Technical Report No.3, Soil Survey administration, Wade Medani Sudan.
- [11] FAO. 2006. Guide lines For Soil Profile Descriptions. 4th edition. FAO, Rome Italy.
- [12] FAO 2014. World Reference Base for Soil Resources. International soil classification system for naming soils and creating legends for soil maps. WORLD SOIL RESOURCES REPORTS No.106. Food And Agriculture Organization Of The United Nations. Rome, Italy 2014
- [13] Kevie, Van der. W. T. Buraymah. 1987. Manual for land suitability classification for Agriculture part (11). Giude line for soil survey parley chief's Technical report No (21). Soil Survey Administration Wad-Medani Sudan.
- [14] **Khalil, A.R. 1986.** Genesis and Ecology of Vertisols in Eastern Sudan Keil, West Germany PhD Thesis.
- [15] Khoddarry, O.E., 1978. Reconnaissance Soil Survey and Land Capability Classification of Elmashisha – and El – Haraza schemes area, Soil Survey Administration, Wad Medani
- [16] Pacheco R. and H. A., Dawoud. 1976. Exploratory, Soil Survey of North and South Kordofan Ministry of Agriculture Fisheries and Forestry. Soil Survey Administration Wad Medani, Sudan.
- [17] Showgi, Omima O. A. 2011. A study on properties and genesis of cracking clay soils in Southern Kordofan, Sudan. Ph.D Thesis. Soil and Water Science Dept. College of Agricultural Studies. Sudan University of Science and Technology. Khartoum.Sudan
- [18] Showgi, O. O. A., E. Doka M. Ali, El-Abbas, H. A. ,Dawoud, and M. S., Mohamedain (2016). Mineralogical properties of cracking clay Soils along toposequence at Northern Upper Clay Plains of Nuba Mountains South Kordofan (Sudan). Int.J.Curr.Microbiol.App.Sci (2016) 5(8): 745-759
- [19] **USDA 1999**. Key to Soil Taxonomy, 2nd, Ed. Agric .Hand Book No .436. USDA – NCS. Washington, D.C.
- [20] Vail, J.R. 1973. Outline of the Geology of the Nuba Mountains and Vicinity Southern Kordofan Province Sudan, Ministry of Industry and Mining, Geology Department, University Of Khartoum.
- [21] **Van der Kevie. 1973.** Climatic Zones in the Sudan Soil Survey Administration Wad Mandeni Sudan

Volume 6 Issue 5, May 2017

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

DOI: 10.21275/7041714