

# Detection of *Toxoplasma gondii* Infection among Pregnant Women in Hail, Saudi Arabia

Omar H. Amer<sup>1</sup>, AliS. AL Salem<sup>2</sup>, Husam K. Albluwe<sup>3</sup>, Bashr K. Alghazi<sup>4</sup>, Ibrahim H. Alshammari<sup>5</sup>

<sup>1</sup>Clinical Laboratory Sciences Dept., Faculty of Applied Medical Sciences, Hail University

<sup>2</sup>Hail Regional Lab

<sup>3,4,5</sup>Faculty of Medicine, Hail University

**Abstract:** *Toxoplasmosis is caused by infection with intracellular protozoa parasite T.gondii . One form of the disease, congenitally acquired toxoplasmosis, can result in sever central nervous system malformation or perinatal mortality. Other forms of the disease range from chronic, asymptomatic toxoplasma infection to ocular infection and adult disseminated cerebral toxoplasmosis. This research shows the number of cases which examined in the regional lab in Hail region, 5537 samples for women in the first few months of pregnancy has been tested for toxoplasmosis by using ELISA technique 1621 samples show positive result for TOXO-IgG which constitutes 9.8% of the total samples, which indicates that the 68% of pregnant women have an old infection. In addition, the samples were examined by TOXO-IgM to find out the number of pregnant women that have been infected by toxoplasmosis recently, the study show 17 pregnant women positive for TOXO-IgM, this means that 0.3% of pregnant women are infected by Toxoplasma parasite recently.*

**Keywords:** *Toxoplasma gondii*, Prevalence, IgG & IgM, Pregnant women, Hail, Saudi Arabia

## 1. Introduction

*Toxoplasma gondii* (*T. gondii*) is an obligate intracellular protozoan parasite causing toxoplasmosis, which is one of the most predominant chronic infections affecting one third of the human around the world (1-3). The importance of toxoplasmosis in pregnant women comes from the high prevalence of *T. gondii* infection and its severe consequences to the fetus and infant (4,5), and prevention from infection of the fetus and complications by antibiotic treatment has not been very effective (6). Diagnosis of *T. gondii* infections during pregnancy is mostly done by detecting immunoglobulin M (IgM) and immunoglobulin G (IgG) antibodies by using serological methods such as latex agglutination test, ELISA, and indirect immunofluorescence antibodies test (7-10). Congenital transmission may occur when an uninfected mother acquires primary infection during pregnancy. Primary infection with *T. gondii* in pregnant women occurs all over the world with frequencies between 0.1% and 1%. (11-12) Few studies have been conducted to explore the seroprevalence of *T. gondii* among pregnant Saudi women, with varying results. (13-16) Pregnant women with primary toxoplasmosis infection are often asymptomatic or have only mild symptoms, and approximately 70%-90% of infants born with congenital toxoplasmosis are asymptomatic at birth. Nevertheless, infection may cause spontaneous abortion, still birth, prematurity or serious fetal damage. (17,18,19) to determine the prevalence of toxoplasma antibodies in pregnant women in Hail city, KSA. In 2013, 6076 pregnant women were examined for IgG and IgM antibodies using ELISA technique. The age range was 19-43 years. The overall IgG seroprevalence was 9.8% and IgM seroprevalence was 0.6%. The IgM is indicative of low recent exposure to the parasite. In conclusion, the overall seroprevalence indicate a very low percentage in pregnant

women living in Hail, KSA. This lowers the risk of contracting *T. gondii* infections which minimize the risk congenital toxoplasmosis. Most recently, in Makkah, a study reported that seroprevalence of IgG was 21.2% and that of IgM was 1.2% in a group of pregnant women with history of previous abortion.<sup>31</sup> This seems to be very near to the findings of the current study, perhaps due to similarity of the selected group of patients, or more similar demographic and culture factors affecting transmission of the infection in both holy Islamic cities. The varying results from those regions of the Kingdom may be due to differences in patient numbers, demographics, history and used techniques, or the results may actually point to more public awareness and implementation of preventive measures against toxoplasmosis, such as in Almadinah Almunawwarah. Epidemiological studies suggest that prevalence of *T. gondii* infection in pregnant women varies substantially among different countries; in Europe it varies from 9% to 63%, 63.2% in Germany (20) 19.8% in Italy (21), and 9.1% in the UK (22). In Asian countries the seroprevalence of toxoplasmosis was reported low: 3.7% in Korea and 11.2% in Vietnam (23-24) while prevalence is as high as 41.6% to 45% in Indian pregnant women (25), 66.9% in Jordan and 53.1% in Kuwait (26, 27). In the American continent, the seroprevalence of toxoplasmosis was reported to be 77.5% in Brazil (28) and 63.5% in Colombia (29).

The aim of this study to find out how many pregnant women in Hail region infected with toxoplasma through examination of serum sample in serology department in regional lab using ELISA technique for toxoplasma IgG and IgM, with trying to find out the number of abortion cases due to this parasite

All of serum samples that have been sent to regional lab from hospitals and dispensaries in hail region which collected from pregnant women to follow up in year 2016

## 2. Material and Methods

### 1-Toxoplasma IgG ELISA by United Diagnostic Industry (UDI)

The UDI EG127 Toxoplasma IgG test system is an Enzyme Linked Immunosorbent Assay kit providing material for the detection of IgG –class antibodies to Toxoplasma gondii parasite in human serum or plasma.

#### Principle of the Test:

The EG127 Toxoplasma IgG kit is based on the ELISA technique. In the assay, calibrators and unknown are incubated in microtitration wells coated with purified and inactivated T.gondii antigen. After incubation and washing, the wells are treated with conjugate, composed of anti-human IgG antibodies labeled with peroxidase. After a second incubation and washing step, the wells are incubated with the substrate tetramethylbenzidine (TMB). An acidic stopping solution is then added and the degree of enzymatic turnover of the substrate is determined by wavelength absorbance measurement at 450 nm. The absorbance measured is directly proportional to the concentration of anti –T.gondii IgG antibodies present.

#### Materials:

##### Antigen –coated microtitration strips:

One strip holder containing 12x8 (96) microtitration wells coated with *T.gondii* antigen

##### Wash concentration:

One bottle containing 100ml of a phosphate buffered saline. Dilute with deionized or distilled water 1:10 prior to use

##### Sample diluent:

One bottle containing 100ml of BSA solution with 0.09% sodium azide as a preservative.

##### Toxoplasma IgG calibrators:

Each 2ml of human serum calibrated according to WHO reference standards Anti –Toxoplasma IgG in a 0.01M phosphate buffer containing BSA with 0.09% sodium azide as a preservative.

##### 2<sup>nd</sup> antibody conjugate:

One bottle containing 12ml of anti –human IgG monoclonal antibodies labeled with peroxidase in a phosphate buffer solution with 0.02% proclin.

##### TMB –substrate:

One bottle containing 12ml of tetramethylbenzidine (TMB) and hydrogen peroxide stabilized in citrate buffer with pH 3.8

##### Stopping solution:

One bottle containing 15ml of 0.3M H<sub>2</sub>SO<sub>4</sub>

### 1-Toxoplasma IgM ELISA by United Diagnostic Industry (UDI)

The UDI EM127 Toxoplasma IgM ELISA is an Enzyme Linked Immunosorbent Assay kit providing material for

detection of IgM –class antibodies to T.gondii in human serum or plasma

#### Principle of the Test:

The toxoplasma IgM assay is based on the principle of the capture of these immunoglobulins and subsequent identification of those, which are specific, making use of their ability to bind an antigen conjugated to peroxidase. The capture is performed using monoclonal antibodies bound to the solid phase (microtitration strips). The antigen is composed of purified an inactivated *Toxoplasma gondii* antigen

#### MATERIALS:

##### Antibody –coated microtitration strips:

One strip holder containing 12x8 (96) microtitration wells coated with anti –human IgM antibodies.

##### Wash concentration:

One bottle containing 100ml of a phosphate buffered saline. Dilute with deionized or distilled water 1:10 prior to use

##### Sample diluent:

One bottle containing 100ml of BSA solution with 0.09% sodium azide as a preservative.

##### Toxoplasma IgM controls:

Three vials, negative, cut off and positive each 2ml of human serum in a 0.01M phosphate buffer with BSA containing 0.09% sodium azide as a preservative.

##### Toxoplasma –HRP –Conjugate:

One bottle containing 12ml of purified toxoplasma antigen conjugated with peroxidase in a phosphate buffer solution with 0.02% proclin.

##### TMB –substrate:

One bottle containing 12ml of tetramethylbenzidine (TMB) and hydrogen peroxide stabilized in citrate buffer with pH 3.8

##### Stopping solution:



One bottle containing 15ml of 0.3M H<sub>2</sub>SO<sub>4</sub><sup>(4)</sup>

## 3. Results

In the present results cases which examined in the regional lab in Hail region, 5537 samples for women in the first few months of pregnancy has been tested for toxoplasmosis by using ELISA technique 1621 samples show positive result for TOXO-IgG which constitutes 9.8% of the total samples, which indicates that the 68% of pregnant women have an old infection.

In addition, the samples were examined by TOXO-IgM to find out the number of pregnant women that have been infected by toxoplasmosis recently, the study show 17 pregnant women positive for TOXO-IgM, this means that 0.3% of pregnant women are infected by Toxoplasma parasite recently. Table.1 and Figure. 1

TOXOPLASMOSIS STATISTIC IN 1437H		
HAIL REGIONAL LAB		
	NEGATIVE	POSITIVE
IgG	758	1621
IgM	3141	17
TOTAL IgG	2379	
TOTAL IgM	3158	
Total Patients Number	5537	



#### 4. Discussion

In the present study, a rapid, inexpensive serodiagnostic test (one step *Toxoplasma*-IgM/IgG test, InTec) was used. This test was simple to perform in only 15 min without special equipment or experienced personnel. (30-33)

In our study, IgG seroprevalence rate among the study subjects in Hail was 29% and IgM seroprevalence was 0.3% among pregnant women. Compared to rates previously reported from other KSA cities, this rate is found to be higher than those of Almadinah Almunawwarah 21.3%, but it is slower than the rates reported from Abha (34) in which IgG seroprevalence was 31.6%, and Makkah; 35.6% and 29.4%, in 2002 and 2006 among pregnant women. (14 and 35). However, some of them develop manifestations at childhood and adolescence. The seroprevalence of *T. gondii* in Hail province is 9.8% which is lower than the other cities with the results of the previous cross sectional studies carried out in other regions in Saudi Arabia and neighboring Gulf

countries, showing that the prevalence of *T. gondii* ranges between 25 and 36% (36-40). Low prevalence rates of 10% were reported in the United Kingdom (22), Korea (23), and Norway (41) and as high as 77.5% in Brazil (28) and 63.5% in Colombia (29). Higher prevalence rates were also reported in some neighboring Arab countries like Jordan 66.9% and Kuwait 53.1% (26,27). The difference in prevalence rate between different countries can be explained by variation of geographical and climatic conditions between different areas as the success of oocysts sporulation better in hotter and wetter areas (42).

#### References

- [1] Sonar SS, Brahmhatt MN. Toxoplasmosis: an important protozoan zoonosis. *Vet World* 2010; 3(9): 436-9.
- [2] Tekkesin N. Diagnosis of toxoplasmosis in pregnancy: a review. Telangana: Herbert Publications Pvt. Ltd.; 2012. [Online] Available from: <http://www.hoajonline.com/journals/pdf/2050-0874-1-9.pdf> [Accessed on 18th December, 2015]
- [3] Robert-Gangneux F, Dardé ML. Epidemiology of and diagnostic strategies for toxoplasmosis. *Clin Microbiol Rev* 2012; 25(2): 264-96.
- [4] Alvarados-Esquivel C, Estrada-Martinez S, Liesenfeld O. *Toxoplasma gondii* infection in workers occupationally exposed to unwashed raw fruits and vegetables: a case control seroprevalence study. *Parasit Vectors* 2011; 4: 235.
- [5] Chaudhry SA, Gad N, Koren G. Toxoplasmosis and pregnancy. *Can Fam Physician* 2014; 60: 334-6. *Fam Physician* 2014; 60: 334-6
- [6] Montoya JG, Remington JS. Management of *Toxoplasma gondii* infection during pregnancy. *Clin Infect Dis* 2008; 47: 554-66.
- [7] SYROCOT (Systematic Review on Congenital Toxoplasmosis) study group, Thiébau R, Leproust S, Chêne G, Gilbert R. Effectiveness of prenatal treatment for congenital toxoplasmosis: a meta-analysis of individual patients' data. *Lancet* 2007; 369(9556): 115-22.
- [8] Liesenfeld O, Wong SY JSR, editors. *Cecil text book of medicine*. 21<sup>st</sup> ed. Philadelphia: Saunders; 2000.
- [9] Liu Q, Wang ZD, Huang SY, Zhu XQ. Diagnosis of toxoplasmosis and typing of *Toxoplasma gondii*. *Parasit Vectors* 2015; 28: 292.
- [10] Gibbs RS. The origins of stillbirth: infectious diseases. *Semin Perinatol* 2002; 26(1): 75-8
- [11] J.S. Remington, R. McLeod, P. Thulliez, G. Desmonts J.S. Remington, J. Klein (Eds.), *Toxoplasmosis, infectious diseases of the fetus and newborn infant* (5th ed.), WB Saunders, Philadelphia, PA (2001), p. 207
- [12] S. Bahador, A.K. Samaneh Severe congenital toxoplasmosis: a case report and strain characterization. *Case Rep Infect Dis* (2015), Article 851085 <http://dx.doi.org/10.1155/2015/851085> 3 pages
- [13] H.M. El Hady. Toxoplasmosis among pregnant women in Abha, Saudi Arabia *J Egypt Soc Parasitol*, 21 (3) (1991), pp. 811-815



- [14] H.O. Ghazi, A.M. Telmesani, M.F. Mahomed TORCH agents in pregnant Saudi women Med PrincPract, 11 (4) (2002), pp. 180–182
- [15] H.I.A.T. Mohammad, M.H. Balaha, M.S. Moghannum Toxoplasmosis among the pregnant women attending a Saudi maternity hospital: seroprevalence and possible risk factors Ann Trop Med Parasitol, 104 (2010), pp. 493–504
- [16] I.M. Ashankyty Seroprevalence of *Toxoplasma gondii* among pregnant women visiting maternity hospital in Hail, KSA Life Sci J, 11 (8) (2014), pp. 355–359
- [17] F. Robert-Gangneux, J.B. Murat, H. Fricker-Hidalgo, M.P. Brenier-Pinchart, J.P. Gangneux, H. Pelloux The placenta: a main role in congenital toxoplasmosis? Trends Parasitol, 27 (2011), pp. 530–536
- [18] C. L'Ollivier, M. Wallon, B. Faucher, R. Piarroux, F. Peyron, J. Franck Comparison of mother and child antibodies that target high-molecular-mass *Toxoplasma gondii* antigens by immunoblotting improves neonatal diagnosis of congenital toxoplasmosis Clin Vaccine Immunol, 19 (2012), pp. 1326–1328
- [19] J.G. Montoya, O. Liesenfeld Toxoplasmosis Lancet, 363 (2004), pp. 1965–1976
- [20] K. Fiedler, C. Hulsse, W. Straube, and V. Briese, “Toxoplasmosis-antibody seroprevalence in Mecklenburg-Western Pomerania,” ZentralblattfürGynäkologie, vol. 121, pp. 239–243, 1999.
- [21] L. Masini, L. Casarella, R. L. Grillo, M. P. Zannella, and G. C. Oliva, “Epidemiologic study on anti-*Toxoplasma gondii* antibodies prevalence in an obstetric population,” Italian Journal of Gynaecology and Obstetrics, vol. 20, no. 3, pp. 159–166, 2008.
- [22] J. Q. Nash, S. Chissel, J. Jones, F. Warburton, and N. Q. Verlander, “Risk factors for toxoplasmosis in pregnant women in Kent, United Kingdom,” Epidemiology and Infection, vol. 133, no. 3, pp. 475–483, 2005.
- [23] K. Han, D.-W. Shin, T.-Y. Lee, and Y.-H. Lee, “Seroprevalence of *Toxoplasma gondii* infection and risk factors associated with seropositivity of pregnant women in Korea,” Journal of Parasitology, vol. 94, no. 4, pp. 963–965, 2008.
- [24] P. Buchy, J.-Y. Follézou, T. X. Lien et al., “Serological study of toxoplasmosis in Vietnam in a population of drug users (Ho Chi Minh City) and pregnant women (NhaTrang),” Bulletin de la Societe de PathologieExotique, vol. 96, no. 1, pp. 46–47, 2003.
- [25] B. Borkakoty, A. Borthakur, and M. Gohain, “Prevalence of *Toxoplasma gondii* infection amongst pregnant women in Assam, India,” Indian Journal of Medical Microbiology, vol. 25, no. 4, pp. 431–432, 2007.
- [26] N. F. Jumaian, “Seroprevalence and risk factors for *Toxoplasma* infection in pregnant women in Jordan,” Eastern Mediterranean Health Journal, vol. 11, no. 1-2, pp. 45–51, 2005.
- [27] J. Iqbal and N. Khalid, “Detection of acute *Toxoplasma gondii* infection in early pregnancy by IgG avidity and PCR analysis,” Journal of Medical Microbiology, vol. 56, no. 11, pp. 1495–1499, 2007.
- [28] A. M. Porto, M. M. de Amorim, I. C. Coelho, and L. C. Santos, “Serologic profile of toxoplasmosis in pregnant women attended at a teaching-hospital in Recife,” Revista da AssociaçãoMédicaBrasileira, vol. 54, no. 3, pp. 242–248, 2008.
- [29] A. T. Castro, A. Congora, and M. E. Gonzalez, “*Toxoplasma gondii* antibodyseroprevalence in pregnant women from Villavicencio, Colombia,” Orinoquia, vol. 12, no. 1, pp. 91–100, 2008.
- [30] S.R. Béla, D.A. Oliveira Silva, P.C.P. Cunha-Júnior, F.A. Chaves-Borges, F. Reis de Carvalho, T. Carrijo de Oliveira, J.R. Mineo Use of SAG2A recombinant *Toxoplasma gondii* surface antigen as a diagnostic marker for human acute toxoplasmosis: analysis of titers and avidity of IgG and IgG1 antibodiesDiagnMicrobiol Infect Dis, 62 (2008), pp. 245–254
- [31] S. Jin, Z.Y. Chang, X. Ming, C.L. Min, H. Wei, L.Y. Sheng, G.X. HongFast Dipstick dye immunoassay for detection of immunoglobulin G (IgG) and IgM antibodies of human toxoplasmosisClinDiagn Lab Immunol, 12 (1) (2005), pp. 198–201
- [32] S.H. Jabbar, F.G. Haider, H.A. Abid Al-Razaq. Evaluation of rapid Chromatographic immunoassay with latex agglutination test and ELISA for diagnosis of human toxoplasmosis. J Fac Med Baghdad, 4 (2010), pp. 469–47
- [33] Y.H. Wang, X.R. Li, G.X. Wang, H. Yin, X.P. Cai, B.Q. Fu, D.L. Zhang Development of an immunochromatographic strip for the rapid detection of *Toxoplasma gondii* circulating antigens ParasitolInt, 60 (1) (2011), pp. 105–107
- [34] H. Aqeely, E.K. El-Gayar, D. Perveen Khan, A. Najmi, A. Alvi, I. Bani, M.S. Mahfouz, S.E. Abdalla, I.M. Elhassan Seroepidemiology of *Toxoplasma gondii* amongst pregnant women in Jazan Province, Saudi Arabia
- [35] S.A. Al-Harathi, M.B. Jamjoom, H.O. Ghazi Seroprevalence of *Toxoplasma gondii* among pregnant women in Makkah, Saudi Arabia Umm Al-Qura Univ. J Sci Med Eng, 18 (2) (2006), pp. 217–227.
- [36] A. R. Al-Qurashi, A. M. Ghandour, O. E. Obied, A. Al-Mulhim, and S. M. Makki, “Seroepidemiological study of toxoplasma gondii infection in the human population in the Eastern Region,” Saudi Medical Journal, vol. 22, no. 1, pp. 13–18, 2001.
- [37] A. M. D. Tonkal, “PCR versus ELISA in diagnosis of human Toxoplasmosis in Jeddah, Saudi Arabia,” Journal of the Egyptian Society of Parasitology, vol. 38, no. 3, pp. 707–714, 2008
- [38] A. R. Al-Qurashi, “Seroepidemiological study of Toxoplasmosis in rural areas in the eastern region of Saudi Arabia,” Journal of the Egyptian Society of Parasitology, vol. 34, no. 1, pp. 23–34, 2004.
- [39] S. A. Uduman, H. N. Mohamed, A. Bener, and F. K. Dar, “The prevalence of *Toxoplasma gondii* specific IgG and IgM antibodies in blood donors in Al Ain, United Arab Emirates indicates a potential risk to recipients,” Journal

of Communicable Diseases, vol. 30, no. 4, pp. 237–239, 1998.

- [40] M. A. Abu-Madi, N. Al-Molawi, and J. M. Behnke, “Seroprevalence and epidemiological correlates of *Toxoplasma gondii* infections among patients referred for hospital-based serological testing in Doha, Qatar,” *Parasites and Vectors*, vol. 1, no. 1, article 39, 2008.
- [41] P. A. Jennum, B. Stray-Pedersen, K. K. Melby et al., “Incidence of *Toxoplasma gondii* infection in 35,940 pregnant women in Norway and pregnancy outcome for infected women,” *Journal of Clinical Microbiology*, vol. 36, no. 10, pp. 2900–2906, 1998.
- [42] K. B. A. Kistiah, J. Winiecka-Krusnell, A. Karstaedt, and J. Frean, “Seroprevalence of *Toxoplasma gondii* infection in HIV-positive and HIV-negative subjects in Gauteng, South Africa,” *Southern African Journal of Infectious Diseases*, vol. 26, no. 4, pp. 225–228, 2011.

