Assessment of Nursing Students Knowledge Regarding Hepatitis B Virus at Iraqi Nursing Colleges

Juma JubarA. Al-Redha, PhD¹, Haider Mohammed Majeed, MSc²

¹Dr. Instructor, Fundamentals of Nursing Department, College of Nursing, University of Baghdad

²Assistant Instructor, Fundamentals of Nursing Department, College of Nursing, University of Baghdad

Abstract: <u>Background</u>: Hepatitis B virus is a serious, is a dangerous, global public health problem that leading to liver cancer / hepatic cirrhosis with a high rate mortality around 600000 individuals in world. Nursing students are also vulnerable to acquire this infection. <u>Objectives</u>: This study sought to evaluate the knowledge of undergraduates nursing students concerning viral B hepatitis infection and routes of transmission andprevention control andtodetermine the relationship between demographic characteristic with students knowledge. <u>Methodology</u>: A cross-sectional study, conducted in the Facultiesof Iraqi nursing; Baghdad University, Babel University, Mesan University, Kirkuk University, Tikrit University, Iraq; from December - April 2016. <u>Results</u>: The vast majority of the participants (88%) of the study sample were at (20-24) years, There were 79 male (63.2%) and 46 females (36.8%), concerning academic year (52%) at fourth class (80%) were single, equal distribution (20%) for university.Residents of urban areas represented 91(72.8%) of the study group.<u>Conclusions</u>: The study concluded that the nursing students have poor knowledge concerning HBV.

Keywords: Assessment, Knowledge, Hepatitis B virus, Nursing Students

1. Introduction

Hepatitis B is an infection of the livertissues caused by the hepatitis B virus (HBV). The two billions of individuals in worldwide it is appraises that have current or past hepatitis B infection, including more than 250 million chronic carriers and more than 600,000 deaths worldwide each year ^[1]·Hepatitis B is widespread in the Asia Pacific region and 10 to 15 million of the people suffer from this disease ^{[2].}Globally, According toWHO stated that that over 2 billion people have been infected with HBV and 170 million HCV infected patients world-wide. [3]. The infectious of viral B istransmission through many routes such as with plasma, blood, vaginal fluids, semen, saliva and mucous membranes. Also HBV can be transmitted from mother to child at birth (perinatal transmission), by sexual intercourse with an infected personify, by using contaminated needles, undependable blood transfusions and between children in the early childhood ^[4]. Individuals that high risk to prone viral infection include health care teams in contact with blood and human body fluid, secretions, hemodialysisunit staff, oncology and chemotherapy nurses, all personnel at risk of needle stick/sharps injuries, which includes those working in theater rooms and laboratories, respiratory therapists, surgeons, nursing, dentists, as well as medical, dental and nursing students. Therefore nursing students are the future healthcare staffsso that risk for infection with HBV exposed during their clinical training [5]. The nursing students because of insufficient training and limited clinical experience are at a much higher risk of exposure to such pathogens. They are liable to spread infection to the patients. So knowledge regarding hepatitis transmission, treatment, and preventive measures is crucial for its prevention. Adequate knowledge and sequent preventive measures could effectively reduce the risk of this important hazard. Infact the quality of nursing care depends to a large degree on the knowledge, skills, attitudes and activities of the practicing nursing staff. With this background, the current study was conducted with the objective to detection of the knowledge among undergraduate students towards HBV.

2. Materials and Methods

A cross sectional design was carried out among third and final year undergraduate students of nursing faculty for duration from December - April 2016infifth Iraqi nursing colleges these were selected randomized that includes (Baghdad nursing college, Bable nursing college, Mesannursing college, Kirkuk nursing college, Tikrit nursing college).To evaluate knowledge about viral B hepatitis among nursing students and to identify the relationship between demographic characteristics with students knowledge. The stratified random sampling method was utilized in selecting the participants. The inclusion criteria for the participants were age of 18 years old and above of both genders, male and female, which include third and fourth stage from each faculty. The sample size calculated was 125, inclusive of the 10% non-response rate. Proportional allocation from third& fourth grade levels (1 to 4 or 1 to 5 in all colleges). The study was done with the help of (15) specially prepared questionnaires which were validated by doing pilot study. Data collection tool was a multi section questionnaire; the first sector was about demographic information such as age, gender, academic year, university, marital status, and home. The second part consists of twenty-oneclose-ended questionsto evaluate the knowledge of the participant about hepatitis B, the knowledge questions, as follows: 21 affirmations, for which an "X" should be marked for one of the possible alternatives, (yes), (no) (yes it was considered as a correct question, (no it was considered as an incorrect question answered. The knowledge score ranged from (0 - 1) with higher score indicating higher knowledge. According to the score, the participants were poor knowledge (score: 0-o.49)

Volume 6 Issue 5, May 2017 www.ijsr.net

and good knowledge (score: 0.5-1).Statistical analysis of the data in this study was performed using SPSS V. 18.

3. Results

Table 1: Characteristics of study participants No Items Variables Frequency Percent							
No	Items	Variables	Frequency	Percent			
		20-24	110	88			
1-	Age	25-29	10	8			
		30-34	3	2			
		35-38	2	1			
		Total	125	100%			
2-	Gender	Male	79	63.2			
		Female	46	36.8			
		Total	125	100%			
	Academic year	Third class	60	48			
3-		Fourth class	65	52			
		Total	125	100%			
4-	Marital status	Single	100	80			
		Married	25	20			
		Total	125	100%			
		Baghdad	25	20			
5-	University	Bable	25	20			
		Mesan	25	20			
		Kirkuk	25	20			
		Tikrit	25	20			
		Total	125	100%			
		Urban	91	72.8			
5-	Living place	Rural	34	27.2			
		Total	125	100%			

Frequency, Percent

Table(1) demonstrate that the highest percentage (88%) of the study sample were at (20-24) years, the majority of them (63%) were male, concerning academic year (52%) at fourth class, (80%) were single, equal distribution (20%) for university, and highest (91%) urban living place.

Table 2: The	knowledge	of study sam	ple concerning	HBV
--------------	-----------	--------------	----------------	-----

No.	Items	Ŋ	Yes		No		Sig
		F	%	F	%		_
1-	Liver is a vital organ that processes nutrients, filters of blood, and fights infections.	57	45.6	68	54.4	0.45	N.S
2-	Hepatitis is most often caused by a virus	62	49.6	63	40.4	0.49	N.S
3-	The common types of viral hepatitis are hepatitis A, B, and C	74	59.2	51	39.8	0.59	S
4-	The incubation period of hepatitis B virus from 45 to 160 days	36	28.2	89	71.8	0.28	N.S
5-	Prevent or treat the commonlytypes ofhepatitis A & B can be prevented by vaccination.	41	32.8	84	67.2	0.32	N.S
6-	The incubation period of hepatitis B virus from 45 to 160 days	29	23.2	96	76.8	0.23	N.S
7-	Chronic hepatitis B	52	41.6	73	58.4	0.41	N.S

9- P							11.0
a h le li 8- H to c iii 9 a n	nd C are serious health problems can ead to cirrhosis , iver cancer Histologic evolution o chronic hepatitis can be demonstrated n approximately 90- 95% of cases of acute hepatitis B in heonates	38	30.4	87	69.6	0.30	N.S

F=Frequency P=Percentage M.S=Mean of score Cutoff point = 0.5

This table shows that the mean of score is significant at the items (1, 3) while it is no significant in other items.

 Table 3: The knowledge of study sample concerning methods of transmission

No	Items		Yes		No	M.S	Sig	
		F	%	F	%			
1-	At least 30% of reported hepatitis B among adults cannot be associated with an identifiable risk factor	51	40.8	74	59.2	0.40	N.S	
2-	The HBV cannot be spread by holding hands, sharing eating utensils, hugging, coughing, sneezing	53	42.4	72	57.6	0.42	N.S	
3-	Identified methods of transmission HBV include (blood transfusion, unsanitary sexually)	78	62.4	47	37.6	0.62	S	
4-	Through intercourse or contact with blood or bodily fluids.	53	42.4	73	57.6	0.42	N.S	
5-	From mother to child by breast feeding	61	48.8	64	51.2	0.48	N.S	
6-	Blood contact can occur by sharing syringes in intravenous drug use, shaving accessories such as razor blades,	44	35.2	81	64.8	0.35	N.S	
7-	Touching wounds on infected persons.	64	51.2	61	48.8	0.51	S	
F=F1 This item	requency P=Percentage M.S=Mean o s table demonstrate that the mear (5) while it is no significant in c	f sco 1 of other	re Cu score	t off is s	point : point :	= 0.5 cant a	t the	

Table 4:	The knowledge of study sample concerning
	Preventive measures of HBV

	Treventive measures of TID v						
No		Ŋ	ſes	l	No	M.S	Sig
	Items	F	%	F	%		
1-	Hepatitis infections can be prevented by avoiding exposure to viruses, and through injectable immune globulins or by vaccines	49	39.2	76	60.8	0.39	N.S
2-	Vaccination is recommended in the first day of life if possible	58	46.4	67	53.6	0.46	N.S
3-	A vaccine is	62	49.6	63	50.4	0.49	N.S

Volume 6 Issue 5, May 2017 www.ijsr.net

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

	available that will prevent infection from hepatitis B for life.						
4-	VaccinationB antigen is administer to stimulate the body's immune system to produce protective antibodies	56	44.8	69	55.2	0.44	N.S
5-	Before giving blood must act test of hepatitis virus infection before infusion to prevent infection	64	51.2	61	49.2	0.51	S

F=Frequency, P=Percentage, M.S=Mean of score Cut off point = 0.5

This table demonstrate that the mean of score is significant at the items (21) while it is no significant in other items

Table 5: The knowledge of study sample concerning
symptoms and diagnostic test

No	Items		Yes	No		M.S	Sig
		F	%	F	%		
1-	An illness that begins with general ill-health, loss of appetite, nausea, vomiting, body aches, mild fever	67	53.5	58	46.5	0.53	S
2-	The clinical manifestation are mild and nonspecific and diagnosis usually is delayed	69	55.2	56	44.8	0.55	S
3-	Symptoms of hepatitis includefatigue, tired. dark urine, light-colored stools,fever, and jaundice	71	56.8	54	43.2	0.56	S
4-	There are three types of blood tests for evaluating patients with hepatitis: liver enzymes, antibodies to the hepatitis viruses, and viral proteins or genetic material (viral DNA or RNA).	57	45.6	68	54.4	0.45	N.S
5-	Liver enzymes (AST, SGOT, ALT and SGPT)widely used blood tests for evaluating patients with hepatitis	79	63.2	46	36.8	0.63	S

This table demonstrate that the mean of score is significant in all items except item (4).

Table 6:	The differences between Demogr	aphic
Chara	cteristics and Student's knowleds	e

					0	
Variables	Source	Sum of	dr	Mean	F	Sig.
	of	secure		score		
	variance					
Age	Between	3.968	3	1.323		
	group					
	Within	306.436	38	8.064	.164	.920
	group					
	Total	310.405	41			
Gender	Between	21.460	1	21.460	2.071	002
	group				2.971	.092

	Within	288.944	40	7.224		
	group					
	Total	310.405	41			
Level of	Between	7.690	2	21.460		
education	group					
	Within	9.350	40	7.224	2.971	.015
	group					
	Total	17.040	42			
Marital	Between	19.921	2	9.960		
status	group					
	Within	290.484	39	7.448	1.337	.274
	group					
	Total	310.405	41			

df =Degree of freedom , F= F-statistics , Sig.=level of Significance

This table shows no significant differences between demographic characteristics and knowledge of students.

4. Discussion

In this study we observed that that majority(88%) of the study sample within age group (20-24). This findings supported by a study conducted to assess the knowledge and practice of students regarding HBV, who stated that (82.6%) of the subjects were of 20 and 21 years of age ^{[6].} Another study done in Egyptthe researcher reported that they mentioned that mean age of the all participated was 8 89 \pm 0.98 years (range 17- 23)^{[7].} In relation of gender the most (63%)of respondents were female, this result is compatible with results of study conducted by other researcher who stated that the participation group consisted of 179 women students (77%) and 54 men students (23%)^[8]. Concerning to academic year most of study sample (52%) at fourth year this similar to finding who reported that (39%) of the sample were fourth academic year^{[9].} About marital status the current study showed that the majority(80%) were single, this result disagree with other study who stated that it was noticed that (60.6% and 65.1% respectively) from his sample were married ^{[10].} Our study demonstrate that (72.8)urban living place it is agree with findings obtains from other research who observed that the vast majority of their studied samples were from urban area^[11]. While this results disagree with another study who mentioned that most of his study sample (52.8%) were rural ^{[7].} Regarding to the students knowledge concerning HBV (table 2) the study shows the participants had poor knowledge about VBH this results agree with other study who stated that the present study showed that the knowledge of students about hepatitis B and C and the use of preventive measures was inadequate before education and improved after participation in the health educational program^{[7].} Another studyconducted by researchersaid that the survey indicated he level of knowledge between students toward hepatitis B and C, was fair knowledge, but gaps in knowledge were identified which need to be strengthened in students^{[12].} In relation to Methods of transmission (table 3) we found that the students had no sufficient prior knowledge, (WHO, 2012) mentioned that it is considered to be a serious gap of knowledge since transmission of HBV through sexual relationships is one of the most common ways of transmission^{[13].} Bari (2001) explains that initially it was believed that transfusion was the most common route of transmission of hepatitis. Other modes of transmission include unsafe injections (unsafe

Volume 6 Issue 5, May 2017 www.ijsr.net

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

syringe), unsafe blades for face or armpit shaving at community barber shops, ear/nose piercing and tattooing^[14].Table(3) shows the student's knowledge had low score in relation to preventive measure, prevention for transmission is extremely important, both for the individual person and seen from a societal perspective. It is therefore important to give the right education to health care personnel (Abiola et al., 2013 & Carter et al., 2013)^[15, 16]. According to the study done by other researcher who reported that people are more likely to get vaccinated against HBV if they believe that the vaccine is protective^{[17].} Regarding to knowledge for HBV the study reveals that the participants had fair knowledge, this result same to Shikure & Fete who said in his study that about the appearance of hepatitis B viral infection symptoms; (27.0%) of the study participants respond correctly that the symptoms doesn't appear within few days always after the entry of hepatitis B virus to the body, while (50.4%) and (22.6%) answer incorrectly and doesn't know about the topic respectively^[9]. Current study indicates that there are no significant differences between demographic characteristics (table 6) this findings disagree with results obtain from other researcher who explain that the academic years of study were significantly associated with knowledge of hepatitis B virus infection $(p=0.000)^{[18]}$.

5. Conclusions

The study concluded that the nursing student have poor knowledge concerning HBV.

6. Recommendations

Our study recommended that the need of strengthen the students knowledge about HBV by health education program and booklets and distribute to the students.

7. Acknowledgements

The authors would like to thank all data collectors and study participants involved in the study.

References

- K Park. Park's Textbook of preventive and Social Medicine, 22nd edition, Jabalpur: Banarsidas Bhanot publishers; 2015:495.
- [2] al-Haq N, Hassali MA, Shafie AA, Saleem F, Farooqui M, Aljadhey H. A cross sectional assessment of knowledge, attitude and practice towards Hepatitis B among healthy population of Quetta, Pakistan. BMC Public Health 2012;12:692.
- [3] World Health Organization. Hepatitis Fact sheet N204 Revised August 2008. Key facts. Where is Hepatitis B most common? Available at: www.who.int/mediacentre/factsheets/fs204/-29k. Accessed 2 Feb 2015.
- [4] The Public Health Authority of Sweden. (2015). Sjukdoms information omhepatit B. Stockholm: Folkhälsomyndigheten. Collected 21 October, 2015, fromhttp://www.folkhalsomyndigheten.se/amnesomrade n/smittskydd-och-

sjukdomar/smittsammasjukdomar/hepatit-b/

- [5] Smelzer SC and Bare B. 2003. Brunner and Suddarth's Textbook of medical surgicalnursing.10th edition.
- [6] Anita Kumari; Assessment of Knowledge and Attitude Regarding Hepatitis B and C among Nursing Students; ADR Journals 2015.
- [7] Ghona Abd El-Nasser and Eman Abed El baset; Assessment of Students Knowledge and Attitude Toward Hepatitis B and C in Sohag University/Egypt; Kufa Journal for Nursing Sciences Vol. 3 No. 1, January through April 2013.
- [8] Elin Dahlström and Ellinor Funegård Viberg; Knowledge about hepatitis B virus infection and attitudes towards hepatitis B virus vaccination among Vietnamese university students in Ho Chi Minh City: The Bachelor Programme of Science in Nursing 180 ECTS credits 2013.
- [9] Hikma Shikure : Assessment of Knowledge, Attitude and Practice Toward Hepatitis B Viral Among Addis Ababa University, Regular Undergraduate Students of College of health Science, Addis Ababa Ethiopia, 2014 Addis Ababa University College of Health Science, School of Allied Health Sciences Department of Nursing and Midwifery.
- [10] Shimaa Abdelrahim, Madiha Mohamed, Safaa Ahmed, Mohamed Zakria; Seroprevalence, Knowledge, Attitude and Practices among Barbers and Their Customers Regarding HCV and HBV in Assiut District, Egypt: IOSR Journal of Nursing and Health Science, Volume 4, Issue 3 Ver. I (May. - Jun. (2015), PP 19-30.
- [11] Jokhio A, Bhatti T and Memon M,; Knowledge, attitudes and practices of barbers about hepatitis B and C transmission in Hyderabad, Pakistan. Eastern Mediterranean Health Journal, Vol. 16 No.10,(2007): p 1080.
- [12] Razi' A, urRehman, R, Naz, S, Ghafoor, F, & Khan, A M U: Knowledge Attitude and Practices of University Students Regarding Hepatitis B and C. Journal of Agricultural and Biological Science, (2010).5(4): 38-43.
- [13] World Health Organization (who.int) Prevention &Control of Viral Hepatitis Infection: Framework for Global Action, Available from http: www who intcsr disease hepatitis GHP_framework .Rawalpindi-Islamabad Pakistan. Trop Med Int Health. (2012) 6: 732-738.
- [14] Bari A, Akhtar S, Rahbar MH, Luby SP.. Risk factors for hepatitis C virus infection in male adults in Rawalpindi-Islamabad Pakistan. Trop Med IntHealth. (2001) 6: 732-738.
- [15] Abiola, A., Akodu, B., &Omoyeni, O. Knowledge, attitude and practice of hepatitis B vaccination among health workers at the Lagos State accident and emergency centre, Toll-Gate, Alausa, Lagos State. West African Journal of Medicine, (2013). 32(4): 257-262.
- [16] Carter, T., Damico, A., DeHart, M., Halsey, N., Marcuse, E., Mergler, M., Navar-Boggan, A., Omer, S., Orenstein, W., Pan, W., Salmon, D., & Taylor, J.; Association of vaccinerelated attitudes and beliefs between parents and health care providers. *Vaccine*, *31*(41): 4591-4595. doi: 10.1016/j.vaccine.2013.07.039
- [17] Ma, G.X., Shive, S.E., Fang, C.Y., Feng, Z., Parameswaran, L., Pham, A. & Khanh, C. (2007).Knowledge, attitudes, and behaviors of hepatitis

Volume 6 Issue 5, May 2017

<u>www.ijsr.net</u>

B screening and vaccination and liver cancer risks among Vietnamese Americans. *Journal of Health Care for the Poor and Underserved*, *18*(1), 62-73. doi:10.1353/hpu.2007.0013.

[18] Taranga Reang, Tanushree Chakraborty, Mousumi Sarker, Amar Tripura: A study of knowledge and practice regarding Hepatitis B among nursing students attending tertiary care hospitals in Agartala city *Int J Res Med Sci. 2015 Jul;3(7):1641-1649*

Volume 6 Issue 5, May 2017 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY