

# Model for Estimation of Intrauterine Fetal Weight for Sudanese

Dr. Elsir Ali Saeed Taha<sup>1</sup>

<sup>1</sup>National University, college of radiographic and medical imaging, Khartoum, Sudan

**Abstract:** *The aim of the present study was to estimate fetal weight antenatally at or near term by using ultrasound evaluation and measurement. The study was a descriptive study conducted on 232 pregnant women selected by simple random sampling who attended the Elribat national hospital and Khartoum diagnostic center in periods from January 2016-till January 2017. Ultrasound examination was performed and FL, BPD, AC and Fetal weight was estimated by using Hadlock's formula ultrasonographically. The mean value of calculated fetal weight was 1526.4±1007.7gm. Finally the correlation was done in order to predict models for estimation for future work.*

**Keywords:** GA-LMP, mensuration, ultrasound, pregnancy

## 1. Introduction

Present day obstetrics has in fact rightly been able to focus on the concept of fetal medicine as distinct and significant entity in view of rapid decline in maternal mortality and morbidity with simultaneous recognition of various forms of fetal handicaps affecting the overall perinatal mortality and morbidity. Growth is a basic fundamental of life. Assessment of fetal weight in utero leads to an improved prospective management of high risk pregnancies and considerable reduction in perinatal mortality and morbidity. It has become increasingly important especially for prevention of prematurity, evaluation of fetopelvic disproportion, induction of labor before term and detection of IUGR. Thus a quick, easy and accurate method for estimating the fetal weight in utero with optimum precision would be of obvious benefit to the clinical practicing modern obstetricians. Estimation of birth weight by Johnson's formula based on symphysiofundal height has advantages of speed, economy and general applicability. Obstetric ultrasound has in fact revolutionized the knowledge of fetal medicine in the present day and can predict fetal weight with a great degree of precision. Sowjanya and Lavanya (2015).

### 1.1. Obstetrical ultrasonography

Early expectations that this method might provide an objective standard for identifying fetus of abnormal size for gestation age was recently undermined by prospective studies that showed sonographic estimates of foetal weight to be no better than clinical palpation for predicting foetal weight (Sherman et al. 1998), Ratanasiri et al (2002), Bossak and Spellacy (1972), Banin et al. (2002), Titapant et al. (2001)). Susiki et al (1974), used ultrasound measurement of fetal heart volume to estimate fetal weight. Paulos et al. (1953) used fetal volume by ultrasound. Sonographic predictions based on algorithms using various combinations of fetal parameters ,such as abdominal circumference (AC), Femur Length(FL), Biparietal Diameter (BPD),and Head Circumference (HC) both singly and in combination have been used (Nzeh et al. (2002), Nahum (2002), Hadlock et al. (1985), Nzeb et al. (1992), Campbells and Wilkin (1975)).When other sonographic fetal measurements are used

for estimating fetal weight e.g. humorous soft tissue thickness, ratio of subcutaneous tissue to, femur length, cheek to cheek distance, these nonstandard measurements do not help to predict birth weight except in special subgroups e.g. Diabetic mothers (Abramowicz et al. (1991)). Multiple sonographic foetal biometry also do not improve prediction (Nzeb et al. (1992), Ratanasiri et al. (2002)). Foetal imaging is limited by maternal obesity, oligohydrammios and anterior placentation. Besides these formulas are obtained from populations which do not include pregnant women of all genetic background resulting in an inherent sampling error.

In some studies, ultrasound has been shown to determine the weight of the fetus to within 10% of the actual birth weight in as many as 75% of pregnancies and within 5% in as many as 40%. Watson et al. (1988).

### 1.2. Clinical risk Factor

This method involves quantitative assessment of clinical risk factors and has been shown to be valuable in predicting foetal weight. In case of foetal macrosomia ,the presence of risk factors ,such as maternal diabetes mellitus ,prolonged pregnancy, obesity ,pregnancy weight gain of >20 kg, maternal age >35 years, maternal height> 5 ft 3 inches, multiparity, male foetal sex and white race should be added. In low estimated birth weight socioeconomic status, constitutionally small mother, poor maternal weight gain, foetal infections, congenital malformations, chromosomal abnormality, teratogenic exposure, maternal anaemia, Anti phospholipid Antibody syndrome and other medical disorders complicating pregnancy should be mentioned.

### 1.3. Maternal Self estimation

In literate society maternal self-estimation of fetal birth weight in multiparous women show comparable accuracy to clinical palpation in some studies for predicting abnormally large fetus (Chauhan et al. (1992), Banin et al. (2002)).

### 1.4. Birth weight Prediction equations:

Various calculations and formulae based on measuring

uterine fundal height above symphysis pubis have been developed. Ojwang et al used the product of symphysiofundal height and abdominal girth measurement at various levels in centimetres above symphysis pubis in obtaining a fairly acceptable predictive value but with considerable variation from the mean. Dare et al simplified and used the product of symphysiofundal height (Mc Donald's measurement) and abdominal girth at the level of umbilicus measured in centimetres and result expressed in grams to estimate foetal weight in utero at term, and the estimation correlated well with birth weight (Dare et al. (1990)).

**Johnson's formula for estimation of foetal weight in vertex presentation is as follows:**

**Foetal weight (grams) = (Mc Donald's measurement of symphysiofundal height in cm - X) x 155** where X = 13, when presenting part was not engaged, X = 12 when presenting part is at 0 station and X = 11 when presenting part was at +1 station. If a patient weighs more than 91 kg, 1cm is subtracted from the fundal height.


**Dawn's formula states that weight (grams) = longitudinal diameter of the uterus x transverse diameter of the uterus x 1.44/2.**

Measurements are made with pelvimeter. Double abdominal wall thickness was also measured pelvimeter. If Double abdominal wall thickness was more than 3 cm, the excess was deducted from the longitudinal diameter.


## 2. Material and Methods

The data from 232 pregnant Sudanese women were evaluated retrospectively during this study in period from January 2016-Till January 2017. Ultrasonography evaluation was conducted by Dr. Elsir Ali Saeed and prof. Dr. Mohamed Elfadil. Women with multiple gestations, diabetes, or growth disorders such as intrauterine growth retardation were excluded. Cross-sectional measurements of each case were used for estimation of fetal weight using ultrasonographic measurement. Biparietal diameter (BPD) was measured in axial plane at the level where the continuous midline echo is broken by the septum pellucidum cavum. Measurements were made from the outer to inner margin of the fetal skull. The abdominal circumference (AC) was measured directly by a plot on transfers section through the fetus abdomen at the level where the umbilical vein and stomach bubble were seen. The femur was measured from the origin to the distal end of the shaft. Then the correlation was developed to develop a linear and logarithmic equation to estimate the fetal weight as follow:


## 3. Result Presentation




**Figure 1:** scatter diagram showed the relation between average value of abdominal circumference (AC) (mm) (x) and FW (y)




**Figure 2:** scatter diagram showed the relation between abdominal circumference (AC) (x) and FW (y)




**Figure 3:** scatter diagram showed the relation between average value of femur length (FL) (x) and FW (y)



**Figure 4:** scatter diagram showed the relation between of femur length (FL) (x) and FW (y)



**Figure 5:** scatter diagram showed the relation between average value of BPD (x) and FW (y)



**Figure 6:** scatter diagram showed the relation between BPD (x) and FW (y)

**Table 1:** statistical measure for study variables

Variables	Mean	Std. Deviation
BPD	72.4	16.5
FL	54.5	15
AC	237.6	66.2
FW	1526.4	1007.7

#### 4. Discussion

The purpose of this study was to develop a model for estimation of fetal weight using ultrasound measurement, in normal Sudanese pregnant women using ultrasound

measurement. The research includes 232 pregnant women. The variables used to establish this study was FA, BPD, fetal FL, and AC, which having mean± SD of (1526.4±1007.7gm), (72.4±16.5mm), and (54.5±15mm), (237.6±66.2mm) respectively. As in table (1).

Fetal weight estimation has become increasingly important especially for the prevention of prematurity, evaluation of fetopelvic disproportion, decision for mode of delivery in breech presentation, induction of labor before term, in complications of pregnancy and in detection of intrauterine growth retardation. A lot of work has been done to find out the accurate methods for estimation of fetal size and weight in utero. The different works include clinical, biochemical, radiographic and ultrasonographic methods. Clinical methods were criticized on the basis of being less accurate and subject to considerable observer variation. Biochemical methods were not found to be satisfactory. Radiography was abandoned because of its hazards to both fetus and mother. Ultrasonography has gained popularity for determination of fetal parameters and wellbeing and also found to be useful for estimation of fetal weight. (P< 0.05).

Predanic et al. (2002) reported a significant improvement in EFWs with more training of residents. Ben-Aroya et al. (2002) reported that residents' fatigue affected the accuracy of clinical, but not sonographic EFWs. This study demonstrated that O&G residents and MFM subspecialists achieved accuracies of 65% and 79%, respectively, for EFWs to within 10% of the actual birth weight for all births.

Average weeks of abdominal circumference (mm) strongly related to the fetal Weight in (Gm) where logarithmic correlation was investigate this relationship reveals that (fetal weight = (0.0459\*Average AC)<sup>2</sup> - (6.6796\* Average AC) + 325.01). This equation further can be used for estimation of fetal weight without any ultrasonographic formula correlation co-efficient equal to (R<sup>2</sup>=0.997).

Also abdominal circumference (mm) strongly related to the fetal Weight in (Gm) where logarithmic correlation was investigate this relationship reveals that (fetal weight = (0.0469\*AC (mm))<sup>2</sup> - (7.3876\*AC(mm)) + 431.3). This equation further can be used for estimation of fetal weight without any ultrasonographic formula correlation co-efficient equal to (R<sup>2</sup>=0.9736). We used the average in state of normal measurement because of some sort of variation that noted in scatter plot and the average date of GA using AC can be used for fetal weight estimation. As in figure (1) and (2).

Linear correlation was intended to assess the relationship between the fetal weight and average values of femur length after grouping the femur length data, so as in figure (3) strong logarithmic correlation was noted at R<sup>2</sup>= 0.9964 and the fetal weight=-(1.0041\*Average FL (mm))<sup>2</sup> - (42.21\*Average FL (mm)) + 620.43). Also when we use the measurement without average values the correlation co-efficient equal to R<sup>2</sup> = 0.9833, indeed these values crospnd the strong relationship between these variables. Where the equations stated that (y = 0.9966x<sup>2</sup> - 41.553x + 608.66)

Where Y= fetal weight and X= FL (mm). As in figure (4 and 3)

Indeed as we already show in other articles the BPD is one of the methods that can be used in assessing the value of ultrasound examination in 2nd and strongly in 3rd trimester in case of fetal weight and GA estimation using ultrasound formulas here was a strong relationship between the Fetal weight and BPD in which direct (logarithmic) relation was reveals that ( $y = 1.2931x^2 - 121.62x + 3202.8$ ) at  $R^2 = 0.9816$ , for BPD in (mm). And ( $y = 1.2949x^2 - 121.81x + 3207.7$ ), at  $R^2 = 0.9934$ . For average value in (mm). As in figure (5) and (6).

## 5. Conclusion

The model can be summarized as:

$$FW (gm.) = 0.0459 * (average AC (mm))^2 - 6.6796 * average AC (mm) + 325.01.$$

$$FW (gm.) = 0.0469 * AC (mm)^2 - 7.3876 * AC (mm) + 431.3.$$

$$FW (gm.) = 1.0041 * (Average FL (mm))^2 - (42.21 * Average FL (mm)) + 620.43.$$

$$FW (gm.) = 0.9966 * (FL (mm))^2 - 41.553 * FL (mm) + 608.66$$

$$FW (gm.) = 1.2949 * (average BPD (mm))^2 - 121.81 * average BPD (mm) + 3207.7$$

$$FW (gm.) = 1.2931 * (BPD (mm))^2 - 121.62 * BPD (mm) + 3202.8.$$

## References

- [1] Kalish RB, Chervenak FA. Sonographic determination of gestational age. *Ultrasound Rev Obstet Gynecol* 2005; 5: 254-8.
- [2] Sowjanya, S. Lavanya Comparative Study of Clinical Assessment of Fetal Weight Estimation Using Johnson's Formula and Ultrasonographic Assessment Using Hadlock's Formula at or near Term, IOSR (Journal of Dental and Medical Sciences (IOSR-JDMS)), Volume 14, Issue 4 Ver. VII (Apr. 2015), PP 20-23.
- [3] Watson WJ, Soisson AP, Harlass FE. Estimated fetal weight of the term fetus. Accuracy of ultrasound vs. clinical examination. *J Reprod Med* 1988; 33:369-71.
- [4] Predanic M, Cho A, Ingrid F, et al. Ultrasonographic estimation of fetal weight: acquiring accuracy in residency. *J Ultrasound Med* 2002; 21:495-500.
- [5] Ben-Aroya Z, Segal D, Hadar A, et al. Effect of Ob/GYN residents' fatigue and training level on the accuracy of weight estimation. *Fetal Diagn Ther* 2002; 17:177-81.
- [6] Sherman DJ, Arieli S, Tovbin J, Siegal G, Caspi E, Bukovsky A. A comparison of clinical and ultrasound estimation of fetal weight. *Obstet Gynaecol* 1998; 91:212-7.
- [7] Chauhan SP, Hendrix NW, Magann EF, Morrison JC, Jenney SP, Devoe LD. Limitations of clinical and sonographic estimation of birth weight: experience with 1034 parturients. *Obstet Gynecol* 1998;91:72-7.
- [8] Nzeh DA, Oyawoye O, Adetoro OO. Ultrasound estimation of birth weight in late pregnancy among African women. *West African J ultrasound* 2000; 1:9-14.
- [9] Hanretty KP, Neilson JP, Fleming EE. Re-evaluation of clinical estimation of fetal weight: a comparison with ultrasound. *J Obstet Gynaecol* 1990; 10:199-201.
- [10] Hendrix NW, Grady CS, Chauhan SP. Clinical versus sonographic estimates of birth weight in term parturients. A randomized clinical trial. *J Reprod Med* 2000; 45:317-22.
- [11] Raman S, Urquhart R, Yusof M. Clinical versus ultrasound estimation of fetal weight. *Aust N Z J Obstet Gynaecol* 1992; 32:196-9.
- [12] Watson WJ, Soissons AP, Harlass FE. Estimated weight of the term fetus. Accuracy of ultrasound vs. clinical examination. *J Reprod Med* 1988; 33:369-711.
- [13] Dawn CS, Modale GC, Galosh A. A simple procedure for determination of antenatal fetal weight. *J Obstet gynecol* 1983; 33:133-7.
- [14] Tiwari R, Sood M. Comparative study of various methods of fetal weight estimation at term pregnancy. *J Obstet Gynaecol* 1989; 39:279-86.
- [15] Nahum G. Estimation of fetal weight-a review article last updated on 11 July 2002 (<http://www.emedicine.com>)
- [16] Mhaskar R, Mhaskar A, Molly SR, Symphysiofundal height (SFH) measurement for prediction of birth weight-A new formula. *J obstet Gynaecol Ind* 2001; 51:73-6
- [17] Mehdizadeh A, Alaghebandan R, Horsan H. Comparison of clinical versus ultrasound estimation of fetal weight. *Am J Perinatol* 2000; 17:233-6.
- [18] Wilcox AJ, Skjaerven R. Birth weight and perinatal mortality: the effect of gestational age. *Am J public health* 1992; 82:378-82.
- [19] Cecatti JG, Machado MR, Dos santos FF, Marussi EF. [curve of normal fetal weight values estimated by ultrasound according to gestational age]. *Cad Saude Publica Brazil* 2000; 16:1083-90.
- [20] Johar R, Rayburn W, Weir D, Eggert I. Birth weight in term infants. A 50-year prospective. *J Reprod Med* 1988; 33:813-6.
- [21] Hulsey TC, Levkoff AH, Alexander GR. Birth weights of infants of black and white mothers without pregnancy complications. *Am J Obstet gynecol* 1991; 164 (pt 1):1299-302.
- [22] Richards M, Hardy R, Kuh D, Wadsworth ME. Birth weight and cognitive function in british 1946 cohort. Longitudinal population based study. *BMJ* 2001; 322:199-203.
- [23] Hadlock FP, Harrist RB, Sharman RS, Deter RL, Park SK. Estimation of fetal weight with the use of head, body and femur measurements-a prospective study. *Am J Obstet gynaecol* 1985; 151:33-7?
- [24] Ebomoyi E, Adetoro OO, Wickremasinghe AR. Birth weight and sociological factors in lorin, Nigeria. *J Bio Soc* 1991; 23:417-23.
- [25] Ojwang S, Ouko BC. Prediction of fetal weights in utero by fundal height/girth measurements. *J Obstet Gynecol East Central Afr* 1984; 3:111.
- [26] Dare FO, Ademowore AS, Ifatureti OO, Nganwuchu A. The value of symphysiofundal height/abdominal girth measurement in predicting fetal weight. *Int J gynaecol Obstet* 1990; 31:243-8.
- [27] Schild RL, Fimmers R, Hansmann M. Fetal weight estimation by three-dimensional ultrasound. *Ultrasound Obstet Gynaecol (England)* 2000; 16:445-52.

- [28] Benacerraf BR, Gelman R, Frigoletto FD, Jr. Sonographically estimated fetal weight: accuracy and limitation. *Am J Obstet Gynecol* 1988; 159:1118-21.
- [29] Chauhan SP, Lutton PM, Bailey KJ, Guerrieri JP, Morrison JC. Intrapartum clinical, sonographic, and parous patients' estimates of newborn birth weight. *Obstet Gynecol* 1992; 79:956-8.
- [30] Nzeb DA, Rimmam S, More WMO, Hunt I. Prediction of birth weight by fetal ultrasound biometry. *Br J Radiol* 1992; 66:987-9.
- [31] Ratanasiri T, Jirapornkul S, Sombooporn W, Seejorn K, Patumnakul P. Comparison of the accuracy of ultrasonic fetal weight estimation by using the various equations. *J Med Assoc Thai* 2002; 85:962-7.
- [32] Bossak WS, Spellacy WN. Accuracy of estimating fetal weight by abdominal palpation. *J Reprod Med* 1972; 9:58-60.
- [33] Lawn J. E, Cousens S and Zupan J. Four Million neonatal deaths: When? Where? Why? *Lancet* 2005; 365:891-900
- [34] Banin JD, Gussman D, Stone P. Clinical and patient estimation of fetal weight vs. ultrasound estimation. *J Reprod Med* 2002; 47:194-8.
- [35] Nahum GG, Stanislaw H, Huffakar BJ. Accurate prediction term birth-weight from prospectively measurable maternal characteristics. *J Reprod Med* 1999; 44:705-12.
- [36] Pernoll ML, Taylor CM. Normal pregnancy and perinatal care. In: DeCherney AH, Pernoll ML, editors. *Current obstetrics and gynaecologic diagnosis & treatment*. 8th ed. Connecticut: Appleton and Lange, 1994: 183-7.
- [37] Abramowicz JS, Sherer DM, Bar-Tov E, Woods JR, jr. The cheek-to-cheek diameter in ultrasonographic assessment of fetal growth. *Am J Obstet Gynaecol* 1991; 165 (pt 1):846-52.
- [38] Abramson JH, Gahlinger M. Computer Programme for Epidemiologist (PEPI) version 3.10. Ilanidloes: Brixton Books, 1999.
- [39] Titapant V, Chawanpaiboon S, Mingmitpatanakul K. A comparison of clinical and ultrasound estimation of fetal weight. *J Med Assoc Thai* 2001; 84: 1251-7.
- [40] Shamley KT, Landon MB. Accuracy and modifying factors for ultrasonographic determination of fetal weight at term. *Obstet Gynecol* 1994; 88: 926-30.
- [41] Johnstone FD, Prescott RJ, Steel JM, Mao JH, Chambers S, Muir N. Clinical and ultrasound prediction of macrosomia in diabetic pregnancy. *Br J Obstet Gynaecol* 1996; 103:747-54.
- [42] Campbells, Wilkin D. ultrasonic measurement of foetal abdominal circumference in the estimation of fetal weight. *Br J Obstet Gynaecol* 1975; 82:689-97.
- [43] Combs CA, Jaekle RK, Rosenn B, Pope M, Miodovnik M, Siddiqi TA. Sonographic estimation of fetal weight based on a model of fetal volume. *Obstet Gynecol* 1993; 82:365-70.
- [44] Ott WJ, Doyle S, Flamm S, Wittman J. accurate ultrasonic estimation of fetal weight. Prospective analysis of new ultrasonic formulas. *A m J Perinatol* 1986; 3: 307-10.
- [45] Warsof SL, Wolf P, Coulehan J, Queenan JT. Comparison of fetal weight estimation formulas with and without head measurements. *Obstet Gynecol* 1986;67:569-73
- [46] Shepard MJ, Richards VA, Berkowitz RL, Warsof SL, Hobbins JC. An evaluation of two equations for predicting fetal weight by ultrasound. *Am J Obstet Gynecol* 1982; 142:47-54?
- [47] Deter RL, Hadlock FP. Use of ultrasound in the detection of macrosomia: a review. *J Clin Ultrasound* 1985;13:519
- [48] Richards M, Hardy R, Kuh D, Wadsworth ME. Birth weight and cognitive function in the British 1946 birth cohort: longitudinal population-based study. *BMJ* 2001; 322:199-203.
- [49] Susuki K, Minei LJ, Schintzer LE. Ultrasonographic measurement of foetal heart volume for estimation of birth weight. *Obstet Gynecol* 1974; 43:867-71.
- [50] Poulos PP, Langstadt JR. The volume of the uterus during labor and its correlation with birth weight. I. A method for the prediction of birth weight. *Am J Obstet Gynecol* 1953; 65:233-44.
- [51] Wikstrom I, Bergstrom R, Bakketaig L. Prediction of high birth weight from internal characteristics, symphysis-fundal health and ultrasound biometry. *Gynaecol Obstet Invest* 1993;35: 27-33.
- [52] Williams Obstetrics 21st edition. Chapter 29. Fetal growth disorders Pg 743-760