

Association between Gram-Negative Enteric Rods, *Porphyromonas gingivalis* and Changes in Clinical Parameters in Chronic Periodontitis: An Observational Study

Bhavya .B¹, Ashwini .S², Shruthi .K .R³

Department of Periodontology, Faculty of dental Sciences, Ramaiah University of Applied Sciences, Bangalore-560054

Abstract: *The correlation between Gram negative enteric rods and Porphyromonas gingivalis in periodontal diseases has received little attention in the literature. The objective of this study was to investigate the association between gram negative enteric rods and Porphyromonas gingivalis and changes in clinical parameters in chronic periodontitis subjects. Occurrence of Gram negative enteric rods, Porphyromonas gingivalis and clinical parameters were examined in 30 subjects with chronic periodontitis. Sub gingival plaque samples were collected from the deepest periodontal pocket and transported in thyoglycolate broth and organisms were cultured. Chi-square, Mann-Whitney and Spearman rank correlation coefficient were used to assess the clinical data. Gram negative enteric rods, Porphyromonas gingivalis were detected in 20 subjects. There were significantly positive correlations between Gram negative enteric rods, Porphyromonas gingivalis and bleeding on probing, probing depth, clinical attachment loss. This study suggests that presence of enteric rods and Porphyromonas gingivalis were related to adverse periodontal conditions. These results could have an impact on periodontal treatment and should be taken into account in the mechanical and antimicrobial treatment of periodontal disease in some populations.*

Keywords: Gram negative rods, Porphyromonas gingivalis, chronic periodontitis

1. Introduction

Periodontitis, a biofilm-related infection with mixed microbial etiology. Sub gingival biofilm hosts a variety of bacterial species and only a few have been associated positively with disease progression. Porphyromonas gingivalis is present in 85% of the diseased sites in chronic periodontitis. (1) They can adhere and rapidly invade oral epithelial cells (2). Its fimbriae mediate initial attachment and subsequent invasion (3). Gram negative enteric rods have shown the capacity to invade human tissue and produce enterotoxins. Hence the aim of the study was to investigate the association between Gram negative enteric rods, Porphyromonas gingivalis and clinical parameters in subjects with chronic periodontitis.

2. Methodology

A total of 30 subjects reported to the Department of Periodontology, Faculty of Dental Sciences, Ramaiah University of Applied Sciences in Bangalore, who were diagnosed with chronic periodontitis were included. Informed and written consents were obtained from each participant. The study design was approved by the Ethical

Committee and study is carried out for one month i.e., from December 2015 – January 2016.


Inclusion criteria's

- Age group 25 - 65 years
- Pocket depth \geq 4mm,
- Bleeding on probing,
- Clinical attachment level \geq 5mm.

Exclusion criteria's

- Subjects with history of any systemic diseases.
- Subjects who smoke.
- With history of periodontal treatment in previous 6 months.
- Who are pregnant or lactating
- Who used antibiotic or other drugs that affect periodontal status in the past 6 months.
- Regularly using CHX mouthwash.

Clinical Evaluation



Bleeding on probing elicited using a straight probe

Probing pocket depth elicited using a UNC-15 probe.

Microbial sampling and isolation of *Porphyromonas gingivalis* by culture

Microbial sampling on periodontitis patients was performed on pockets ≥ 4 mm. The deepest pockets were selected for sampling. After removing supragingival plaque with curettes and isolating the area with cotton pellets, the absorbent paper points were inserted into each periodontal pocket for 20 seconds. The paper points were transferred to a tube with thyoglycolate medium. All samples were labeled properly and processed within four hours after sampling. The samples were analyzed using microbial culture techniques for the

presence of periodontopathic bacteria, most samples were processed at room temperature (25°C) and incubated in CO₂ and anaerobic culture systems. The Trypticase Soy Serum Bacitracin Vancomycin agar medium was incubated in 10% CO₂ at 37°C for four days. The colonies are seen on culture plates which were seen as black pigmented colonies after gram staining. Total viable counts (TVC) were defined as the total number of colony forming units obtained on non selective media plates. Species found on selective media were enumerated and their percentage of TVC was calculated.



Collecting sub-gingival plaque using absorbent paper point.

Collected sample is transferred into a transport medium, (thyoglycolate)




TSBV Media is poured on plates


Transferring sample on to medium



Incubating sample in 10% CO₂ at 37°C for 4days



Colonies formed after incubation



After incubation inoculate micro organisms on non-selective media

Growth of different colonies on non-selective media

Growth of black-pigmented colonies on non selective blood agar

Isolation of Gramnegative enteric rods by culture:

After placement for 20 s, the paper points were pooled into a vial containing thymoglycolate transport medium. The sample vials were maintained at room temperature, transferred to the laboratory, and processed within 4 h after sampling. After the vials were placed in an incubator for 30 min at 37°C, bacterial plaque was mechanically dispersed with a test tube mixer at the maximal setting for 60 s. Serial 10-fold dilutions were prepared in

pepton water, and aliquots were plated on MacConkey agar. The plates were incubated aerobically at 37°C for 24 h. Each isolate was characterized according to colonial and cellular morphology and Gram stain characteristics. Gram-negative enteric rods were speciated using a standardized biochemical test. Total viable counts were defined as the total number of colony forming units obtained on non-selective media plates. Species found on selective media were enumerated and presented as counts $\times 10^5$.




Macconkey agar plate

Streaking the plate

Processing the sample with in 4hrs

Incubated at 37°C for 24hours



Colonies formed on mac conkey agar plates

Colony forming units are counted

3. Statistical Analysis

Data were entered into an Excel, The database was subsequently locked, imported into Statistical Package for Social Sciences (SPSS) for Windows, formatted, and analyzed. Indicators of descriptive statistics were used, such as frequencies, percentage, average, variance, and standard deviation. The presence of *Porphyromonas gingivalis* and Gram-negative enteric rods positive individuals were described as the percentage of individuals with at least infected pocket. PD and CAL differences and the presence or absence of *Porphyromonas gingivalis* and Gram-negative enteric rods were determined by the Mann-Whitney test. Association among *Porphyromonas gingivalis* and Gram-negative enteric rods was expressed through a nonparametric correlation coefficient (Spearman rank). Only


sites presenting concomitantly CAL and PD of 4 mm or more at baseline were considered in the analyses of CAL, PD, and BOP. The significance level was set at 0.05 for all tests.

4. Results

Among 30 patients examined, Gram-negative enteric rods and *Porphyromonas gingivalis* were detected in 20 individuals, respectively. A total of 20 (66.66%) patients harbored both microorganisms studied. Gram-negative enteric rods in periodontal pockets was highly significant and positively correlated with presence of *P.gingivalis* ($r=0.652$, $P<0.0001$) and also both organisms were highly significant and positively correlated with PD, CAL [Table 1 and fig 1].

Table 1: Comparison of presence and absence of *Porphyromonas gingivalis* and Gram-negative enteric rods with clinical parameters by Mann-Whitney U test

Variables	Absence of p.gingivalis and gram-ve enteric rods			Presence of p.gingivalis and gram -ve enteric rods			U-value	Z-value	P-value
	Mean	SD	Sum of ranks	Mean	SD	Sum of ranks			
BOP	1.05	0.09	241.50	1.03	0.13	223.50	103.50	-0.3733	0.7089
PD	5.53	0.99	145.50	7.87	1.73	319.50	25.50	-3.6086	0.0003*
CAL	6.27	1.22	145.50	7.40	1.35	284.00	61.00	-2.1361	0.0327*



Porphyromonas gingivalis is highly and significantly correlated with PD when compared to CAL and BOP (Table 2 and figure 2)

Table 2: Comparison of presence and absence of *Porphyromonas gingivalis* with clinical parameters by Mann-Whitney U test

Variables	Absence of p.gingivalis			Presence of p.gingivalis			U-value	Z-value	P-value
	Mean	SD	Sum of ranks	Mean	SD	Sum of ranks			
BOP	1.07	0.10	180.00	1.03	0.11	285.00	75.00	-1.0999	0.2714
PD	5.10	0.74	68.50	7.50	1.67	396.50	13.50	-3.8055	0.0001*
CAL	6.20	1.40	68.50	7.15	1.31	351.00	59.00	-1.8038	0.0713




Table 3: Comparison of presence and absence of gram negative enteric rods with clinical parameters by Mann-Whitney U test

Variables	Absence of gram negative enteric rods			Presence of gram negative enteric rods			U-value	Z-value	P-value
	Mean	SD	Sum of ranks	Mean	SD	Sum of ranks			
BOP	1.06	0.07	169.50	1.03	0.12	295.50	85.50	-0.6379	0.5235
PD	5.80	1.03	115.50	7.15	1.98	349.50	60.50	-1.7378	0.0823
CAL	6.70	1.25	115.50	6.90	1.48	315.00	95.00	-0.2200	0.8259




Table 4: Correlation between CFU counts of *p. gingivalis* and gram -ve enteric rods with clinical parameters by Spearman's rank correlation

Clinical parameters	N	Spearman R	t-value	p-level
BOP	30	-0.3325	-1.8657	0.0726
PD	30	0.7820	6.6385	0.0001*
CAL	30	0.2529	1.3834	0.1775





Table 5: Correlation between CFU counts of *Porphyromonas gingivalis* with gram negative enteric rods by Spearman's rank correlation

Variables	Correlation between CFU counts of <i>p. gingivalis</i> with Gram negative enteric rods			
	N	Spearman R	t-value	p-level
	30	0.5297	3.3046	0.0026*



5. Discussion

In this study, we investigated the relationships between *P.gingivalis* and Gram negative enteric rods and clinical parameters from patients with untreated chronic periodontitis. Information from the present study may have therapeutic implications for the treatment of non oral infections caused by oral pathogens. Dissemination of periodontal pathogens to other body sites frequently occurs and may cause serious diseases.

The ecology of oral sub gingival communities in health and periodontitis and elucidate the relationship between inflammation and the sub gingival micro biome. Supra-and sub gingival bacterial biofilm development assumes a crucial part in the improvement and movement of the infection, with gram-negative anaerobic rods and spirochetes commanding sub gingival polymicrobial biofilms in chronic periodontitis. The results of present study also showed a positive correlation of presence of gram negative enteric rods in sub gingival plaque samples (4)

Carlos M et.al in 2012 found that there were significantly positive correlations between enteric rods and presence of *P. gingivalis* and both microorganisms were significantly and positively correlated with clinical parameters. The results of the present study shows a significant positive correlation between *Porphyromonas gingivalis*, gram negative enteric rods and clinical parameters.(5)

Carlos M et.al in 2011 showed that the mean probing depth (mm) of the sampled sites was significantly deeper in patients with presence of *P. gingivalis* and Gram-negative enteric rods. The present study also shows that the mean probing depth of the sampled sites were significantly deeper

in patients with presence of *P.gingivalis* when compared to Gram negative enteric rods.(6)

Lafaurie et.al in 2007 showed that *P. gingivalis* occurred in 71.5% and enteric rods occurred in 34.5% of individuals with chronic periodontitis. The present study shows that both *P.gingivalis* and gram negative enteric rods occurred in 66.66% of individuals with chronic periodontitis subjects.(8)

In a study by Thomasae in 1997 showed negative association between gram-negative enteric rods and *A. Actinomyces comitans* related to the ecological interrelationships that occur among subgingival microorganisms inhabiting deep periodontal pockets in humans. The results of present study showed *P.gingivalis* are inhabited more in deepest pockets than gram negative enteric rods in chronic periodontitis patients.(7,9)

The investigation of the subgingival microbiota in a specific nation gets to be related to distinguish its conceivable effect on results after treatment. A larger examination would be more appropriate to study connections between Gram negative enteric rods and *P.gingivalis* further. Contrasts in host response, oral cleanliness, oral therapeutic administrations access, and microbial species may clear up these refinements in the clinical articulation of periodontitis in the population concentrated. More thorough examinations tending to the relationship amongst periodontitis and natural and inherited variables are required.(10)

6. Conclusion

This study concludes that presence of gram negative enteric rods and *P.gingivalis* were related to cause periodontal conditions. These results could have an impact on

periodontal treatment and should be taken into account in the mechanical and antimicrobial treatment of periodontal disease in some populations.

7. Future Scope

Further long term studies have to be carried out for establishment of specific treatment strategies in larger populations in different parts of other countries.

References

- [1] Hajishengallis, G., Liang, S., Payne, M.A., Hashim, A., Jotwani, R., Eskan, M.A., McIntosh, M.L., Alsam, A., Kirkwood, K.L., Lambris, J.D. and Darveau, R.P., 2011. Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota and complement. *Cell host & microbe*, 10(5), pp.497-506.
- [2] Belton, C.M., Izutsu, K.T., Goodwin, P.C., Park, Y. and Lamont, R.J., 1999. Fluorescence image analysis of the association between Porphyromonas gingivalis and gingival epithelial cells. *Cellular microbiology*, 1(3), pp.215-223.
- [3] Tsuda, K., Furuta, N., Inaba, H., Kawai, S., Hanada, K., Yoshimori, T. and Amano, A., 2008. Functional Analysis of ALPHA. 5. BETA. 1 Integrin and Lipid Rafts in Invasion of Epithelial Cells by Porphyromonas gingivalis using Fluorescent Beads Coated with Bacterial Membrane Vesicles. *Cell structure and function*, 33(1), pp.123-132.
- [4] Abusleme, L., Dupuy, A.K., Dutzan, N., Silva, N., Burleson, J.A., Strausbaugh, L.D., Gamonal, J. and Diaz, P.I., 2013. The subgingival microbiome in health and periodontitis and its relationship with community biomass and inflammation. *The ISME journal*, 7(5), pp.1016-1025.
- [5] Carlos M. Ardila et.al (2012), *Journal of Indian Society of Periodontology*, Relationship between Gram negative enteric rods, Aggregatibacter actinomycetemcomitans, and clinical parameters in periodontal diseases, volume 16, Issue 1, page 65-69.
- [6] Carlos M. Ardila et.al (2011), *Acta Odontologica Latinoamericana*, Positive correlations between presence of gram negative enteric rods and porphyromonas gingivalis in sub gingival plaque, volume 24, page 15-19.
- [7] Herrera D et.al (2008) *Journal of clinical periodontology*, Subgingival microbial profiles in chronic periodontitis patients from Chile, Colombia and Spain, volume 35, Issue 10, page 106-113.
- [8] Gloria Ine's Lafaurie et.al (2007) *Journal of periodontology*, Demographic, Clinical, and Microbial Aspects of Chronic and Aggressive Periodontitis in Colombia: A Multicenter Study, volume 78, number 4, page 629-639.
- [9] Thomase. Rams et.al (1997) *clinical infectious diseases*, oxford university press, subgingival microbial associations in severe human periodontitis, volume 25, supplement 2, page 224-226.
- [10] Mager, D.L., Haffajee, A.D. and Socransky, S.S., 2003. Effects of periodontitis and smoking on the microbiota of oral mucous membranes and saliva in systemically

healthy subjects. *Journal of clinical periodontology*, 30(12), pp.1031-1037.

Author Profile



Corresponding author:-Dr. Bhavya. B is Reader, Department of Periodontology, Faculty of Dental Sciences, Ramaiah University of Applied Sciences, Bangalore-560054, India



Dr. Ashwini. S is Professor and Head of the Department, Department of Periodontology, Faculty of Dental Sciences, Ramaiah University of Applied Sciences, Bangalore-560054



Dr. Shruthi. K. R is Post graduate Student, Department of Periodontology, Faculty of Dental Sciences, Ramaiah University of Applied Sciences, Bangalore-560054