

Study of Some Parameters on the Rate of the Catalyzed Decomposition of Hydrogen Peroxide by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex

Magdy A. Wassel¹, Abdou S. El-Tabl², Ahmed S. Elzaref³, Mahmoud M. Arafa⁴, Abdullah A. Alhalib⁵

^{1,3}Chemistry Department, Faculty of Science, Al-Azhar University, Cairo, Nasr city, Egypt

^{2,4,5}Chemistry Department, Faculty of Science, Menoufia University, Shebin El-Kom, Egypt

Abstract: Catalytic decomposition of H_2O_2 by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex was studying under different experimental conditions such as different concentration of H_2O_2 (10%, 15% and 20%), different temperature (25, 35 and 45°C) and different amounts of complex as a catalyst (0.1, 0.3, 0.5 and 0.7 gm). It was observed from the experimental results that the catalytic decomposition rate constant was found to increase by increasing the amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst. This can be explained by higher hydroxyl radicals are formed in the reaction medium, and also increase in temperature accelerates the Kinetics of catalytic decomposition of H_2O_2 . Also the thermodynamic parameters: (E_a), (ΔH), (ΔG) and (ΔS) were determined.

Keywords: Metal complexes, Decomposition, H_2O_2 , Kinetics, Mechanism

1. Introduction

Hydrogen peroxide (H_2O_2) is involved in all life's vital processes as it is versatile in its uses and applications. Its varied utility ranges from fenton's reagent, bleaching, catalytic activity oxidant for industrial chemical effluents, biological and medical applications. [1]

The kinetics of H_2O_2 Catalytic decomposition over [Cu(II) fulvate] complex as a catalyst was investigated. The effect of different concentration of H_2O_2 , different amount of catalyst and different temperature were studied Based on the rate constants obtained at different temperatures; the empirical Arrhenius expression of H_2O_2 decomposition was derived. The reaction was further studied and the values of (E), (ΔH), (ΔG) and (ΔS) were determined, its well described by an autocatalytic first order equation. [2]

The catalytic decomposition of H_2O_2 in presence of $Fe_2O_3-MoO_3$ (2:4) departs from first order kinetics in the sense of an autocatalytic. In the higher stage of H_2O_2 decomposition the first order kinetics equation is obeyed. [3], [4]

H_2O_2 is widely recommended in aero-space field for propulsive applications for the high energy density and for its "green" nature due to low toxicity and low environmental impact in comparison with conventional propellants (N_2H_4 , N_2O_4). Nevertheless, the presence of a catalyst is necessary for H_2O_2 decomposition due to the low kinetics of the homogeneous reaction. The catalytic activity of innovative materials for the H_2O_2 decomposition has been studied under the vapor phase condition at 200°C, catalysts are based on (MnO_x) dispersed on monolithic zirconia substrates. They have been developed using a precipitation technique with an improved procedure, content of the active phase was from 0.5 to 2.0 wt %. Results obtained showed a high activity of the catalyst. The convention was strongly depending on the space velocity and on MnO_x content on the support. [5]

The catalytic and photocatalytic activities of ramsdellite type manganese oxide, R- MnO_2 were studied from the initial rate of decomposition of H_2O_2 in aqueous solution. The kinetic study was followed by the production of O_2 via gas pressure monitoring (O_2 – monitored method) in the dark and under visible light irradiation ($\lambda \geq 400$ nm) at 20 °C. Experimental data showed that the rate of H_2O_2 decomposition, under visible light irradiation and in the dark, obeys the first order kinetic law. While in the dark the H_2O_2 decomposition rate by mass unit was around $6.88 \text{ mol g}^{-1} \text{ s}^{-1}$, under visible light in addition the H_2O_2 decomposition rate achieved a maximal value of $15.13 \text{ mol g}^{-1} \text{ s}^{-1}$. The increase in the H_2O_2 decomposition by effect of the visible light was observed in range of catalyst concentration from 16 to 374 mg L^{-1} . These results reflect the potential of R- MnO_2 as photoactive catalyst, being a consequence of pressured photo education of Mn^{4+} centers. [6]

The decomposition rate of H_2O_2 by Fe(III)-nitrilotriacetate complex (Fe^{III} NTA) has been investigated over a large range of experimental conditions: $3 < \text{pH} < 7$, $[Fe(III)]_{T,0} : 0.05-1 \text{ mM}$; $[NTA]_{T,0} / [Fe(III)]_{T,0}$, molar ratios: 1-250; $[H_2O_2]_0 : 1 \text{ mM} - 4 \text{ M}$ and concentration of OH radical scavengers: 0-53 mM. Spectrophotometric analyses revealed that reactions of H_2O_2 with Fe^{III} NTA (1mM) at neutral pH immediately lead to the formation of intermediates (presumably peroxocomplexes of Fe^{III} NTA) which absorb light in the region 350-600 nm where Fe^{III} NTA and H_2O_2 were first order with respect to H_2O_2 and that the apparent first-order rate constants were found to be proportional to the total concentration of Fe^{III} NTA complexes.

Were at a maximum at $\text{pH} = 7.95 \pm 0.10$ and depend on the $[NTA]_{T,0} / [Fe(III)]_{T,0}$ molar ratios. The addition of increasing concentration of tert-butanol or sodium bicarbonate significantly decreased the decomposition rate of H_2O_2 , suggesting the involvement of HO^\cdot radicals in the decomposition of H_2O_2 . The decomposition of H_2O_2 by Fe^{III} NTA at neutral pH was accompanied by a production of

dioxygen and by oxidation of NTA. The degradation of the organic ligand during the course of the reaction led to a progressive dcomplexation of Fe^{III}NTA followed by a subsequent precip-itation of iron (III) oxyhydroxides and by a significant decrease in the catalytic activity of Fe(III) species for the decomposition of H₂O₂. [7]

2. Experimental

• Materials and methods

All chemical were of analytical grade and were employed without further purification. The experimental apparatus as in Fig(1). It contains : (1) sort ball neck (reactor) borosilicate glass 500 ml Volume. (2) A condenser to prevent evaporation of the reaction medium and maintaining constant the content of H₂O₂. (3) A heater thermally regulated flask to set the desired temperature (4) Thermometer . (5) A syringes to introduce the oxidant and take sample.

Working solutions of H₂O₂ were prepared weekly by volumetric dilution of 30% (v/v) H₂O₂ (AR, grade, Merck) and were standardized daily by titration with potassium permanganate KMnO₄ it was found that the natural decomposition rate of aqueous H₂O₂ Solution was less than 1% in 24h.

The decomposition of H₂O₂ catalyzed by [H₅LFe₂(II)(SO₄)₂.H₂O] complex as in Fig(2) can be kinetically monitored by removing aliquots of the reaction mixture at predeterm9;99999ined intervals and titrating the undecomposed H₂O₂with standard KMnO₄ solutions (0.04-0.05N) standardized with(COONa)₂ (primary standard). The decomposition of H₂O₂ was carried out in a thermostatic cell at different temperatures between 25 and 45 °C .




Figure 1: Experimental Apparatus




Figure 2

• Preparation of catalyst [H₅LFe₂(II)(SO₄)₂.H₂O] complex

A) Ligand preparation (H₅L)

Sodium salt added to hydrazine by different amount with stirring the suspension at 70 °C for one hour, The product is filtered and gives a start, starting with glucose with stirring at 60 °C for an hour and filtered product theLigand as Fig(3).




Figure 3

B) Preparation of metal complex (H₅L):-

Iron(II)sulphate tetrahydrate dissolved in ethanol warmed at 60 °C was added to H₅L in the same solvent by different amount (1 mol). The mixture was warmed at 60 °C with stirring for 1 hour, then the solution was filtered off and dark brown precipitate was obtained by evaporated the solution.Fig(2)

3. Results and Discussion

(A) Kinetics of the catalytic decomposition of H₂O₂:

Many experiment were carried out to study the effect of H₂O₂ concentration, effect of the amount of catalyst [H₅LFe₂(II)(SO₄)₂.H₂O]complex and the effect of different temperature (25-45 °C) on the rate of catalytic decomposition of H₂O₂. Although the decomposition of H₂O₂ is very complex, it has been shown that under certain operating conditions, the kinetics of decomposition of H₂O₂ can be described by a kinetic of pseudo-first order as described in the following the equation(1)

$$d[H_2O_2]/dt = -k_{app} [H_2O_2] \quad (1)$$

Where (k_{app}): the apparent rate constant of the first order, (t): reaction time and [H₂O₂]: the concentration of hydrogen

peroxide at time (t). After integration we obtain the equation(2)

$$\ln([H_2O_2]_o/[H_2O_2])=k_{app}t \quad (2)$$

where $[H_2O_2]_o$: initial concentration of hydrogen peroxide.[8]

Kinetic studies in the conditions mentioned above showed that the initial rate of the decomposition of H_2O_2 is proportional to the concentration of hydrogen peroxide and the amount of catalyst $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst and the proposed mechanism of this reaction may be as the following :

$$V_o = -d[H_2O_2]/dt = k_{dec} [H_5LFe_2(II)(SO_4)_2.H_2O] [H_2O_2] \quad (3)$$

4. Studies the different factors effecting in the Catalytic Decomposition of H_2O_2

1- Study the effect of temperature

The experiments were performed with 10 % H_2O_2 and 0.3 g $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex. Temperatures of our experimental are carried out between 25 to 45 C. As can be seen, the increase in temperature accelerates the kinetics of decomposition of hydrogen peroxide. Indeed, at 45 °C the decomposition is slow. The rate constants of H_2O_2 decomposition as a function of temperatures are given in (Table 1) and (Figure 4-11).

Table 1: Rate constant at different temperature

Temp.	Time (min)				k	Logk	1/T
	10	20	30	40			
	$\ln([H_2O_2]_{dec})$						
25 C°	2.166	2.194	2.200	2.240	0.0228	-2.638	0.00335
35 C°	2.180	2.200	2.220	2.230	0.0017	-2.769	0.00324
45 C°	2.240	2.270	2.280	2.287	0.0015	-2.823	0.00314

When we draw $(\ln([H_2O_2]_{dec}))$ versus time (t) at constant temperature 25,35 and 45° C as shown in Figure (4, 5 and 6) its clear straight line :




Figure 4: The rate constants of H_2O_2 decomposition temperature 25°C.




Figure 5: The rate constants of H_2O_2 decomposition temperature 35°C.




Figure 6: The rate constants of H_2O_2 decomposition temperature 45°C.




Figure 7: Plot of $\log(k)$ Vs $(1/T)$ for the decomposition of H_2O_2 by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex ; $[H_2O_2]_0 = 10\%$

The activation energies of the catalytic decomposition of H_2O_2 at temperature range 25-45C° are calculated from Arrhenius plots and the values obtained are given in Fig (7). It seems that the activation energy (E_a) evaluated from (k) values, (E_a) indicating that the decomposition of H_2O_2 , we see that Arrhenius equations:

$$\log k = -E_a/2.303RT + \log A \quad (10)$$

Given with slope equal to $(-E_a/2.303R)$ and intercept equal to $(\log A)$. This result are illustrated in Fig (7).

2- Study the Thermodynamic Parameters for the Catalytic Decomposition of H_2O_2

The thermodynamic parameters (E_a), (ΔH), (ΔG) and (ΔS) were also determined and illustrated in Table (2). The automatically first order plots derived from following equation:

$$\log k/T = \log k_B/h + \Delta S/R - \Delta H/RT \quad (11)$$

Here, (k) is the rate constant for the reaction, (T) is absolute temperature, (k_B) is Boltzmann's constant, (h) is Planck's constant, (ΔS) and (ΔH) are activation entropy and enthalpy, respectively. The slope of the line obtained using this equations proportional to the activation enthalpy, while the activation entropy can be extracted from the intercept. By performing each reaction at a series off our temperatures and using the empirical rate law to determine the rate constant at each temperature, activation entropy and enthalpy can be calculated. Are illustrated in Fig. (8).and equation (12):

$$\Delta G = \Delta H - T\Delta S \quad (12)$$

Table 2: Thermodynamic parameter for decomposition of hydrogen peroxide by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as a catalyst

Temp. (K°)	ΔG (KJ/mol)	ΔS (J/mol k)	ΔH (KJ/mol)	E_a (KJ/mol)
298	-14.54	15.74	19.23	16.96
308	-14.38			
318	-14.22			




Figure 8: Plot of $\log(k/T)$ vs $(1/T)$ for the decomposition of H_2O_2 by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex ; $[H_2O_2]_0 = 10\%$




Figure (9): Effect of temperature on the catalytic decomposition of H_2O_2 (at $[H_2O_2]_0 = 10\%$, various amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst).




Figure 10: Effect of temperature on the catalytic decomposition of H_2O_2 (at $[H_2O_2]_0 = 15\%$, various amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst)




Figure 11: Effect of temperature on the catalytic decomposition of H_2O_2 (at $[H_2O_2]_0 = 20\%$, various amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst)

The heterogeneous as catalytic decomposition of H_2O_2 over $Fe_2O_3-MoO_3(2:4)$ is characterized by the presence of an induction period during which the active centers (Mo^{5+}) are being formed at the start of reaction [9] in all cases studied an induction period has been encountered before the reaction assumes a significant rate. These results showed also that the induction period (t_{ind}) is largely by the amount of catalyst and the temperature of the reaction.

3- Study The effect of the amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst:

The level of concentrations of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex used in this study is (0.1, 0.3, 0.5 and 0.7 g) A significant increase in the rate of decomposition was observed by increasing the dose of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex. For high concentrations of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex, the kinetics of decomposition of H_2O_2 is fast. This can be explained by the fact that the higher the $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex, the higher OH^- hydroxyl radicals are formed in the reaction medium. The relationship between the amount of catalyst

and the catalytic decomposition of H_2O_2 at different temperature were illustrated in Fig (12-14).




Figure 12: Effect of the amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex on the catalytic decomposition of H_2O_2 (At various initial concentration of H_2O_2 , at $25^\circ C$).




Figure 13: Effect of the amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex on the catalytic decomposition of H_2O_2 (at various initial concentration of H_2O_2 , at $35^\circ C$)




Figure 14: Effect of the amount of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex on the catalytic decomposition of H_2O_2 (at various initial concentration of H_2O_2 , $45^\circ C$).

4- Study the Role of H_2O_2 Concentration

In this case, the concentration of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex is set equal to 0.1, 0.3, 0.5 and 0.7 g respectively, the temperature is varied between (25, 35, and $45^\circ C$) and the initial concentration of hydrogen peroxide is variable from (10, 15, and 20 %). The evolution of the H_2O_2 concentration is reported in Figure (15-18).

As the figure shows the decomposition rate increased by increasing H_2O_2 concentration. This may be due to the competition between the (OH^-) radicals and other compounds. For high concentrations of H_2O_2 the kinetics of decomposition of hydrogen peroxide is fast. This is explained by the Fenton reaction to high concentrations of H_2O_2 .




Figure 15: Effect of initial concentration of H_2O_2 on the catalytic decomposition of hydrogen peroxide by 0.1g $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as a catalyst)




Figure 16: Effect of initial concentration of H_2O_2 on the catalytic decomposition of hydrogen peroxide by 0.3g $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as a catalyst).




Figure 17: Effect of initial concentration of H_2O_2 on the catalytic decomposition of hydrogen peroxide by 0.5g $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as a catalyst)





Figure (18): Effect of initial concentration of H_2O_2 on the catalytic decomposition of hydrogen peroxide by 0.7g $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as a catalyst

The catalytic decomposition of H_2O_2 in presence of $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex as a catalyst, depends upon the concentration of the catalyst, temperature and slowed with high concentrations of H_2O_2 . The rate of the reaction is of pseudo first order with respect to the concentration of the catalyst and temperature.

(B) Mechanism of the Catalytic Decomposition of H_2O_2 :

We suggested the reaction mechanism of decomposition of H_2O_2 by $[H_5LFe_2(II)(SO_4)_2.H_2O]$ Complex as the following :



It is assumed that step (3) is the rate determining step of the reaction, as the reduction of Fe(II) requires geometrical changes around the metal center. This mechanism is in agreement with the suggestion for other macromolecular systems and can explain the change in rate in going to more alkaline conditions, as the dependence of the redox potential of the couple $HO_2^-(O_2^-)/H_2O_2$ (HO_2^-) on pH is very well known. It is obvious that the oxidation of H_2O_2 to HO_2^- or O_2^- by Fe(II) is more favorable in more alkaline conditions. This kind of mechanism does not involve any diffusible radical species in accordance to previous proposed mechanism, but several very rapid intermediate step between (3) and (4) may exist which include the participation of radicals.

Complexes of Fe(II) especially with various ligands acting as catalysts have been investigated in depth and disagreements over mechanistic details, involving intermediate radicals or complexes, have lasted for decades. The formation of iron peroxide complexes both in acidic and alkaline solutions has been confirmed and a mechanism not involving any radicals has been suggested. The difference in reactivity of Fe(II) complexes towards H_2O_2 is due to the change in the redox potential of Fe(II) ions as a result of ligation with different ligands. It was supposed that a superoxide iron (I) complex $[HO_2-Fe(I)(Ligand)]$ is formed from the complex $[HO_2-Fe(II)(Ligand)]$ in which an instantaneous electron transfer occurs.

The plot of $\log(dx/dt)/(a-x)^2$ against (t) , must be a straight line with slopes equal to $k/2.303$ and intercept equal to $-\log a/k_0$. The autocatalytic first order plots derived from the following equation (16) are illustrated in Fig.(4,5 and 6) and evaluated.

$$\log(dx/dt)/(a-x)^2 = k t / 2.303 - \log a/k_0 \quad (9)$$

The specific reaction rate (k) were evaluated from the plot of $(\ln[H_2O_2]_{dec})$ against time (t) as shown in , Table (1), where :

- $[H_2O_2]_{dec} = [H_2O_2]_0 - [H_2O_2]_{unreact}$
- $[H_2O_2]_{dec}$: concentration of H_2O_2 decom-position at time t
- $[H_2O_2]_0$: initial concentration of H_2O_2
- $[H_2O_2]_{unreact}$: concentration of remaining H_2O_2

5. Conclusion

The catalytic decomposition of H_2O_2 $[H_5LFe_2(II)(SO_4)_2.H_2O]$ complex depends upon the concentration of the catalyst temperature and slowed with high concentration of H_2O_2 .

The rate of the reaction is of pseudo first order with respect to the Fe(II) to Fe(I), during the slow rate determining step.

References

- [1] E.Linley,S.P.Danger, G.McDonnell,C.Simons and J.Mailard;“Use of hydrogen peroxide as a biocide: new consideration of its mechanisms of biocidal action”, J.of Anti. Chem 24, pp.1-8. 2012.
- [2] M. A. Wassel, A.S.El.Table, M.A.Wahba and M.M.Arafa ;“Kinetic Study of The Catalytic Decomposition of Hydrogen Peroxide Over [Cu(II)Fulvate] Complex As a Catalyst” ,J.of chem.and chem. Sci 5(11), pp. 629-638. 2015.
- [3] M.A.Wassel, F.I.zidan and M.El.Wardani.“Effect of the composition of $Fe_2O_3-MoO_3$ catalysts on the decomposition of H_2O_2 ”, J.seib. chem. Soc 54 (8), pp. 425-433.1989.
- [4] M.A.Wassel, F.I.zidan and M.El.Wardani,“decomposition of H_2O_2 on $Fe_2O_3-MoO_3$ (2:4)and the effect of the amount of catalys and temperature on the induction period”, Rasseгна chimica N.4.Luglio. Agosto.1990.
- [5] L.Micoli and M.Turco;“Decomposition of H_2O_2 on Monolithic $Mnox/zro_2$ Catalysts for Aerospace”, Application Chem.Eng.Tran 43, pp.1819-1824.2015.
- [6] H.E. Hamshary, M.H.El.Newehy and S.S.Al-Deyab;“Oxidation of phenol by hydrogen peroxide catalyzed by metal-containing poly (amidoxime) grafted starch”, Mole 16, pp. 9900-9911. 2011.
- [7] J.De laat, Y.H.Dao,N.H. Elnajjar and C.Daou;“Effect of some parameters on the rate of the catalyzed decomposition of hydrogen peroxide byiron(III)-nitrilotriacetate in water”, water research 45, pp. 5654 - 5664.2011.
- [8] O. Zeineb ,B. A. Hedi, M. R. Jeday,C. Cheker ;“Kinetic study of the catalytic decom-positionof H_2O_2 in phosphoric acid medium”, Int. J. o f. hydrogen energy 40, pp. 1278-1282.2015.
- [9] M. Fouda, M. Abd-Elzaher, M. Shakdofa, F. El Saied, M. Ayad, A. S. El Tabl ;“Synthesis and characterization of transition metal complexes of N' -[(1,5-dimethyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-4-yl)methylene]thiophene-2-carbohydrazide”, Tran. Met. Chem. 33, pp. 219-228. 2008.