CNS Stimulant Property of Natural Products – A Systemic Review

Srineeraja .P

Intern, Saveetha Dental College and Hospitals, Chennai, India

Abstract: The use of herbal medicines world wide has provided an excellent opportunity for India to look for therapeutic lead compounds from ancient systems of therapy.CNS stimulants include caffeine, cocaine, and various amphetamines. They are used to enhance mental alertness and reduce drowsiness and fatigue. This review aims to investigate the CNS stimulant activity of various natural productssuch as Ginkgo biloba, Bacopa Monnieri, Aswagandha, Acacia Catechu, Acacia Nilotica, Neem, Curcumin, Cucurbita Maxima, Barleria prionitis Linn.,Alpinia galanga and Diplazium Esculentum.

Keywords: CNS Stimulants, Ginkgo biloba, Bacopa Monnieri, Aswagandha, Acacia Catechu, Acacia Nilotica, Neem, Curcumin, Cucurbita Maxima, Barleria prionitis Linn.,Alpinia galanga and Diplazium Esculentum

1. Introduction

Over past few decades, the affinity towards the herbal drugs has been grown by utilization of traditional medicinal plant to heal some critical diseases and it is turning out to be better medicine with respect to synthetic drugs that assure numerous side effects for prolong treatment. . Herbal medicines are also in great demand in the developed world for primary health care because of their efficacy, safety and lesser side effects. They also offer therapeutics for agerelated disorders like memory loss, osteoporosis, immune disorders, etc. for which no modern medicine is available.

CNS disorders can affect either brain or the spinal cord resulting in neurological or psychiatric disorders. Causes of CNS diseases are trauma, infection, degeneration, autoimmune disorders, structural defects, tumours and stroke, neurodegenerative diseases, mood disorders, schizophrenia and autism [1]. Central nervous system (CNS) stimulants are medicines that speed up physical and mental processes. Central nervous system stimulants are used to treat conditions characterized by lack of adrenergic stimulation, including narcolepsy and neonatal apnoea. The majority of CNS stimulants is chemically similar to the neurohormone, norepinephrine and simulates the traditional "fight or flight" syndrome associated with sympathetic nervous system arousal.

1) BACOPA MONNIERI

Bacopa monnieri, also known as Water Hyssop (Brahmi), is prominently used in Ayurveda, a holistic system of medicine originating from India. It an adaptogen, a physiological agent that naturally increases the body's resistance to physical and emotional stress.[2]

Strengthens adrenal processes that facilitate carbohydrate metabolism. It is a powerful blood purifier and is specific for chronic skin diseases, including leprosy and syphilis, as well as eczema and psoriasis.

CNS STIMULANT ACTIVITY

It is the main revitalizing herb for the nerves and the brain cells. The herb is known for its ability to build and improve

mental performance. Improves both short-term and longterm memory. It increases intelligence, longevity, circulation in the brain; it decreases senility and ageing. It fortifies the immune system, both cleansing and feeding it.[3]

2) ASWAGANDHA

Ashwagandha (Withaniasomnifera, fam. Solanaceae) is commonly known as "Indian Winter cherry" or "Indian Ginseng". It is one of the most important herb of Ayurveda (the traditional system of medicine in India) used for millennia as a Rasayana for its wide ranging health benefits. It is used for various kinds of disease processes and specially as a nervine tonic.[4]

CNS STIMULANT ACTIVITY

It enhances the function of the brain and nervous system and improves the memory. It improves the function of the reproductive system promoting a healthy sexual and reproductive balance. Being a powerful adaptogen, it enhances the body's resilience to stress. Ashwagandha improves the body's defense against disease by improving the cell-mediated immunity. It also possesses potent antioxidant properties that help protect against cellular damage caused by free radicals.[5]

3) ACACIA CATECHU

The Acacia catechu is known as Cutch tree, Terra Japonica as well as Black Catechu. In Hindi it is called Khair and Khadira in Sanskrit.

This herb was previously known as Kat or Cacho. This tree was a very important export product that was sent from India to China, Persia and Arabia in the early 16th century particularly. This plant was used majorly for the purpose of dyeing and tanning.[6]

CNS STIMULANT ACTIVITY

The extract of heartwood, flowering tops, young shoots, the bark, fruits and the gum of the plant are used to create products for use. These extracts are used as an anodyne, bactericide, refrigerant, detergent, astringent, styptic, masticatory, expectorant, stimulant and as an antiphlogistic. The plant of Acacia catechu contains tannins and Flavonoids

Volume 6 Issue 7, July 2017 www.ijsr.net Licensed Under Creative Commons Attribution CC BY majorly.Catechin and rutin are most important constituents which are free radical scavengers.[7][8]

4) ACACIA NILOTICA

It is a medium sized tree characterized by dark brown or black and longitudinally fissured bark. Leaves bipinnate, petiole 2.5-5 cm long, pinnae in 4-9 pairs, stipular spines, white, sharp and straight. The decoction of bark yields spongy gum which is useful in sore throat, for washing ulcers, to stop bleeding from wounds, skin diseases, as an astringent for diarrhoea and leucorrhoea. Powdered gum is also given in dysentery and diabetes. The fried gum is considered a nutritive tonic, particularly in sexual debility. It also soothes inflamed membranes of the pharynx, alimentary canal and genito-urinary organs. It is prescribed for chest complaints. Pods are used to stop bleeding from bites of leeches. Tender shoots tops are used to treat cough.[9]

CNS STIMULANT ACTIVITY

It has Alexeteric, Algicide, Amebicide, Antibacterial, Antiedemic, Antihistaminic, Antinflammatory, Antibacterial, Aphrodisiac, Astringent, Decongestant, Demulcent, Expectorant, Hemostat, Hepatotonic, Hypoglycemic , Hypotensive, Lactagogue, Molluscicide, Neurostimulant, Protisticide, Stimulant, Taenicide, Teratologic , Vermifugeactivity.[10]

5) NEEM

The beneficial properties of Neem (Azadirachtaindica A. Juss) have been recognized in the Indian tradition. Each part of the neem tree has some medicinal property. Biswas et al (2002) have recently reviewed the biological activities some of the neem compounds, pharmacological actions of the neem extracts, clinical study and plausible medicinal applications of neem along with their safety evaluation. It is Antiarthritic; Anti-inflammatory; Antipyretic; an Hypoglycaemic; Antigastric ulcer; Spermicidal; Antifungal; Antibacterial; Diuretic; Antimalarial; Antitumour; Immunomodulatory etc.[11][12]

CNS STIMULANT ACTIVITY

It contains isoprenoids (like diterpenoids and triterpenoids containing protomeliacins, limonoids, azadirone and its derivatives, gedunin and its derivatives, vilasinin type of compounds and C- secomeliacins such as nimbin, salanin and azadirachtin) and non-isoprenoids, which are proteins (amino acids) and carbohydrates (polysaccharides), sulphurous compounds, polyphenolics such as flavonoids and their glycosides, dihydrochalcone, coumarin and tannins, aliphatic compounds, etc. Varying degrees of central nervous system (CNS) depressant activity in mice was observed with the leaf extract. Fractions of acetone extract of leaf showed significant CNS depressant activity.[13]

6) CURCUMIN

Curcumalonga, a perennial herb of the Zingibera-ceae (ginger family), contain curcuminoids and cur-cumin. Turmeric is used as a spice in curry powders and mustard. It is being investigated in clinical trials for the treatment and prevention of cancers, particularly of the GI tract, and for treatment of colitis and Alzheimer and Huntington diseases.[14][15]

CNS STIMULANT ACTIVITY

Rhizomes of Curcuma longa Linn.have been used to treat depression in past. Curcumin is low molecular weight polyphenol, is considered as active principle of Curcuma longa Linn. Curcuma longa Linn.is reported to have antioxidant, antiepileptic and neuroprotective properties.[16-18]

7) CUCURBITA MAXIMA

Cucurbita maxima is a short lived shrub which belongs to family Cucurbitaceae. Seeds are traditionally used as a bitter tonic, oil obtained used in debility nervous disorders.

CNS STIMULANT ACTIVITY

Cucurbita maxima has a potential CNS stimulant effect that can be explored for therapeutic advantage as an alternative treatment in medical conditions associated with dizziness and sedative.[19]

8) BARLERIA PRIONITIS LINN.

Barleria prionitis Linn. (Acanthaceae) is widely distributed throughout India, Sri Lanka, Africa and tropical Asia. The crude extract of this plant is commonly used treat whooping cough. The plant extract acts as diaphoretic and expectorant. The plant has also antirespiratory syncytial virus, antiarthritic, antiinflammatory and anti-fertility activities. The leaves and the tender branches are used for treatment of toothache, strengthening of gums, whooping cough and premature ejaculation. Whole-plant extracts contain iridoid glycosides, barlerin and verbascoside, which have shown potent activity against respiratory syncytial treating fever and several respiratory diseases.[20]

CNS STIMULANT ACTIVITY

The ethanol extract of Barleria prionitis exhibit antidepressant activity.

9) ALPINIA GALANGA

The plant Alapinia Galanga (Zingiberaceae) is a herb primarily used in cooking. It is widely distributed in various parts of India and Southeast Asia. It is 5 feet long with narrow leaves. The rhizome has the pharmacological actions such as antitumour, antifungal, antiulcer, antiallergic, antibacterial and antiviral activities. It is used for cough, asthma, bronchitis, headache, inflammation, rheumatoid arthritis etc.[21]

CNS STIMULANT ACTIVITY

The have kampheride, alpinin, galangin, methyl cinnamate, cincole and tannins like phlobaphines, flavonoids terpinoids and saponins which acts against cns disorders.

10)DIPLAZIUM ESCULENTUM

The vegetable fern plant (Diplazium esculentum) is a species found and used in East to South Asia and Oceania. It is a cold sensitive plant suitable for the warmer regions and tender to freezing temperatures. Diplazium esculentum (Retz.)Sw (FamilyAthyriaceae) consists of creeping and branched rhizome; scales brown, lanceolate and upto 7 to 15 mm in length; stripes fragile, straw coloured, 10-35 cm long.[22-26]

CNS STIMULANT ACTIVITY

The rhizome is considered as strong haemoptysis, used in cough, asthma, phthisis, fever, dyspepsia, stomachache, diarrhea and as antidysenteric, insect and pest repellant. [27]Young tips of fronds are used as tonics for health. Decoction of rhizome and young leaves are useful haemoptysis and constipation and used as antibacterial. It is found to be a good cnsstimultant.[28]

11)GINKGO BILOBA

Referred to as yinxing in Chinese medicine, the leaves of the ginkgo tree have long been used by herbalists for their effects as a brain tonic helpful for improving memory and the ability to concentrate. As a blood vessel dilator, it works directly with the circulatory system to increase oxygen supply to the brain and is additionally supportive to the heart and lung meridian systems.

CNS STIMULANT ACTIVITY

Ginkgo is known primarily for its beneficial effect on circulation, and it is through this function that it increases the energy of those taking it. At the same time, it can actually increase the oxygen capacity of the blood and improve mood.[29] It has also been shown to be very beneficial for those with early-stage Alzheimer's disease. It improves the vision of those with normal tension glaucoma.

References

- [1] Antidepressant activity of standardized extract of B. monnieri in experimental models of depression in rats Phytomedicine 2002; 9(3): 207-11.
- [2] Anju.,Bacopamonnieri a Preliminary Study Evaluating Its Anti-Stress Activity in Swiss Albino Mice, October – December 2011 RJPBCS Volume 2 Issue 4 Page No. 787, ISSN: 0975-8585.
- [3] Anti-stress effects of bacosides of B. monnieri. Phytother Res 2002; 16(7): 639-45
- [4] ChanghadiGovardhan Sharma, author. Ashwagandharishta -RastantraSarEvamSidhyaprayogSangrah - Krishna-Gopal Ayurveda Bhawan (Dharmarth Trust) Nagpur: 1938. pp. 743–744.
- [5] Narendra Singh, MohitBhalla, Prashanti de Jager,^{*} and MarilenaGilca., An Overview on Ashwagandha: A Rasayana (Rejuvenator) of Ayurveda.,Afr J Tradit Complement Altern Med. 2011; 8(5 Suppl): 208–213.
- [6] Muhammad AnisHashmat, RabiaHussain., A review on Acacia catechu Willd., Institute of Interdisciplinary Business Research 594 MAY 2013 VOL 5, NO 1
- [7] Guleria, S., Tiku, A., Singh, G., Vyas, D., & Bhardwaj, A., (2011). Antioxidant Activity and Protective Effect Against Plasmid DNA Strand Scission of Leaf, Bark, and Heartwood Extracts from Acacia catechu. Journal of Food Science. 76(7):C959-64. doi: 10.1111/j.1750-3841.2011.02284.x. Epub 2011 Aug 1.
- [8] Hazra, B., Sarkar, R., Biswas, S., & Mandal, N. (2011). Antioxidant and iron chelating potential of Pongammiapinnata and its role in preventing free radical induced oxidative damage in plasmid DNA. International Journal of Phytomedicine. 3(2):240-53.

- [9] Encyclopedia of Traditional Chinese Medicines -Molecular Structures by Jiaju Zhou, GuirongXie, Xinjian Yan.
- [10] Medicinal properties of Neem: New Findings by D.P. Agrawal.
- [11] P. SudhirKumar1 ,Debasis Mishra1 , Goutam Ghosh1 and Chandra S. Panda2,. Biological action and medicinal properties of various constituent of Azadirachtaindica (Meliaceae)" an Overview., Annals of Biological Research, 2010, 1 (3):24-34.
- [12] Kausik Biswas, Ishita Chattopadhyay, Ranajit K. Banerjee* and UdayBandyopadhyay., Biological activities and medicinal properties of neem (Azadirachtaindica)., CURRENT SCIENCE, VOL. 82, NO. 11, 10 JUNE 2002.
- [13] Immunomodulatory potenial of Curcuma longa: A review .,S.Mehrotra, G. Agnihotri, S. Singh and F. Jamal.,South Asian J ExpBiol; 3 (6) Special: 299-307; 2013 ISSN: 2230-9799 Vol. 3, Issue 6.
- [14] Job git ahi wang'ombe., evaluation of the central nervous system depressant effects of hydrazinocurcumin.,u29/27 4212009
- [15] Ashish p. Anovadiya1, jayesh j. Sanmukhani2, vishal k. Vadgama3, c. B. Tripathi.,evaluation of antiepileptic and memory retention activity of curcumin per se and in combination with antiepileptic drugs., asian j pharm clin res, vol 6, suppl 2, 2013, 145-148
- [16] Xia X, Cheng G, Pan Y, Xia ZH, Kong LD. Behavioral, neurochemical and neuroendocrine effects of the ethanolic extract from Curcuma longa L. in the mouse forced swimming test. J Ethnopharmacol 2007;110:356-63.
- [17] Tayyem RF, Heath DD, Al-Delaimy WK, Rock CL. Curcumin content of turmeric and curry powders. Nutr Cancer 2006; 55:126-31.
- [18] Doke p.p.*, tare h.l., sherikara.k., shendev.s., deores.r., damag.y. Central nervous system stimulant effect of the oils obtained from seeds of cucurbita maxima., journal of pharmaceutical biology, 1(1), 2011, 30-36
- [19] The Central Nervous System Activity of Barleria prionitis Linn. on the Locomotor Activity of Swiss Albino Mice using ActophotometerAmitesGangopadhyay*, JadupatiMalakar, Amitava Ghosh, Jyotirmoy Deb, Sanjay Dey, SupriyaDatta, Prabir Kumar Datta International Journal of Pharmaceutical & Biological Archives 2012; 3(2):403-405.
 [20] Savansaha, et al. Central pervous system stimulant
- [20] Sayansaha et al Cental nervous system stimulant ACTION of alpinia galanga L rhizome – a preliminary study.,international journal of experimental biology.,vol 15 october 2013 pp 828-832.
- [21] Singh HB. Potential medicinal Pteridophytes of India and their chemical constituents. J Econ Taxon Bot 1999; 23(1):63-786.
- [22]Bir SS. Key notes address on Ferns of India, their wealth exploration, diversity, growth conditions and conservation, Indian Fern Journal 1992; (9):IV-VI.
- [23] Kaushik P, Dhiman AK. Common Medicinal Pteridophytes. Indian ferns Journal 1995; (12):139145.
- [24] Quisumbing E. Medicinal Plants of Philippines, Manila Dept. Arg. Nat Res Bull 1951; 16(2):174176.

Licensed Under Creative Commons Attribution CC BY

- [25] Singh S, Dixit RD, Sahu TR, Some Medicinally Important Pteridophytes of Central India. Int J For Usuf Magt 2003; 4(2):41-50.
- [26] SemwalAmit ,FarswanMamta Singh .,In-Vitro Anthelmintic Activity of Diplazium esculentum (Retz.) Swiss Rhizome Extract ., Journal of Pharmacognosy and Phytochemistry., Vol. 1 No. 4 2012.
- [27] Semwal A, Kaushik S, Bhatt SP, Negi A, Antibacterial Activity of Diplazium esculentum(Retz.) Sw. Pharmacognosy Journal 2011; 21(4):74-76.
- [28] Laura L. Boles Ponto; Susan K. Schultz., Ginkgo biloba Extract: Review of CNS Effects., Annals of Clinical Psychiatry; Jun 2003; 15, 2.