

The Influence of ZnO NPs on Reproductive System Tissues of Albino Male Mice. Histopathological Study

Ruqayah Ali Salman

University of Al-Forat Al-Awsat Techniques

Abstract: ***Objective:** To evaluate the influence of Zinc Oxide nanoparticles (ZnO NPs) on Albino male mice reproductive system. **Methodology:** Albino male mice were used in this study, divided to three groups (6 mice to each): control group, first treated (150mg/kg.bw.) and second treated (350mg/kg.bw.). **Result:** After animal dissected, we tacked organs of reproductive system (testis, prostate, seminal vesicles and epididymis). The change in all tissues of reproductive system showed in (350mg/kg. bw.) dose treated with ZnO NPs, While the changes showed only in seminal vesicles and epididymis in the groups treated with (150mg/kg.bw.ZnONPs). **Conclusion:** The study established that a significant negative effect of ZnO NPs on Albino male mice reproductive tissues. **Recommendation:** Study the mechanism of ZnO NPs inside the body to know how can effects on body tissues.*

Keywords: Zinc Oxide nanoparticles, reproductive system, male mice

1. Introduction

Spermatogenesis is a complex process proliferation and differentiation of germ cell which leads to the production and release of spermatozoa from the testis. This elaborate process dependent on hormonal and dynamic interactions between in the Sertoli cell and the germ cells [1,2]. Many recent in vivo and in vitro studies identified that most nanoparticles (NPs) have a toxic effect on male germ cells [3,4]. Recent studies have shown that administration of NPs to mice result in their accumulation in the various tissues including the brain and the testis. This confirm that they easily pass through the blood-brain and blood-testis barriers [5,6]. Not all NPs will necessarily demonstrate an adverse effect leading to toxicity for example. Some NPs shown a beneficial or nontoxic effect on spermatogenesis [7,8]. It has been reported that nano selenium diet supplementation produced positive effects on sperm quality in male goats [7]. Same times nano particles can be beneficial or be damaging at the. Materials could be toxic and hazardous when they are converted into nano from [9]. In addition, the small size of nanoparticles allows them to overcome the defense barriers of the body without facing any obstacles [10]. ZnO nanoparticles have other applications besides the environmental usage, including rubber industry, glaze, cosmetics and medicine. Additionally, due to their antibacterial properties, ZnO nanoparticles can be used in preventive medicine against microbes associated with diseases and infections [11]. While Delouise, 2012 [12], report, ZnO nanoparticles, have toxic effects on cultured dermal fibroblast cell. The present study aims investigate side effects of various doses (150, 350 mg/kg) of ZnO NPs (80nm) on reproductive system.

2. Material and Method

In this experimental study how examined 18 adult albino male mice in age 3 week (mean weight=25±5g), purchased from the animal house of biotechnological center in Al-

Nahrain university. The animal divided to 3 groups (6 mice to each) allow free take pelleted and distal water under 23±2 °C (With 12 hours light and 12 hours/darkness). Control group was taken water and food, and (first and second treated) group respectively reactively received doses (150, 350 mg/kg) of ZnO nanoparticles prepared in distal water, after 15 days all animals was dissected, testis, prostate, seminal vesicles and epididymis tissues section was fixed in formalin solution, then embedded in paraffin, sectioned and stained with haematoxylin and eosin (H&E) for histopathological study.

3. Result

The study showed the influence of ZnO NPs on reproductive system tissues in Albino male mice. The tissue sections in the first treated with ZnO NPs (150mg/kg.bw.) indicate several changes in the tissue of seminal vesicles and compare with control group, histopathological section of seminal vesicles showed focal hyperplasia of epithelial cells (figure 6). While don't show any changes in testicular tissues, prostate and epididymis tissues in the first treated compare with control group (5,7,8,9). On the other hand, second treated with ZnO NPs (350mg/kg.bw) resulted incomplete spermatogenesis in certain seminiferous tubules, with vaculation of epithelial layer of testicular tissue (figure 10,11), While the histological sections of these seminal vesicles showed tissue mild to moderate proliferation of epithelial cells, mononuclear cells infiltration in the stroma (figure 12,13). In addition to the observed changes in the tissue of the prostate by the occurrence pinkish homogenous material in dilated of their acini, hyperplasia of epithelial lining cells of their acini that from papillary projection into their lumen (figure 14,15,16). It is also observed epididymis sections that their ked fill with sperms (figure 17,18).




Figure 1: Histopathological section in the testis showed normal structures, H & E stain. (a 100X, b 400X). (control group)




Figure 5: Histopathological section in the testis showed complete spermatogenesis with sperm in the lumen of seminiferous tubules (H & E stain 400X). (first treated)




Figure 2: Histopathological section in the seminal vesicles showed normal structures, H & E stain. (100X) (control group)



Figure 6: Histopathological section in the seminal vesicles showed focal hyperplasia of epithelial cells, H & E stain (400X). (first treated)




Figure 3: Histopathological section in the prostate showed normal structures, H & E stain. (100X) (control group)




Figure 7: Histopathological section in the prostate showed normal structure H & E stain (400X). (first treated)




Figure 4: Histopathological section in the epididymis showed normal structures, H & E stain. (400X). (control group)




Figure 8: Histopathological section in the epididymis showed their tubules filled with sperm, H & E stain. (100X) (first treated)




Figure 9: Histopathological section in the epididymis showed their tubules filled with sperms, H & E stain.(400X)(first treated)




Figure 13: Histopathological section in the seminal vesicles showed mononuclear cells infiltration in the stroma, H & E stain.(400X)(second treated)




Figure 10: Histopathological section in the testis showed incomplete spermatogenesis in certain seminiferous tubules, H & E stain.(400X)(second treated)




Figure 14: Histopathological section in the prostate showed dilated of their acini, H & E stain.(100X)(second treated)




Figure 11: Histopathological section in the testis showed that certain seminiferous tubules expressed incomplete spermatogenesis with vacuolation of epithelial layer, H & E stain.(400X)(second treated)




Figure 15: Histopathological section in the prostate showed pinkish homogeneous material in dilated of their acini, H & E stain.(400X)(second treated)




Figure 12: Histopathological section in the seminal vesicles showed mild to moderate proliferation of epithelial cells, H & E stain.(400X)(second treated)



Figure 16: Histopathological section in the prostate showed hyperplasia of epithelial lining cells of their acini that form papillary projection into their lumen, H & E stain.(400X)(second treated)




Figure 17: Histopathological section in the rete epididym showed sperm fill their lumen, H &E stain.(400X) (second treated)




Figure 18: Histopathological section in the epididymis showed their lumen filled with sperm, H & E stain.(400X)(second treated)

4. Discussion

Aparajitha J., 1992[13], showed the metals like Zinc(Zn), gold(Au) and nickel (Ni) play a significant role in male reproductive functioning and deficiency of these trace metals have negative impact on spermatogenesis and semen quality. Hajar S. *et al.*, 2014 [14] demonstrated that the influence of ZnO NPs on testicular tissues by sloughing of cell layers (stare) and impairment in the production of spermatogenesis in seminiferous tubules. ZnO NPs in our study, through their impact on Sertoli cells and occurrence of vacuolation phenomenon and the following apoptosis, have caused reduction in various cell lines including spermatogonia, round and elongated spermatids [14]. Also impact of ZnO NPs on reproductive organs in female animals [15]. It can be inferred that ZnO NPs are associated with oxidative stress (ROS), which result in an increase in DNA double-strand breakage and a decrease in sperm motility, ROS may penetrate across the cell membrane and inhibit the activity of some vital enzymes such as glucose 6-phosphatase dehydrogenase G6PDH. The ROS may also lead to mutations such as point mutations and polymorphism and thus reduce semen quality. This study agrees with Yoshida M, *et al.*, 2012[16], who studied the effect of NPs reduction is related to Leydig cell degeneration and the damage to the seminiferous tubules. As Talebiet AR, *et al.* 2013,[17] reported, regarding epididymis sperm characteristics including mobility, the percentage of normal cells compared to abnormal ones.

According to their study, low dose (150mg/kg bw.) administration of ZnO NPs don't show changes in tissue of testis and prostate gland. While the changes found in seminal vesicles. These effects were less significant compared to the dose of 350mg/kg bw. Concentration and

exposure mode have an important role in the toxicity of ZnO NPs [18].

5. Conclusion

This study has established that ZnO NPs have cytotoxic effects on testicular tissue in a dose dependent manner. But in general, ZnO NPs showed significant negative effects on albino male mice reproductive system.

6. Recommendation

- 1) More study need to know the mechanism by which ZnO NPs exerts its effects.
- 2) Study along time exposure to ZnO NPs, to know the impact of NPs on fertility in males.
- 3) Take different dose levels from ZnO NPs.

References

- [1] Franca LR; Ghosh S; Ye SJ and Russell LD. (1993). Surface and surface to volume relationships of the Sertoli cells during the cycle of the seminiferous epithelium in the rat. *Biol. Reprod.*, 49 (6):1215-28.
- [2] Boekelheide K; Fleming SL; Johnson KJ; Patel, SR and Schoenfeld HA. (2000). Role of Sertoli cells in injury-associated testicular germ cell apoptosis. *Exp. Biol. Med.*, 225(2): 105-15.
- [3] Braydich-Stolle LK; Lucas B; Schrand A; Murdock RC; Lee T and Schlarger JJ. (2010). Silver nanoparticles disrupt GDNF/Fyn kinase signaling in spermatogonial stem cells. *Toxicol. Sci.*, 116 (2): 577-89.
- [4] Braydich-Stolle L; Hussain S.; Schlager JJ and Hofmann MC. (2005). In vitro cytotoxicity of nanoparticles in mammalian germline stem cells. *Toxicol. Sci.*, 88(2): 412-9.
- [5] Borm PJ and Kreyling W. (2004). Toxicological hazards of inhaled nanoparticles potential implications for drug delivery. *J. Nanosci. Nanotechnol.*, 4(5): 521-31.
- [6] Chen Y; Xue Z.; Zheng D.; Xia K.; Zhao Y. and Liu T. (2003). Sodium chloride modified silica nanoparticles as a non-viral vector with a high efficiency of DNA transfer into cells. *Curr. Gene Ther.*, 3 (3): 273-9.
- [7] Shi L. and Xue W. (2011). Effect of sodium selenite, Se-yeast and nano-elemental selenium on growth performance, Se concentration and antioxidant status in growing male goats. *Small Ruminant Res.*, 96(1): 49-52.
- [8] Shi L.; Yang R.; Yue W.; Xun WJ.; Zhang CX. And Ren YS. (2010). Effect of elemental nano-selenium quality, glutathione peroxidase activity, and testis ultrastructure in male Boer goats. *Anim Reprod. Sci.*, 118 (2-4): 248-54.
- [9] Fiorito S.; Serafino A.; Andreola F. and Bernier P. (2006). Effects of fullerenes and single-wall carbon nanotubes on murine and human macrophages. *Carbon*, 44(6): 1100-1105.
- [10] Reddy LS.; Nisha MM.; Joice M. and Shilpa PN. (2014). "Antimicrobial activity of zinc oxide (ZnO) nanoparticle against *Klebsiella pneumoniae*". *Pharm. Biol.*, 15: 1.

- [11] Nadafi K.; Zare M.; Ali Mohammad BM.; Rastkari N. and Rastkari N. and Mousavi N.(2011). Evaluation of the toxicity of ZnO nanoparticles and titanium oxide by biological tests using Escherichia coli bacteria ATCC35218 and staphylococcus aureus ATCC25923, Health and Environment magazine, 2: 171-180.
- [12] Delouise LA, 2012. Applications of nanotechnology in dermatology. J. Investig Dermatol. 132(3p12): 964-75.
- [13] Apgar J. (1992). Zinc and reproduction: An update. J. Nutr. Biochem., 3: 266-78.
- [14] Hajar S.; Ali N. and Amir Masoud Mashayekh (2014). The effect of ZnO nanoparticles on the growth and puberty of newborn male wistar rats. International J. Basic Sci. Applied Research, 3(sp): 180-185.
- [15] Esmaeillou M; Moharamnejad M. and Hsankhani R. (2013). Toxicity of ZnO nanoparticles in healthy adult mice. Environmental toxicology and pharmacology, 35: 67-71.
- [16] Yoshida M.; Yoshida S.; Sugawara L. and Takeda K. (2002). Maternal exposure to diesel exhaust decrease expression of steroidogenic factor-1 and mulevian inhibiting of substance in the murine fetus. J. Heal. Sci., 48: 317-24.
- [17] Talebi AR; Khorsandi L. and Moridian M. (2013). The effect of Zinc oxide nanoparticles on mouse spermatogenesis. J. Assist. Reprod. Genet., 30(9): 1203-9.
- [18] Adamcakova, DA.; Stebounova LV.; Sung KJ.; Vorrink SU.; Ault AP.; OShaughnessy PT.; Grassian, VH. And Thorne, PS. (2014). Toxicity assessment of zinc oxide nanoparticles using sub-acute and sub-chronic murine inhalation models. Particle Fiber Toxicol., 11:1-15.

