

Cardiology, Hematology and Oncology Applications of Lactate Dehydrogenase in Serum

Thualfeqar Almohanna¹, Ali Sabi², Dhirgam Humaidy³

¹University of Kufa, College of Medicine, Biochemistry Department

²University of Kufa, College of Pharmacy, Biochemistry Department

³Al-Nahrain University, College of Science, Chemistry Department

Abstract: The major focus of this paper is to give a description of the various guidelines for a rationalized use of the lactate dehydrogenase in the various isoenzymes that are involved in the process of diagnosis coupled with the follow up. The estimation of the similarities specific diseases and lactate dehydrogenase that are presented as specificity, sensitivity, and the survival were all extracted. The application of the serum lactate dehydrogenase was found to be more applicable in myocardial infraction, ovarian dysgerminoma, and hemolytic anemia diagnosis. For the process of monitoring, the lactate dehydrogenase is seen to be relevant. Therefore rational use of the of the lactate dehydrogenase is seen to be equitably realized through ensuring that the various requests of the process of its determination are quite limited and the conditions that are mentioned above. Therefore, there was not any rationalized mechanism for the measurement of the lactate dehydrogenase and lactate dehydrogenase isoenzymes.

1. Introduction

The lactate dehydrogenase (LDH) is present in broad variety of organisms and the enzyme commission number is EC1.1.1.27. Moreover, EC1 is oxidoreductase, EC1.1 acts on the CH-OH group of the donor as well. EC1.1.1 is with NAD⁺ or NADP⁺ as acceptor, and EC1.1.1.27 is L-lactate dehydrogenase. However, LDH catalyzes the reaction of reduction pyruvate to lactate and oxidation lactate to pyruvate accompanied the NAD⁺ and NADH (Figure 1). According to Dawson *et al.*, the lactate dehydrogenase (LDH; E.C. 1.1.1.27) is a decisive enzyme necessitated in anaerobic metabolism; energy metabolism in skeletal muscle tissue is reduced pyruvate by LDH. This process produces ATP via glycolysis during anaerobic by regenerating NAD⁺.¹

The most detailed structural investigation and kinetic on a plant of LDH has been carried out from potato tuber tissue. This enzyme has five isozymes. LDH also consists of four subunits that are equal in size (Hoffman and Hanson, 1986).² Lactate dehydrogenases (LDH) have been exhibited to behave likely with respect to their effector (fructose 1,6-diphosphate) and substrates (pyruvate & NADH).³ Furthermore, Pridgaret *al* propose that several methods have been applied of purification. Those methods have separated pure LDH, such as ion exchange chromatography, affinity chromatography, and isoelectric focusing. Besides, the isoelectric focusing is not useful for LD-1 because it undergoes acid denaturation, while this does not happen with LD-5.⁴

2. Goals and Importance

The goals and objective of this paper is to give a description of the various guidelines for a rationalized use of the lactate dehydrogenase in the various isoenzymes that are involved in the process of diagnosis coupled with the follow up. This will ensure that there is a proper way to reduce the number of the requests that are forwarded in the diagnostic tests of the laboratory conducted tests. These will ensure that there is a clinical use of the provided information through a focus on the various approaches that can be formulated. The research are as in the abstract of the paper is the clinical value of the lactate dehydrogenase in serum.⁵ This is entirely based on the review of a number of the relevant literatures that are profitable to aid the analysis. The information originated from publications that are scientific having been obtained from the national library of medicine database of medicine. These are concerned with the applications clinically which does cover a number of the diagnosis, treatment and the monitoring of the disease for the case of the lactate dehydrogenase and isoenzymes for of measurements that are associated with the serum in various clinical fields. These capture cardiology, hematology, herpetology, and oncology. The studies that are presented in this paper involve proper definition of the populations of the tested patients, sampling




Figure 1: (A) The structure of LDH. PDB: 1cet. Image generated in PyMOL; (B) the reaction of conversion of pyruvate to lactate and vice versa.

timelines, and frequency and the various characteristics of the tests that are presented in methodologies. In order to address these a rational use of the of the lactate dehydrogenase is seen to be equitably realized through ensuring that the various requests of the process of its determination are quite limited and the condition that hinder the effectiveness of the study.⁶

Therefore, a number of the descriptions of the results of the systematic relevant literature review of the clinical associated values that are associated with the determination of the lactate dehydrogenase coupled with the isoenzymes that are found in the serum and therefore does give an appropriate indication of the appropriate time to span when and when not to make a request of the laboratory tests that have been conducted.⁷

3. Experimental Approach

In reference to a number for textbooks that are associated with medicine, the major clinical fields that are associated with the application of lactate dehydrogenase are found to be cardiology, hematology, herpetology, and oncology. The lactate dehydrogenase through a number of the evaluations that were made through the review of the literatures of the field, we were able to adequately obtain information. This was coupled with the aid of the bibliographic search systems that are computerized.⁸ These were aimed at providing a valid and proper judgment of the data that was provided. Therefore, according the methodologies that were provided, there effectiveness of the case with lactate dehydrogenase was found to be more pronounced in the clinical fields. These were largely pressed by the methods of sensitivity and specificity. The remission and rate of survival in the case with oncology and hematology were used frequently as criterion for test. The basic goal was to ensure that we found out the for the publications that are scientific and in English that did provide the relation existing between lactate dehydrogenase isoenzyme and lactate dehydrogenase measurements that are found in human serum and the clinical specialties as below.

Hepatology

The manuscript title was found to be having terms of the lactate dehydrogenase which was combined with the neoplasm invasiveness, anemia, and leukemia from the list thesaurus. The heading of the subject did contain the blood analysis, the therapy and the diagnostic use. Years of 1985-1996 were used for the search with the strategy thereby containing a combination of lactate dehydrogenase found with anemia, Hodgkin, anemia, and the non-Hodgkin with the blood analysis and the therapy.⁹

Cardiology

The titles of the manuscript did contain the lactate dehydrogenase and the liver. The main headings of the subject were liver function tests and the diseases of the liver and these were combined main subject heading of lactate dehydrogenase. The strategies were therefore formulated in the process of the search.

Hematology

The medical subject heading was further found to be having the term lactate dehydrogenase which was combined with

the terms of neoplasm invasiveness, and anemia. These were all found to be contained the subject analysis with all the above aspects.

Oncology

The mesh term was found to be having lactate dehydrogenase was further combined with various tumor-marker biology elements and the invasiveness. These were all found to be contained in the list of the thesaurus. Therefore the heading of the subject did contain the analysis of the blood, the therapy, and the blood diagnosis.¹⁰

Outcomes and Impacts

Cardiology

As for the case with cardiology, the research criteria that was formulated gave results of the in over the 35 publications. The population that was tested population for most of the conducted studies, it was found that a number of consecutive patients were found prevalent with the admissions to the coronary unit care. However, the populations prevalent with patients of geriatric and those that were collected by computer laboratories were also made a description of. The authors forwarded different criterions with revealed that a number of the manuscript tests did contain the mentions of specificity and sensitivity.¹¹

Hepatology

It was found out that the activity of the lactate dehydrogenase in the liver cells of which represented over 95% of the lactate dehydrogenase. This represented over 1/3 of the cells of myocardial with over 1800 times that are found to be prevalent in the serum. Therefore, lactate dehydrogenase was seen to be an effective use of the diagnosis in the case of the liver diseases.

Hematology

The search that was formulated in this case did contain over 90 publications of the forwarded measurement of the various activities of the lactate dehydrogenase isoenzymes and lactate dehydrogenase in serum among all the various types of the blood cells. These studies gave a report of the prevalence of leukemia which were both acute and chronic myeloid and lymphatic leukemia. These were coupled with deficiencies in the megaloblastic and iron. This gave over a total of over 24 publications.⁹

Oncology

The criteria presented for the search did result into over 208 articles that did cover a very wide field and a number of different diseases. Over 16 publications about the small cell cancer of the lung, ovarian dysgerminoma and the testicular germ tumor and the various non-seminoma were made a selection of.

4. Discussion

The distribution of the enzyme dehydrogenase was found to be wide. This is a fact which is coupled with a general trend that is forwarded to rationalize the lab. Requests and therefore often times raise the various doubts on the relevance of the lactate dehydrogenase in the field of medicine.⁹

In order to identify the clinical situations of which they are found to present the real values of lactate dehydrogenase and the compounding isoenzymes in serum. These were through a review of the literature that are found to be obtained through the computerized search of the bibliographic system for the search. The conclusions that can be final in this case of the relevant indications for the process of determining the lactate dehydrogenase. This is in prevalent with the various diagnosis of the hemolytic and megaloblastic anemia.¹⁰

5. Conclusion

The paper did therefore present the a formidable and relevant with the analysis of the description of the various guidelines for the case with the rationalized use of the lactate dehydrogenases in a number of the various isoenzymes in process of diagnosis. The paper did yield relevant results to validate the distribution of the enzyme dehydrogenase. This validated the fact of the prevalent doubts on the relevance of the lactate dehydrogenase in the medicine field.

References

- [1] Dawson, N. J., Bell, R. V., & Storey, K. B. (2013). Purification and Properties of White Muscle Lactate Dehydrogenase from the Anoxia-Tolerant Turtle, the Red-Eared Slider, *Trachemys scripta elegans*. *Enzyme Research*, 2013(c), 784973.
- [2] Hoffman, N. E., & Hanson, A. D. (1986). Purification and properties of hypoxically induced lactate dehydrogenase from barley roots. *Plant physiology*, 82(3), 664-670.
- [3] Sommer, P. A. S. C. A. L., Klein, J. P., Schöller, M., & Frank, R. M. (1985). Lactate dehydrogenase from *Streptococcus mutans*: purification, characterization, and crossed antigenicity with lactate dehydrogenases from *Lactobacillus casei*, *Actinomyces viscosus*, and *Streptococcus sanguis*. *Infection and immunity*, 47(2), 489-495.
- [4] Pridgar, E. M., Moses, G. C., & Henderson, A. R. (1984). Purification of lactate dehydrogenase isoenzymes one, two, and three from human erythrocytes. *Clinical chemistry*, 30(8), 1353-1357.
- [5] Cooker, L. A., Brooke, C. D., Kumari, M. E. E. N. A., Hofmann, M. C., Millan, J. L., & Goldberg, E. (1993). Genomic structure and promoter activity of the human testis lactate dehydrogenase gene. *Biology of reproduction*, 48(6), 1309-1319.
- [6] Chung, F. Z., Tsujibo, H., Bhattacharyya, U., Sharief, F. S., & Li, S. S. (1985). Genomic organization of human lactate dehydrogenase-A gene. *Biochemical Journal*, 231(3), 537-541.
- [7] Azuma, M., Shi, M., Danenberg, K. D., Gardner, H., Barrett, C., Jacques, C. J., ... & Zhang, W. (2007). Serum lactate dehydrogenase levels and glycolysis significantly correlate with tumor VEGFA and VEGFR expression in metastatic CRC patients.
- [8] Galardo, M. N., Regueira, M., Riera, M. F., Pellizzari, E. H., Cigorraga, S. B., & Meroni, S. B. (2014). Lactate regulates rat male germ cell function through reactive oxygen species. *PLoS one*, 9(1), e88024.
- [9] Robergs, R. A., Ghiasvand, F., & Parker, D. (2004). Biochemistry of exercise-induced metabolic acidosis. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 287(3), R502-R516.
- [10] Nageh, T., Sherwood, R. A., Harris, B. M., Byrne, J. A., & Thomas, M. R. (2003). Cardiac troponin T and I and creatine kinase-MB as markers of myocardial injury and predictors of outcome following percutaneous coronary intervention. *International journal of cardiology*, 92(2), 285-293.
- [11] Lippi, G., Mattiuzzi, C., Comelli, I., & Cervellin, G. (2013). Glycogen phosphorylase isoenzyme BB in the diagnosis of acute myocardial infarction: a meta-analysis. *Biochemica medica*, 23(1), 78-82.