

Modification on Two Parameters Kappa with Methods of Estimation

Rafid I. Ahmed

Al – Turath University College

Abstract: This paper deals with transforming two parameters Kappa (α, β) into three parameters Kappa (α, β, λ) using exponential of given CDF of Kappa by a new parameter (λ) then the new generated three parameters Kappa is obtained, also its cumulative distribution function (CDF), and then the derivation of r^{th} moment formula about origin is also derived after application of certain formulas of integral, and then we derive the formulas for maximum likelihood estimators of three parameters (α, β, λ), all the derivation required are explain.

Keywords: Modified three parameters Kappa, Moments estimators, Maximum likelihood Estimators.

1. Introduction

Kappa probability distribution was introduced by Mielke and Mielke and Johson, it is one of family distribution that represent data with positive skewed model^[2]. Many research work, on powering the given CDF by a power, from parameter to obtain a new exponential class family, Gupta et al^[1], (1998) work on generalization of standard exponential distribution to obtain a new exponentiated exponential family, Gupta and Kundu^[5] in (2001) discuss the properties of exponential family, Nadarajah, S.^[8] (2005) work on transforming Gumbel distribution to exponentiated one, Nadarajah, S. and Gupta, A. K.^[9,11] (2007), derive exponentiated gamma with application, Samir, K. A., El – Sayed, A. Elsherpiny and Yassmen, Y. A./11] (2009) work on maximum likelihood estimators (MLE's) for the unknown parameters and the corresponding asymptotic variance covariance matrix of the three-parameter Kappa distribution will be obtained under type II censored sample^[11]. Also D. S. Hassun, Inam A. N. and Layla M. N^[3] (2014) work on introducing three methods, for estimating the two parameters (α, β), Kappa distribution, these methods are maximum likelihood, maximum entropy, and L – moments.

The r^{th} non-central moments are derived, all the required formula for integrals needed explained, especially for maximum entropy method, since it requires four steps to be applied, using Lagrange multipliers and certain constraints.

In our research, we work on extending two parameters Kappa to one with three parameters, using exponentiated. The P.D.F is obtained which have three parameters (β) scale parameter and two shapes parameters (α, λ), also CDF is derived and r^{th} Moments about origin and maximum likelihood estimators are obtained.

2. Definition

The two-parameter Kappa distribution (α, β) is^[13]:

$$f(x) = \frac{\alpha}{\beta} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{-\frac{\alpha+1}{\alpha}} \quad x, \alpha, \beta > 0 \quad \dots (1)$$

$$F(x) = \frac{x}{\beta} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{-\frac{1}{\alpha}} \quad x > 0 \quad \dots (2)$$

A new family can be obtained from;

$$G(x) = [F(x)]^\lambda \quad \dots (3)$$

This implies;

$$G(x) = \frac{x^\lambda}{\beta^\lambda} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{-\frac{\lambda}{\alpha}} \quad x, \alpha, \beta, \lambda > 0 \quad \dots (4)$$

New generated P.D.F is;

$$\begin{aligned} g(x) &= \frac{x^\lambda}{\beta^\lambda} \left[\left(\frac{-\lambda}{\alpha} \right) \left(\alpha + \left(\frac{x}{\beta} \right)^\alpha \right) \right]^{-\frac{\lambda}{\alpha}-1} \left(\alpha \frac{x^{\alpha-1}}{\beta^\alpha} \right) \\ &\quad + \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}} \frac{\lambda x^{\lambda-1}}{\beta^\lambda} \\ g(x) &= \frac{\lambda}{\beta^\lambda} x^{\lambda-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}} - \frac{\lambda x^{\lambda+\alpha-1}}{\beta^{\lambda+\alpha}} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}-1} \\ g(x) &= \frac{\lambda}{\beta^\lambda} x^{\lambda-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}} \left[1 - \left(\frac{x}{\beta} \right)^\alpha (\alpha + \left(\frac{x}{\beta} \right)^\alpha)^{-1} \right] \end{aligned} \quad \dots (5)$$

This new generated three parameters kappa (α, β, λ);

$$m' r = e(x^r) = \int_0^\infty x^r g(x) dx \quad x, \alpha, \beta, \lambda > 0 \quad \dots (6)$$

$$\begin{aligned} &= \int_0^\infty \frac{\lambda}{\beta^\lambda} x^{r+\lambda-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}-1} dx \\ &\quad - \frac{\lambda}{\beta^{\lambda+\alpha}} \int_0^\infty x^{r+\lambda+\alpha-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}-1} dx \end{aligned} \quad \dots (7)$$

$$\text{Let } Z = \alpha + \left(\frac{x}{\beta} \right)^\alpha \rightarrow z - \alpha = \left(\frac{x}{\beta} \right)^\alpha$$

$$\begin{aligned} z - \alpha &= \left(\frac{x}{\beta} \right)^\alpha \rightarrow dz = \frac{\alpha x^{\alpha-1}}{\beta^\alpha} dx \\ m' r &= \frac{\lambda}{\beta^\lambda} I_1 - \frac{\lambda}{\beta^{\lambda+\alpha}} I_2 \end{aligned} \quad \dots (8)$$

$$I_1 = \int_0^\infty x^{r+\lambda-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}-1} dx \quad \dots (9)$$

$$I_2 \Rightarrow \int_0^\infty x^{r+\lambda+\alpha-1} \left[\alpha + \left(\frac{x}{\beta} \right)^\alpha \right]^{\frac{-\lambda}{\alpha}-1} dx$$

Now

$$(z - \alpha)^{\frac{1}{\alpha}} = \frac{x}{\beta}$$

$$\begin{aligned}
 x &= \beta(z - \alpha)^{\frac{1}{\alpha}} \\
 dx &= \frac{\beta}{\alpha}(z - \alpha)^{\frac{1}{\alpha}-1}dz \\
 \therefore I_1 &= \frac{\lambda}{\beta^\lambda} \int_0^\infty \beta[(z - \alpha)^{\frac{1}{\alpha}}]^{r+\lambda-1} z^{\frac{-\lambda}{\alpha}} (\alpha + (\frac{x_i}{\beta})^\alpha)^{-1} dz \\
 &\quad \frac{\lambda}{\alpha\beta^{\lambda-2}} \int_0^\infty [(z - \alpha)^{\frac{r+\lambda-1}{\alpha}} z^{\frac{-\lambda}{\alpha}} (\alpha + (\frac{x_i}{\beta})^\alpha)^{-1} dz \\
 &\quad \frac{\lambda}{\alpha\beta^{\lambda-2}} \int_0^\infty [(z - \alpha)^{\frac{r+\lambda}{\alpha}-1} z^{\frac{-\lambda}{\alpha}} dz \\
 &\quad \frac{\lambda}{\alpha\beta^{\lambda-2}} (\alpha^{\frac{r+\lambda}{\alpha}-1}) \int_0^\infty z^{\frac{-\lambda}{\alpha}} [(z - \alpha)^{\frac{r+\lambda}{\alpha}-1} dz \\
 &\quad \frac{\lambda}{\beta^{\lambda-2}} (\alpha^{\frac{r+\lambda}{\alpha}-2} (-1)^{\frac{r+\lambda}{\alpha}-1} \int_0^\infty u^{\frac{-\lambda}{\alpha}} [(1-u)^{\frac{r+\lambda}{\alpha}-1} du \\
 &\quad \frac{\lambda}{\beta^{\lambda-2}} (\alpha^{\frac{r+\lambda}{\alpha}-2} (-1)^{\frac{r+\lambda}{\alpha}-1} \frac{d(1-\frac{\lambda}{\alpha}) d(\frac{r+\lambda}{\alpha})}{d(1+\frac{r}{\alpha})} \quad \dots (10) \\
 I_2 &= \frac{\lambda}{\beta^{\lambda+\alpha}} \int_0^\infty x^{r+\lambda+\alpha-1} [\alpha + (\frac{x}{\beta})^\alpha]^{\frac{\lambda}{\alpha}-1} dx \quad \dots (11) \\
 z &= \alpha + \left(\frac{x}{\beta}\right)^\alpha
 \end{aligned}$$

$$(z - \alpha)^{\frac{1}{\alpha}} = \frac{x}{\beta}$$

$$dx = \frac{\beta}{\alpha}(z - \alpha)^{\frac{1}{\alpha}-1}dz$$

$$\begin{aligned} \text{Let } u &= \frac{z}{\alpha} \\ dz &= \alpha du \end{aligned}$$

$$\begin{aligned}
 I_2 &= \frac{\lambda}{\beta^{\lambda+\alpha}} \int_0^\infty \beta[(z - \alpha)^{\frac{1}{\alpha}}]^{r+\lambda+\alpha-1} z^{\frac{-\lambda}{\alpha}-1} \frac{\beta}{\alpha} (\alpha + (\frac{x_i}{\beta})^\alpha)^{-1} dz \\
 &\quad \frac{\lambda}{\beta^{\lambda+\alpha}} \frac{\beta^{r+\lambda+\alpha}}{\alpha} \int_0^\infty (z - \alpha)^{\frac{r+\lambda}{\alpha}} z^{\frac{-\lambda}{\alpha}-1} dz \\
 u &= \frac{z}{\alpha} \rightarrow \alpha du = dz \\
 \frac{\lambda}{\alpha} \beta^r \int_0^\infty (\alpha u - \alpha)^{\frac{r+\lambda}{\alpha}} (\alpha u)^{\frac{-\lambda}{\alpha}-1} \alpha du \\
 \lambda \beta^r (\alpha)^{\frac{r}{\alpha} + \frac{\lambda}{\alpha} - \frac{\lambda}{\alpha} - 1} \int_0^\infty (u - 1)^{\frac{r+\lambda}{\alpha}} u^{\frac{-\lambda}{\alpha}-1} du \\
 \lambda \beta^r \alpha^{\frac{r}{\alpha}-1} (-1)^{\frac{r+\lambda}{\alpha}} \int_0^\infty (1-u)^{\frac{r+\lambda}{\alpha}} u^{\frac{-\lambda}{\alpha}-1} du \quad \dots (12)
 \end{aligned}$$

$$I_2 = \lambda \beta^r \alpha^{\frac{r}{\alpha}-1} (-1)^{\frac{r+\lambda}{\alpha}} \frac{\Gamma(\frac{r+\lambda}{\alpha} + 1) \Gamma(1 - (\frac{\lambda}{\alpha} + 1))}{\Gamma(\frac{r}{\alpha})} \quad (13)$$

Therefore, the general formula of r^{th} moment for new modified three parameters kappa is;
 $m'r = I_1 + I_2$

Which represent equation (10) and equation (13) knowing that ($\alpha > \lambda$).

3. Estimation by Maximum Likelihood Method

Let $x_1, x_2 \dots x_n$ be a random sample from $g(x)$ defined by equation (5), then [7];

$$L = \prod_{i=1}^n g(x_i)$$

$$L = \lambda^n \beta^{-n\lambda} \prod_{i=1}^n x_i^{\lambda-1} \prod_{i=1}^n \left[\left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right) \right]^{-\frac{\lambda}{\alpha}} \prod_{i=1}^n [1 - xi\beta\alpha(\alpha+xi\beta)\alpha-1] \quad \dots (14)$$

$$\log l = n \log \lambda - n \lambda \log \beta + (\lambda - 1) \sum_{i=1}^n \log x_i$$

$$- \frac{\lambda}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$+ \sum_{i=1}^n \log \left(1 - \frac{\left(\frac{x_i}{\beta} \right)^\alpha}{\alpha + \left(\frac{x_i}{\beta} \right)^\alpha} \right)$$

$$\log l = n \log \lambda - n \lambda \log \beta + (\lambda - 1) \sum_{i=1}^n \log x_i - \frac{\lambda}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right) \dots (15)$$

$$\Rightarrow n \log \lambda - n \lambda \log \beta + (\lambda - 1) \sum_{i=1}^n \log x_i - \frac{\lambda}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$+ \left(\frac{x_i}{\beta} \right)^\alpha + n \log \alpha$$

$$- \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$= n \log \lambda - n \lambda \log \beta + (\lambda - 1) \sum_{i=1}^n \log x_i + n \log \alpha - (1$$

$$+ \frac{\lambda}{\alpha}) \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

Then;

$$\frac{\partial \log l}{\partial \lambda} = \frac{n}{\lambda} - n \log \beta + \sum_{i=1}^n \log x_i - \frac{1}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$\frac{\partial \log l}{\partial \beta} = \frac{-n\lambda}{\beta} - [(1 + \frac{\lambda}{\alpha}) \sum_{i=1}^n \frac{x_i^{-\alpha} \beta^{-\alpha} (-1) \log \beta}{\left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)}]$$

$$\frac{\partial \log l}{\partial \beta} = \frac{n}{\alpha} - [(1 + \frac{\lambda}{\alpha}) \sum_{i=1}^n \frac{1 + \left(\frac{x_i}{\beta} \right)^\alpha (1) \log \left(\frac{x_i}{\beta} \right)}{\left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)}]$$

$$+ \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \left(\frac{-\lambda}{\alpha^2} \right) \right)$$

$$\Rightarrow \frac{n}{\alpha} - [(1 + \frac{\lambda}{\alpha}) \sum_{i=1}^n \frac{1 + \left(\frac{x_i}{\beta} \right)^\alpha \log \left(\frac{x_i}{\beta} \right)}{\left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)}]$$

$$+ \frac{\lambda}{\alpha^2} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$\frac{\partial \log l}{\partial \lambda} = 0 \Rightarrow$$

$$\frac{n}{\lambda} = n \log \beta - \sum_{i=1}^n \log x_i + \frac{1}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)$$

$$\hat{\lambda}_{MLE} = \frac{n \log \beta - \sum_{i=1}^n \log x_i + \frac{1}{\alpha} \sum_{i=1}^n \log \left(\alpha + \left(\frac{x_i}{\beta} \right)^\alpha \right)}{n}$$

$$\frac{\partial \log l}{\partial \beta} = 0 \Rightarrow$$

$$\frac{n\lambda}{\beta} = (1 + \frac{\lambda}{\alpha}) \sum_{i=1}^n \frac{x_i^{-\alpha} \beta^{-\alpha} \log \beta}{(\alpha + (\frac{x_i}{\beta})^\alpha)}$$

Finally

$$\begin{aligned} \frac{n}{\hat{\alpha}} &= [(1 + \frac{\lambda}{\alpha}) \sum_{i=1}^n \frac{(1 + (\frac{x_i}{\beta})^\alpha) \log (\frac{x_i}{\beta})}{(\alpha + (\frac{x_i}{\beta})^\alpha)} \\ &\quad - \frac{\lambda}{\alpha^2} \sum_{i=1}^n \log^2 (\alpha + (\frac{x_i}{\beta})^\alpha)] \end{aligned}$$

Now we explain different terms of $p.d.f$ and $c.d.f$ according to different sets of parameters that is necessary to explain the shape and behavior of $p.d.f$ and $c.d.f$ as shown in the two figures below;




Figure 1: The C.D.F result is obtained have three parameters (β) scale parameter and two shapes parameters (α, λ).




Figure 2: The P.D.F result is obtained

4. Conclusion

We explain different figures representing the *p.d.f* of modified Kappa, and also c.d.f, using different of value. We see that all the curves are flexible, this is due to modification of Kappa.

Modification through exponent gives new *p.d.f* which is necessary when the values of random variable lie between zero and one.

We suppose the variation of estimators $\alpha=0.2, \beta = 1.8, \lambda = 0.8$, the results can show in Fig. (1) and Fig. (2) which is in red color.

Then suppose the variation of estimators $\alpha = 0.1, \beta = 0.6, \lambda = 1.2$, the results can show in Fig. (1) and Fig. (2) which is in green color.

We suppose the variation of estimators are $\alpha=0.4, \beta = 1, \lambda = 0.6$, the results can show in Fig. (1) and Fig. (2) which is in blue color.

References

- [1] A.Slam,M. and Shahbazi.Q.M.,(2007). “Economic Reliability test plans using the generalized exponential distribution”, Journal of statistics Vol 14.
- [2] BungonKumphon. (2012). “Maximum Entropy and Maximum Likelihood Estimation for the Tree-parameters Kappa distribution”, Open Journal of statistics, 2012,2,415-419.
- [3] D. S. Hassun, InamA.N.and Layla M. N. (2014),“Introducing Different Estimators of two Parameters Kappa-distribution”,International Journal of Advanced Research in Engineering and Technology 12,107-115.
- [4] Firouzabadi, Morteza, Katebi, Serajddin, (2013), “An extension of a discretization method to solve fuzzy fractional differential equation”, 13th Iranian Conference on Fuzzy Systems (IFSC) 1 – 5.
- [5] Gupta, R.D. and Kundu, D. (2001),“Exponentiated exponential Family; an alternative to gamma and Weibull”, Biometrical Journal 43,117-130.
- [6] Kundu,D.andGupto,R.D.(2005), “Estimator of $p(y < x)$ for generalized exponential distribution”, MetrikaVol.61.No.(3).291-308,2005.
- [7] Marco Cello, Giorgio Gnecco, Mario Marchese, and Marcello Sanguineti, (2011), “CAC with Nonlinearly-Constrained Feasibility Regions”, IEEE COMMUNICATIONS LETTERS, VOL. 15, NO. 4.
- [8] Nadarajah, S. (2005),“The exponentiatedGumbel distribution with climate application. Environ Metrics 17,13-23.
- [9] Nadarajah, S .and Gupta, A.K. (2007), “The exponentiated gamma distribution with application on drought data”, Calculate statistical association Bulletin 59,29-54.
- [10] Nan-Cheng Su, Arjun K. Gupta, (2015), “On some sampling distributions for skew-normal population”,

Journal of Statistical Computation and
Simulation Volume 85, Issue 17

- [11] Samir, K. A., El-Sayed, A. Elsherpieny and Yassmen, Y. A. (2009), "Parameter Estimation for Three-Parameter Kappa Distribution Under Type II Censored Samples", *Journal of Applied Sciences Research* 10, 1762-1766.
- [12] Tomasz J. Kozubowski. (2008), "Multitude of Laplace distributions", *Statistical Papers*, 3/11/2008
- [13] Yazhuo Zhang and Baodong Zheng, (2013), "The Stability Criteria with Compound Matrices", *Abstr. Appl. Anal.* Volume 2013, Special Issue (2013), Article ID 930576, 5 pages.
- [14] www.e-sciencencentral.org.