

Awareness of Adverse Effects of Corticosteroids Among Dental Undergraduates: A Survey

Faazila Fathima¹, Anitha Roy²

¹II BDS, Saveetha Dental College and Hospital, Saveetha University, Chennai

²Department of Pharmacology, Saveetha Dental College and Hospital, Saveetha University, Chennai

Abstract: ***Objective:** The objective of this study was to assess and evaluate the knowledge on adverse effects of corticosteroids among dental undergraduates. **Methods and materials:** A pretested questionnaire containing 17 questions were used to assess knowledge, attitude and practice (KAP) on awareness of adverse effects of corticosteroids was administered to 120 dental undergraduates studying in a private hospital between the age group of 17-25 years after explaining about the study. The questions were carefully studied and the corresponding answers were marked and was tabulated. **Result:** It was found that 95% of the students knew about corticosteroids. 60% of the students were sure about the adverse effects of corticosteroids and about 10% of them were unsure and 30% of them didn't know about the adverse effects. 75-80% of the students were aware of the different types like inhalational and oral corticosteroids and their advantages and disadvantages. 65% of the students were aware that corticosteroids has to be gradually stopped after chronic use. **Conclusion:** Our study shows that majority of the students were aware of what corticosteroids are and their adverse effects and can use this knowledge in their future practices. Corticosteroids being one of the most commonly prescribed anti inflammatory, anti allergic and immune suppressant drug is used for a variety of health conditions. Where all possible it is given as inhalation but mostly it is given through oral route. Thus the prescriber must be aware of the adverse effects so that unnecessary complications can be prevented.*

Keywords: corticosteroids, adverse, effect, knowledge, attitude, dental undergraduate

1. Introduction

Steroids, referred to as corticosteroids, are substances that are naturally produced in our body. They are produced by the adrenal glands and regulate our immune system and the salt-water balance and water in our system.(1) They help to reduce inflammation.(2) Corticosteroids includes two main classes which are glucocorticoids and mineralocorticoids. Glucocorticoids such as cortisol affect carbohydrate, fat, and protein metabolism, and have anti-inflammatory, immunosuppressive, anti-proliferative, and vasoconstrictive effects.(3) Mineralocorticoids such as aldosterone are primarily involved in the regulation of electrolyte and water balance by modulating ion transport in the epithelial cells of the renal tubules of the kidney.(3) Since their discovery in the 1940s, corticosteroids have become one of the most widely used and effective treatments for various inflammatory and autoimmune disorder. (3) Most of the anti-inflammatory and immunosuppressive actions of glucocorticoids are attributable, either directly or indirectly, to their interaction with the cytosolic glucocorticoids receptor, which alters gene transcription to either induce or repress gene transcription in both inflammatory leukocytes and in structural cells, such as epithelium. (4,5,6) Based on route of administration, there are many different corticosteroids like oral, topical and inhalation. Each of these have their advantages and disadvantages. Currently corticosteroids are widely used for clinical conditions. Steroids are known to cause many pathological effects including chemically induced diabetes. Considering the important risk factors with the use of these, this study evaluates the level of awareness among dental undergraduates.

2. Methods and materials

A pretested questionnaire containing 17 questions were used to assess knowledge, attitude and practice (KAP) on awareness of adverse effects of corticosteroids and was administered to 120 dental undergraduates studying in a private hospital between the age group of 17-25 years after explaining about the study. Questions regarding whether the student used corticosteroids, if they were aware of their side effects, different types of corticosteroids available, which route of administration has faster action, the advantages and disadvantages of the different types of corticosteroids were asked. The questions were carefully studied and the corresponding answers were marked and was tabulated.

3. Result

It was found that 95% of the students knew about corticosteroids (figure1). 60% of the students were sure about the adverse effects of corticosteroids and about 10% of them were unsure and 30% of them didn't know about the adverse effects (figure2). 75-80% of the students were aware of the different types like inhalational and oral corticosteroids and their advantages and disadvantages (figure3). Also, 35% of the students didn't know corticosteroids can be gradually stopped after chronic use (figure4). It was observed that 61% of the students didn't take corticosteroids. Most of the students (57%) preferred the oral route of administration and its efficiency. 52% of the students were not aware of the different combinations of corticosteroids and the adverse effects of corticosteroids overdose. Awareness of antidote for corticosteroids was minimal.




Figure1




Figure2




Figure3




Figure4

4. Discussion

This study was conducted to evaluate the level of awareness on adverse effects of corticosteroids among dental undergraduates. It was observed that majority of them were aware about the adverse effects. The different types of corticosteroids available and their uses were familiar to more than average number of the individuals. Few students were not aware that corticosteroids can be gradually stopped. Corticosteroids are commonly used in the treatment of cancer, primarily owing to their anti-inflammatory activities (7). In a prospective study the most common side effects associated with corticosteroid use were oral candidiasis, oedema, cushingoid facies, dyspepsia, and weight gain (8). A separate study states that hyperglycaemia occurs in a majority of hospitalised patients receiving high doses of corticosteroids (9). Many scientific and literature evidences highlight how the administration of corticosteroids results in a high incidence of mood elevation, satisfaction, and optimism.(10) Less frequently, euphoria, insomnia, and increase in motor activity may occur.(11) Corticosteroids have wide range of uses in dentistry. Steroids are used in intracanal medicaments such as Ledermix to reduce pulpal inflammation and prevent root resorption; widely used in oral medicine such as in vesiculobullous diseases, orofacial granulomatosis, temporal arteritis and other oral mucosal disorders.(12)Hooley and Hohl described several instances of steroid use in prevention of post-operative oedema and topical use on the lips and corners of the mouth to prevent ulceration and excoriation as a consequence of retraction during surgery.(13)Steroid administration influences the occurrence of root resorption. The percentage of root resorption has been found to be more on treatment with corticosteroids.(14)However steroid-antibiotic combinations like Ledermix have also been used as intracanal medicaments for management of root resorption with reasonable success. Steroids like hydrocortisone are also mixed with zinc oxide eugenol to be used as root canal sealers. It appears that the action of steroids on root resorption is chemistry dependent.(1)Despite the known numerous side effects, the use of corticosteroid is widely

spread considering the broad spectrum of clinical indications. Psychiatric adverse reactions are underestimated and therefore it is not always possible to identify the effective dose and at the same time the most secure. It seems only right to recall how the spontaneous reporting of adverse reactions by health professionals and patients is the easiest way to integrate the missing information on the potential and dangers of drugs.(15)

5. Conclusion

The present study shows that majority of the students were aware of what corticosteroids are and their adverse effects and can use this knowledge in their future practices. Corticosteroids being one of the most commonly prescribed anti inflammatory, anti allergic and immune suppressant drug is used for a variety of health conditions. Where all possible it is given as inhalation but mostly it is given through oral route. Thus the prescriber must be aware of the adverse effects so that unnecessary complications can be prevented.

6. Acknowledgement

I thank the participants of this study for giving their valuable time to fill the questionnaire sincerely.

7. Conflict of Interest

NIL

References

- [1] Vigneshwar Sambandam, Prasanna Neelakantan - Steroids in dentistry, Int. J. Pharm. Sci. Rev. Res., 22(2), Sep – Oct 2013
- [2] Sweetman SC et al., Martindale, The Complete Drug Reference, Pharmaceutical Press, 37th edition, 2011.
- [3] Liu, Dora et al "A practical guide to the monitoring and management of the complications of systemic corticosteroid therapy". Allergy, Asthma, and Clinical

Immunology : Official Journal of the Canadian Society of Allergy and Clinical Immunology. 9 (1): 30.

- [4] Singh N, Rieder MJ, Tucker MJ: Mechanisms of glucocorticoid-mediated antiinflammatory and immunosuppressive action. *Paed Perinatal Drug Ther.* 2004, 6: 107-115.
- [5] Newton R, Leigh R, Gienbycz MA: Pharmacological strategies for improving the efficacy and therapeutic ratio of glucocorticoids in inflammatory lung diseases. *Pharmacol Ther.* 2010, 125: 286-327.
- [6] Coutinho AE, Chapman KE: The anti-inflammatory and immunosuppressive effects of glucocorticoids, recent developments and mechanistic insights. *Mol Cell Endocrinol.* 2011, 335: 2-13.
- [7] Rhen, T. and Cidlowski J.A., 2005. Antiinflammatory action of glucocorticoids—new mechanisms for old drugs. *New Engl J Med*, pp353:1711-1723.
- [8] Drof, T.B. and Crawford, E.D., 2013. Management and challenges of corticosteroid therapy in men and metastatic castrate-resistant prostate cancer. *Annals of Oncology* pp. 24; 31–38.
- [9] Donihi, A.C., Raval, D., Saul, M., Korytkowski, M.T., De Vita, M.A., 2006. Prevalence and predictors of corticosteroid related hyperglycaemia in hospitalised patients. *Endocr Pract.* pp 358-62.
- [10] Lewis DA, Smith RE. Steroid-induced psychiatric syndromes: A report of 14 cases and a review of the literature. *J Affect Disord.* 1983;5:319–33.
- [11] Varney NR, Alexander B, Macindoe JH. Reversible steroid dementia in patients without steroid psychosis. *Am J Psychiatry.* 1984;141:369–72.
- [12] Veseau PJ, Medicating post extraction sockets, *J Oral Maxillafac Surg*, 58, 2000, 531-537.
- [13] Holley JR, Hohl TH, Use of steroids in the prevention of some complications after traumatic oral surgery, *J Oral Surg*, 32, 1974, 864-866.
- [14] Verna C, Hartug LE, Kalia S, Melsen B, Influence of steroid drugs on orthodontically induced root resorption, *Orthod Croniofac Res*, 9, 2005, 57-62.
- [15] Ciriaco, Miriam et al. "Corticosteroid-Related Central Nervous System Side Effects." *Journal of Pharmacology & Pharmacotherapeutics* 4.Suppl1 (2013): S94–S98. PMC. Web. 13 Feb. 2017.