

Comparative Study of Larvicidal Activity of Clerodendrum Inerme Leaves Extracts and Crude Oil of Roasted Coconut Shell Individually as Well as Mixture of Both

Rajendra Prajapati¹, Sidhesh Gaonkar², Dr. Satish S Kolte³, Dr. Mira Ramaiya⁴

^{1,2,3}K.C. College, Department of Chemistry, Churchgate, Mumbai

⁴Department of Zoonosis, Haffkine Institute, Parel (East), Mumbai

Abstract: Various fractions of leaves and crude oil of coconut shell were obtained in organic solvents like Petroleum ether, Chloroform and Methanol and were subjected to larvicidal activity individually as well as mixture of both against 3rd and 4th instar larvae of *Culex pipen fatigan* and *Aedes aegypti* mosquito larvae. The study of larvicidal activity were done by using World Health Organization (WHO) method. They were tested for 24-48 hrs for their mortality. The concentration ranged from 50ppm to 600ppm. Comparative study shows that individual extract of leaves and crude oil with Petroleum ether gave highest activity against *Aedes aegypti* and *Anopheles stephensi* whereas mixture of extracts of leaves and crude oil with Petroleum ether gave maximum activity against *Culex pipen fatigan* mosquito larvae.

Keywords: Fractionated extraction, *Culex pipen fatigan*, *Aedes aegypti*, *Anopheles stephensi*

1. Introduction

Mosquitoes are responsible for the spread of a wide range of diseases including Malaria, Yellow fever, Dengue, Chikungunya, Japanese encephalitis (JE), West Nile virus and Rift Valley fever.^(1,2) *Aedes aegypti* and *Aedes albopictus* are implicated as disease vectors for the spread of dengue, chikungunya and Zika virus and breed in a variety of containers. It is a common species in city premises and recorded breeding is predominantly in discarded tyres and solid waste containers in slum areas.⁽³⁾ Japanese encephalitis, formerly endemic in north-east in India, is currently spreading fast across the state of north region of India with confirmed cases and a high case-fatality rate affecting all ages which is caused by *Culex* species of mosquito.⁽⁴⁾ The incidence of malaria which is caused by *Anopheles* species is, however, gradually receding, with a consistent decline in cases over the past few years where as the no. of Dengue, Chikungunya and Japanese encephalitis cases has increased gradually due to climate change, rapid urbanization, a growing population and lack of sanitary waste and water disposal are all factors which are responsible for spread of mosquito borne diseases in India.

Clerodendrum inerme is a small tree, shrubs and herbs occurs predominantly in the mangrove forests of coastal India.^(5,6) The plant has many medicinal importance such as it is used for treatment of rheumatism, fever, hypertension, cough, skin rashes, muscular pain, dysentery and its Roots are used for venereal diseases.^(7,8,9) Also the leaves extract of plant is reported as active against mosquito larvae.^(10,11) The *Cocos nucifera* also known as 'Kalpa vriksha' in India is widely present in the coastal region of India. Its each and every part are used by human in various ways such as food, timber, medicine, fibre, oil and charcoal.^(12,13) Various studies

shows its highly medicinal value and it's larvicidal activity.⁽¹⁴⁾

As both the plants are active against mosquitos, it is an attempt to study their effect on mosquito larvae individually as well as mixture of both.

2. Materials and Methods

The leaves were sun dried and 1KG of leaves were fractionated with solvents like Petroleum ether, Chloroform and Methanol respectively by refluxion method using soxhlet extraction technique. Crude fractionated solvents extract (Petroleum ether, Chloroform and Methanol) then allowed to evaporate remaining solvent from it and to dry at room temperature. All three concentrated fractionated leaves extract are used to prepare samples of concentration 10ppm to 600 ppm and were tested against mosquito larvae.

1KG of dried coconut shell broken into pieces and subjected to bioactive crude oil extraction by heating in the earthen pot using patented method.⁽¹⁵⁾ The dark viscous crude oil then subjected to fractionated extraction using solvent such as Petroleum ether and Chloroform respectively. Crude fractionated solvents extract (Petroleum ether and Chloroform) then allowed to evaporate remaining solvent from it and to dry at room temperature. The concentrated fractionated solvent extracts of crude oil and whole crude oil were used to prepare samples of concentration 10ppm to 600 ppm and were tested against mosquito larvae.

Mixture was prepared by mixing Petroleum ether leaves extract with Petroleum ether crude oil extract, Chloroform leaves extract with Chloroform crude oil extract and Methanol leaves extract with whole crude oil respectively in the ratio of 1:1.

Bioassay

Bioassay was conducted in 500ml glass beakers containing 250 ml of dechlorinated water for each species separately. The test solution with four replicates was prepared according to the guidelines of WHO (2005) ⁽¹⁶⁾. A batch of 3rd and early 4th instar larvae (10 in no.) of *Culex pipen fatigan*, *Aedes aegypti* and *Anopheles stephensi* were collected from the field and introduced in each test solution and in an appropriate control solution. The results were observed at 24hour.

3. Results and Discussion

Among all samples Petroleum ether extract (Individual as well as combination sample) shown maximum activity against all three species of mosquito. With LC₅₀ – 31.85 Petroleum ether extract of crude oil(individually) had shown maximum activity for *Culex pipen fatigan* species of

mosquito larvae followed by Chloroform extract of combination sample with LC₅₀ – 34.55 and Petroleum ether extract of combination sample. For *Aedes aegypti* and *Anopheles stephensi*, Petroleum ether extract of crude oil (individually) found more active with LC₅₀ – 295.54 and LC₅₀ – 298.54 respectively. followed by Chloroform extract of crude oil(individually) and whole crude oil.

4. Conclusion

Present comparative study suggests that Petroleum ether extract individually as well as in combination is quite effective against all three types of mosquito species larvae and possess effective larvicidal properties. Hence can be a better replaceable option to chemical substituent which can be used as herbal insecticide for treating the mosquito larvae.

Table 1: Lethal concentration of organic solvent extracts of *C. inermis* plant leaves and crude oil of roasted coconut shell against third and fourth instar larvae of *Culex* mosquito species

Sr. No.	Sample/Extracts	Mosquito species	Sample concentration	% Mortality	LC50	LC90 (Estimated)	Regression equation
1	Petroleum ether (Leaves)		300ppm	100	91.38	196.384	Y=1.7465x-2.9662
2	Chloroform(Leaves)		300ppm	100	81.43	175.593	Y=1.5687x-1.8169
3	Methanol(Leaves)		600ppm	0	-	-	-
1	Petroleum ether (Crude oil)		100ppm	100	31.85	61.283	Y=2.617x-4.06
2	Chloroform (Crude oil)	<i>Culex pipen fatigan</i>	400ppm	100	176.30	346.264	Y=1.961x-5.14
3	Whole Crude oil		300ppm	100	149.35	260.014	Y=2.213x-6.08
1	Petroleum ether (Leaves+ Crude oil)		100ppm	100	36.525	86.525	Y=2.3271x-3.1674
2	Chloroform (Leaves+ Crude oil)		100ppm	100	34.55	85.162	Y=2.3748x-3.4123
3	Methanol+ Whole crude oil (Leaves+ Crude oil)		-	-	-	-	-

Table 2: Lethal concentration of organic solvent extracts of *C. inermis* plant leaves and crude oil of roasted coconut shell against third and fourth instar larvae of *Aedes* mosquito species

Sr. No.	Sample/Extracts	Mosquito species	Sample concentration	% Mortality	LC50	LC90 (Estimated)	Regression equation
1	Petroleum ether (Leaves)		600ppm	50	434.96	884.609	Y=1.064x-2.8647
2	Chloroform(Leaves)		600ppm	0	-	-	-
3	Methanol(Leaves)		600ppm	0	-	-	-
1	Petroleum ether (Crude oil)		600ppm	90	295.54	543.023	Y=2.958x-11.18
2	Chloroform (Crude oil)	<i>Aedes aegypti</i>	600ppm	80	295.07	656.865	Y=2.385x-8.56
3	Whole Crude oil		600ppm	80	565.56	691.059	Y=1.367x-4.0
1	Petroleum ether (Leaves+ Crude oil)		600ppm	0	-	-	-
2	Chloroform (Leaves+ Crude oil)		600ppm	0	-	-	-
3	Methanol+ Whole crude oil (Leaves+ Crude oil)		600ppm	0	-	-	-

Table 3: Lethal concentration of organic solvent extracts of *C. inerme* plant leaves and crude oil of roasted coconut shell against third and fourth instar larvae of *Anopheles* mosquito species

Sr. No.	Sample/Extracts	Mosquito species	Sample concentration	% Mortality	LC50	LC90 (Estimated)	Regression equation
1	Petroleum ether (Leaves)		600ppm	0	-	-	-
2	Chloroform(Leaves)		600ppm	0	-	-	-
3	Methanol(Leaves)		600ppm	0	-	-	-
1	Petroleum ether (Crude oil)		600ppm	90	298.54	784.609	Y=2.958x-11.18
2	Chloroform (Crude oil)	<i>Anopheles stephensi</i>	600ppm	50	746	1242.526	Y=2.385x-8.56
3	Whole Crude oil		600ppm	60	725.37	964.335	Y=1.367x-4.0
1	Petroleum ether (Leaves+ Crude oil)		600ppm	0	-	-	-
2	Chloroform (Leaves+ Crude oil)		600ppm	0	-	-	-
3	Methanol+ Whole crude oil (Leaves+ Crude oil)		600ppm	0	-	-	-

References

- [1] Mereta ST, Yewhalaw D, Boets P, Ahmed A, Duchateau L, Speybroeck N et al. Physico-chemical and biological characterization of anopheline mosquito larval habitats (Diptera: Culicidae): implications for malaria control. *Parasites & Vectors* 2013; 6(1).
- [2] P. Suganthi, M. Govindaraju, V. Thenmozhi and B. K. Tyagi. Survey of mosquito vector abundance in and around tribal residential areas. *Journal of Entomology and Zoology Studies* 2014; 2 (6): 233-239.
- [3] Dutta P, Mahanta J. Potential vectors of dengue and the profile of dengue in the north-eastern region of India: an epidemiological perspective. *Dengue Bulletin*. 2006; 30: 234-242.
- [4] V Dev, VP Sharma, K Barman. Mosquito-borne diseases in Assam, north-east India: current status and key challenges, WHO South-East Asia Journal of Public Health | January-June 2015 | 4 (1-2).
- [5] Neeta Shrivastava, Tejas Patel (2007), *Clerodendrum* and Healthcare: An Overview, Medical and Aromatic Plant Science and Biotechnology. Global Science Book, 1(1),142-150.
- [6] S.R. Harish and K. Murugan (2011), Biochemical and Genetical Variation in the Mangrove Associate *Clerodendron inerme* (L) Gaertn. Under Different Habitats of Kerala. *ASIAN J. EXP. BIOL. SCI.* 2(4), 553-561.
- [7] Jasvinder Kaur Chahal, Renu Sarin and Manvi malwal (2010), Efficacy of *Clerodendrum inerme* l.(Garden queen) against some human pathogenic strain, *International Journal of Pharma and Bio Science*, 1(4),219-223.
- [8] Chethana G S, Hari Venkatesh K R, S M Gopinath (2013), Review on *Clerodendrum inerme*, *Journal of Pharmaceutical and Scientific Innovation*, 2(2), 38-40.
- [9] Nilima Shukla, Chandra Shekhar Panda and Sagar Kumar Mehta (2014), *Pharmacognostical Studies on C. inerme*, 6(5),191-197.
- [10] Mr. RAJENDRA G PRAJAPATI, DR. SATISH S KOLTE, Mr. SIDHESH R GAONKAR. Larvicidal Activity of Fractionated *Clerodendrum Inerme* Leaves Extract. Volume: 4 | Issue: 7 | July 2015,203-204.
- [11] P.B. Patil, S.V. Kallakar, V.L. Kallakar, S.N. Holihosur (2013), *Clerodendrum inerme* Gaertn. Plant as an effective natural product against *Aedes aegypti* L. mosquito. *International Journal of Natural Products Research*. 2(2),36-38.
- [12] Miss. Pallavi P. Borate, Siddheshwar D. Disale and R. S. Ghalme. *International Journal of Advanced Scientific and Technical Research*. Studies on Isolation, Analysis and Antimicrobial Properties of Coconut Shell Oil. Issue 3 volume 2, March-April 2013.
- [13] E Prabhakar Reddy, T.Mohana Lakshmi. Coconut Water - Properties, Uses, Nutritional Benefits in Health and Wealth and in Health and Disease: A Review. *Journal of Current Trends in Clinical Medicine & Laboratory Biochemistry*: April-June 2014, Volume 2, Issue 2, page no. 6-18.
- [14] Rajendra Prajapati, Dr. Satish S Kolte, Sidhesh Gaonkar, Dr. Mira Ramaiya. Larvicidal Activity of Roasted Coconut Shell's Extracts against Different Species of Mosquitoes Found in and around Mumbai. *Imperial Journal of Interdisciplinary Research (IJIR)* Vol.2, Issue-1, 2016, pg. no. 137-139.
- [15] Satish Kolte (1998), Application of HPLC and HPTLC in the study of some pharmaceutical products and study of biologically active compounds from crude coconut shell oil, 230.
- [16] WHO, World Health Organization, Guidelines for laboratory and field testing of mosquito larvicides. WHO/CDS/whopes/GCDPP 2005;13.