# **International Journal of Science and Research (IJSR)**

ISSN (Online): 2319-7064

Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

# Sideropenic Dysphagia

# Tsander Tito Prince BDS<sup>1</sup>, K. Srinivaas BDS<sup>2</sup>, K. Thafseera BDS<sup>3</sup>, Dr J. Jananee MDS<sup>4</sup>

<sup>1</sup>CRRI, Asan Memorial Dental College & Hospital, Chengalpattu

<sup>2</sup>CRRI, Asan Memorial Dental College & Hospital, Chengalpattu

<sup>3</sup>CRRI, Asan Memorial Dental College & Hospital, Chengalpattu.

<sup>4</sup>Professor, Department of Oral Pathology Asan Memorial Dental College &Hospital, Chengalpattu

Abstract: Siderophic dyspahagia is a premalignant condition due to vitamin deficiency. If untreated may lead to Squamous cell carcinoma, the most commonest oral cancer in India. A through knowledge of this pre cancerous condition is an must for all the clinical practioners which is discussed in detail in this article

Keywords: Precancerous condition, Plummer Vinson syndrome

#### 1. Introduction

Plummer–Vinson syndrome (PVS), also called Paterson–Brown–Kelly syndrome or sideropenic dysphagia, is a rare disease characterized by difficulty in swallowing, iron deficiency anaemia, glossitis, cheilosis and oesophageal webs<sup>1</sup>.

The reduction in the prevalence of PVS has been hypothesized to be the result of improvements in nutritional status and availability in countries where the syndrome was previously described. It generally occurs in postmenopausal women. Its identification and follow-up is considered relevant due to increased risk of squamous cell carcinomas of the oesophagus and pharynx <sup>2,3</sup>.

## 2. Etiology

The cause of PVS is unknown; however, genetic factors and nutritional deficiencies may play a role. It is more common in women particularly in middle age (peak age is over 50). In these patients, oesophageal squamous cell carcinoma risk is increased therefore, it is considered a premalignant process<sup>2,4</sup>.

The condition is associated with koilonychia, glossitis, inflammation of the lips (cheilitis), and splenomegaly. Oesophageal web in Plummer-Vinson syndrome is found at upper end of oesophagus (post cricoid region) and Schatzki ring may be found at the lower end of oesophagus.

# 3. Clinical and Radiographic Features

Most reported patients with PVS are between 30 and 50 years of age. Patients typically complain of burning sensation associated with the tongue and oral mucosa. Sometimes the discomfort is so severe that dentures cannot be worn.

Angular Cheilitis is often present and may be severe. Marked atrophy of the lingual papillae, which produces a smooth, red appearance of the dorsal tongue, is seen clinically<sup>5,6</sup>.

Patients also frequently complain of difficulty in swallowing (Dysphagia) or pain on swallowing. An evaluation with endoscopy or oesophageal barium contrast radiographic studies usually shows the presence of abnormal bands of tissue in the oesophagus, called oesophageal webs.

Another sign is an alteration of the growth pattern of the nails, which results in a spoon shaped configuration (koilonychia). The nails may also be brittle<sup>7</sup>.

Symptoms of anaemia may prompt patients with Plummer-Vinson Syndrome to seek medical care. Fatigue, shortness of breath, and weakness are characteristic symptoms<sup>8</sup>.

## 4. Diagnosis

Following clinical presentations may be used in the diagnosis of this condition.

- 1) Dizziness
- 2) Pallor of the conjuctiva and face
- 3) Erythematous oral mucosa with burning sensation
- 4) Breathlessness
- 5) Atrophic and smooth tongue
- 6) Peripheral rhaggades around the oral cavity

The following tests are helpful in the diagnosis of Plummer-Vinson syndrome.

#### 5. Treatment

Treatment is primarily aimed at correcting the irondeficiency anaemia. Patients with PVS should receive iron supplementation in their diet. This may improve dysphagia and pain. If not, the web can be dilated during upperendoscopy to allow normal swallowing and passage of food 9,10,11

### 6. Conclusion

As this condition occurs due to vitamin deficiency, it can be diagnosed at the earliest as the symptoms can present early in the oral cavity. Thus, as far as dentists are concerned, they play a major role in early diagnosis of the Plummer-Vinson

Volume 6 Issue 9, September 2017

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

Paper ID: ART20176884 1373

# **International Journal of Science and Research (IJSR)**

ISSN (Online): 2319-7064

Index Copernicus Value (2015): 78.96 | Impact Factor (2015): 6.391

syndrome. So the symptoms can be suppressed and early treatment can be commenced as soon as possible, which would decrease the rate of occurrence of the disease. So, the dentists should be much aware of these conditions and help the physicians to undergo preventive treatment as early as possible.

#### References

- [1] S. Hasan, N. I. Khan, and A. Siddiqui, "Plummer Vinson Syndrome-A premalignant condition-an overview of literature," Journal of Medical and Dental Sciences, vol. 1, no. 1, pp. 28–30, 2013.
- [2] S. S. Bakshi, "Plummer vinson syndrome—is it common in males?" Arquivos de Gastroenterologia, vol. 52, no. 3, pp. 250–252, 2015.
- [3] S. K. Swain, R. Panigrahy, and M. C. Sahu, "Plummer vinson syndrome in a male and his chromosomal study—a case report," Egyptian Journal of Medical Human Genetics, vol. 16, no. 3, pp. 283–286, 2015.
- [4] G. Novacek, "Plummer-vinson syndrome," Orphanet Journal of Rare Diseases, vol. 1, no. 1, article 36, 2006.
- [5] F. Demirci, M. C. Savaş, N. Kepkep et al., "Plummervinson syndrome and dilation therapy: a report of two cases," Turkish Journal of Gastroenterology, vol. 16, no. 4, pp. 224–227, 2005.
- [6] S. M. Naik, M. C. Shivakumar, M. K. Appaji, S. Ravishankara, S. S. Naik, and C. de Souza, "A case of plummer-vinson syndrome esophageal web dysphagia treated by dilatation with cuffed endotracheal tube," International Journal of Head and Neck Surgery, vol. 2, no. 3, pp. 161–165, 2011.
- [7] A. Samad, N. Mohan, R. V. Balaji, D. Augustine, and S. G. Patil, "Oral manifestations ofplummer-vinson syndrome: a classic report with literature review," Journal of International Oral Health, vol. 7, no. 3, pp. 68–71, 2015.
- [8] D. Lawoyin, E. Reid, F. Sam, T. Obayomi, and A. Bonnick, "Plummer vinson syndrome: a case report and literature review," The Internet Journal of Dental Science, vol. 5, no. 2, 2006.
- [9] Y.-C. Wu, Y.-P. Wang, J. Y.-F. Chang, S.-J. Cheng, H.-M. Chen, and A. Sun, "Oral manifestations and blood profile in patients with iron deficiency anemia," Journal of the Formosan Medical Association, vol. 113, no. 2, pp. 83–87, 2014
- [10] K. Vittal, S. S. Pandian, and T. Malarkodi, "lummer–vinsonsyndrome: a case report and medical management," IJSS Case Reports & Reviews, vol. 2, no. 6, pp. 10–13, 2015. <u>View at Google Scholar</u>
- [11] C. G. Mahesh, "Plummer vinson syndrome in males: a rare case," Medica Innovatica, vol. 4, no. 2, pp. 41-42, 2015.

Volume 6 Issue 9, September 2017 www.ijsr.net