

The Effects of Using Mandibular Free End Frame Denture First and Second Class Lever toward on Level IL-1 β of Abutment Teeth

Lisda Damayanti¹, Harmas Yazid Yusuf², Agus Susanto³, Gian Nur Alamsyah⁴

^{1,2,3,4} Universitas Padjadjaran, Faculty of Dentistry, Jl. Sekeloa Selatan 1, Bandung Indonesia

Abstract: *The purpose of this research is to analyze the effects of using mandibular free end frame denture first and second lever towards on levels IL- β of abutment teeth. **Materials and Steps:** 38 cases of missing mandibular free end teeth, either unilateral or bilateral, were made a free end frame denture design retainer first and second lever on each groups (19 cases). Before insertion and after 12 weeks of using mandibular free end frame denture, they were taken gingival crevicular fluid to measure the levels of IL-1 β d by using ELISA (Enzyme Linked Immunosorbent Assay) method. This research is an experimental design research, using single blind which is given actions by using mandibular free end frame denture first and second class lever. Statistical analysis begins with doing characteristic test of both groups. Both groups' numerical data are done to compare each groups' first and second class lever with non-paired test (data distributes normally) and Mann Whitney (data doesn't distribute normally). According to non-paired test, there were obtained p value 0.000 (p value < 0.005) which has statistic differentiation for IL-1 β level using mandibular free end frame denture first and second class lever. IL-1 β level of teeth abutment of first class lever has higher than second class lever. **Conclusion:** First class lever has a greater influence on elevated inflammation cytokin level of IL-1 β teeth abutment.*

Keywords: Mandibular free end frame denture, first class lever, second class lever, IL-1 β level

1. Introduction

Missing maxilla and mandibular posterior teeth effects mastication function a lot, the losing of continuity of dental arch results not balanced stomatognathic system inside oral cavity.¹ One of undertaken efforts to replace missing mandibular posterior teeth is by making removable partial denture (RPD).^{1,2,3} Free end RPD/free end denture is a removable denture which supported by either the teeth or tissue that covers residual ridge where on one or both saddle sides are expanded towards posterior. This kind of removable denture doesn't get the whole support from the teeth, but also depends on residual ridge as a support. Abutment teeth is a relatively passive support, while the soft tissue which covers residual ridge has a bigger and more variated moving degree.^{4,5,6}

The most common issues of free end denture is that the removable denture stays unstable, it moves easily either horizontally or vertically (rotating). This caused by tissue compressibility differences between anterior and posterior free end saddle. The movement of soft tissue under removable denture base approximately 350-500 μ m, while movement of the teeth around 20 μ m with the same pressure.⁷ Then, the unstable removable denture will cause issues towards remained oral cavity health. Because there are differences of support compressibility between one and other mucosa, between mucosa and abutment teeth periodontal tissue, the removable denture will lifted vertically when masticating. Also, because there's no teeth beside saddle's distal, its free end will move more freely.^{1,3,5,13}

Leverage towards apical from unstable removable denture's base will cause the masticatory forces distribute unequally. Apical towards masticatory forces will concentrate more on the posterior side (free end side), it causes too much forces (overload/overfunction) which next causes greater resorption

of alveolar residual ridge on the side.^{1,14}

Leveraged removable denture, besides causing greater resorption of alveolar residual ridge, also leverages abutment teeth. The main factor which causes greatest leverage force is the pressure/masticatory force, that causing leverage on the saddle side will move towards apical/alveolar residual ridge. First leverage, where its retention point and masticatory loads are across the fulcrum point/line, the free end removable denture will move apically and the abutment teeth as if rotated and pulled towards posterior. Because this happens continually, there will be damages on abutment teeth's periodontal tissue.

Second leverage happens when retention point and masticatory loads (on free end saddle) are on one side with the fulcrum point/line. On this circumstances, if the free end saddle pressured apically because of masticatory force, retentive arms will also going down towards apical, so that the abutment teeth won't leveraged, but causing removable denture becomes more unstable compared first leverage when masticatory force happened towards apical. Leverages caused by masticatory force towards apical are concentrated around alveolar residual ridge.^{1,3,5,14,15}

Bio-mechanic principles which must be noticed are; the emerged of removable denture's leverage may causes pressures towards abutment teeth, when free end removable denture's saddle is going down when accepting masticatory loads will causes leverage on abutment teeth. The forces obtained by that abutment teeth also called torque force. Occlusal loads distribution on teeth and alveolar bones covered by soft tissue, factors affecting the value of distributed loads on abutment teeth are; the longer the removable denture, then the bigger forces obtained by abutment teeth, a wide alveolar residual ridge will obtain more functional forces so there will be minimal forces

obtained by abutment teeth, a healthy mucosa with normal thickness endure occlusal loads better compared to thin mucosa, atrophy and occlusion factor are balanced.^{1,22} Hill and friends pointed that mucosal inflammation that caused by tissue under removable denture base which got pushed, may cause a variety of chronic inflammations, immune reactions, and broken periodontal tissue through the making of arachidonat acid metabolic or Interleukin (IL), and also has a strong activity of alveolar bone resorption IL-1 β acts as the key of mediator inflammation to determine periodontal health parameter.^{23,24,26}

2. Materials and Steps

The research is done to 38 cases of missing mandibular free end posterior teeth/mandibular free end denture, along with missing unilateral and bilateral teeth cases, the subjects' age range are around 35 until 54 years old. Cases missing free end bilateral teeth are around 68,4%. The research's plot includes the preparation of research's subjects which are protocol preparation, research form and permissions from research's ethical commissions, and also mouth preparation on research subjects. The making of mandibular free end frame denture first and second leverage are in a private lab. Gingival crevicular fluid were taken to measure IL-1 β and level with ELISA method which has been done before and 12 weeks after using mandibular free end frame denture first and second leverage. Before the fluid was taken, the first and second mandibular premolar teeth were isolated with cotton roll to prevent samples from salivary contamination and blood, then those were dried with air sprayer. Sterilized paper point were put into gum sulcus in the depth of 1 mm for 20 seconds. Paper point is gone inside the tube eppendorf which already filled with Phosphat Buffer Solution (PBS) and kept it inside a box filled with ice, and then the sample got moved to -70 Celcius degrees until the time of concentration test is done.

Examination and measurement with ELISA method for IL-1 β are done through the making of standard assay IL-1 β curve.

The making of standard assay IL-1 β curves by creating 6 dilution multilevel variations, which are 125 ; 62,5 ; 31,2 ; 15,6 ; 7,8 ; and 3,9 pg/mL from IL-1 β standard with 250 pg/mL concentration, also with Calibrator RD5-5. Put the standard/blank which has been made into the well in amount of 200 β L. Do the incubation with room temperature for 2 hours with see saw. Throw away the solution, then wash it with Wash Buffer 1x for 3x repeating. Add IL-1 β Conjugate as many as 200 β L, then incubate it on room temperature for 1 hour with see saw tool. Throw away the solution again and wash it using wash buffer 1x for 3x repeating. Add 200 β L substrat solution A and substrat solution B with comparison 1:1. Do the incubation for 20 minutes and protect it from sunlight (the color will turn blue). Add stop solution into the well, then there will be changing from blue to yellow. Measure it before 30 minutes on 450 nm and 490 nm.

Procedure of examination and measurement of gingival crevicular fluid's IL-1 β level

Put the standard/blank which has been made into the well in

amount of 200 β L. Throw away the solution, then wash it with Wash Buffer 1x for 3x repeating. Throw away the solution again and wash it using wash buffer 1x for 3x repeating. Add 200 β L substrat solution A and substrat solution B with comparison 1:1. Do the incubation for 20 minutes and protect it from sunlight (the color will turn blue). Add stop solution into the well, then there will be color changing from blue to yellow. Measure it before 30 minutes on 450 nm and 490 nm.

3. Result of The Research

Table 1: Research Subject's Characteristic (N=38)

Age	
Mean \pm SI	44.76 \pm 7.820
Median	47.000
Interval (min-max)	34.00-54.00
Gender	
Male	10(26.3%)
Female	28(73.7%)
Missing Mandibular Posterior Teeth Cases	
Unilateral free end	12 cases (31.6%)
Bilateral free end	13 cases (68.4%)

Numerical data are presented with mean, standard intersection, median, while categorical data are presented with value/frequency and percentage; SI: Standard Intersection

Table 1 describes the characteristic of overall research patient subjects according to their ages and genders. The average number of age is 44.76 \pm 7.820. There are 10 males or 26.3% and 28 females or 73.7%. The cases of missing mandibular unilateral free end posterior teeth are 31.6% and bilateral free end are 68.4%.

Table 2: Delta of IL-1 β Level of Abutment Teeth between First and Second Class Lever

IL-1 β (pg/ml)	Group		P Value
	First Class Lever	Second Class Lever	
	N=19 Cases	N=19 Cases	
Delta IL-1β			0.003*
Mean \pm Std	173.67 \pm 93.442	143.35 \pm 84.773	
Median	144.860	150.520	
Range (min-max)	43.31-373.53	-19.49-315.91	

The average of delta IL-1 β level for first class lever: 173.67 \pm 93.442 pg/ml, second class lever: 143.35 \pm 84.773 pg/ml. According to non paired test, there were obtained p value 0.003 (p value<0.005) which has statistic differentiation for IL-1 β level using mandibular free end frame denture first and second class lever. IL-1 β level of teeth abutment of first class lever has higher than second class lever.

4. Discussion

Interleukin-1 (IL-1) is a multifunctional cytokines with a wide activity on various tissues and also a mediator of immune cell that functioned on controlling bone resorption and increase prostaglandin synthesis.¹⁴ IL is a mediator key from body respond to micro-bacterial , immunology

reactions, and tissue trauma. IL-1 consists of 2 peptides, which are α and β , interleukin 1 β (IL-1 β) secreted with monocyte, some macrophages, endothelial cells, fibroblasts, and epidermal cells which are activated by some kind of stimulus.¹² IL-1 β increases PMN bond and monocyte/macrophage towards endothelial cells, stimulating prostaglandin E2 production, release lysosomal enzymes, and stimulating bone resorption.^{12,13} The value of IL-1 β will increase on various kinds of immune responses, including bone resorption and inflammation responses such as gingivitis and periodontitis.^{14,15,16}

This result shows that there were significant differences and patient's IL-1 β level also increased from before and after using mandibular free end frame denture either on first and second class lever (table 2). This result suits Grieve and friends' research which stated that if there is any mechanical pressure, then IL-1 β production will increase, periodontal tissue cells will produce IL-1 β if it got pressured and will increase inside the gingival crevicular fluid.¹⁷ Kurtis and friends' research stated that IL-1 β on periodontal tissue and gingival crevicular fluid will increase along with the severity of the disease.^{9,16} This result shows the increase of IL-1 β level on gingival crevicular fluid after using mandibular free end frame denture first class lever is higher than second class lever (table 2). This higher IL-1 β is caused because on first class lever, the force direction from distal to mesial, fulcrum on distal, are suitable with biomechanical law, fulcrum point put in the middle of retention point and loads, so if there any loads that touch the base of removable denture apically, then the retentive hands will move occlusally to balance this movement.⁹ The end of retentive hands which located beside distal will keep leveraging the abutment teeth which will cause inflammation on that abutment teeth, this shows that mechanical pressure on tissue will increase the level of IL-1 β .¹⁶

Kurtis and friends' research stated that mechanical pressure causes inflammation on abutment teeth's supporting tissue, which will cause damage on adhesion between supporting tissue and alveolar bone.¹⁸ The using of removable denture may also increase the accumulation of plaque and bacteria that are inside the plaque and interacts on epithelial cell, leucocytes, and fibroblasts that stimulate cytokines production such as IL-1 β .¹⁶

5. Conclusion

The result of this research showed level of IL-1 β ($p=0.003$) on teeth abutment higher on first class lever in the using of mandibular free end frame denture.

References

- [1] Carr AB, McGivney GP, Brown DT. McCracken's removable partial prosthodontics. 11th ed. St.Louis Mosby.2005;3-299.
- [2] Singla SG, Lal J. Removable partial dentures designing: forces as primary concern. JIPS.2006;6(1):179-84.
- [3] McGivney GP, Castleberry DJ. Removable partial prosthodontics.9th ed. R.R Donnelly and Sons Co.,USA.1995;1,114,160.
- [4] Fueki K, Igarashi Y, Maeda Y, Baba K, Koyano K, Sasaki K et al. Effect of prosthetic restoration on oral health-related quality of life in patients with shortened dental arches: a multicentre study. J Oral Rehabil.2015;4(1)15-22.
- [5] McCracken WL. Contemporary partial denture designs. J Prost Dent.2004; 92 (5):409-17.
- [6] The Academy of Prosthodontics. The Glossary of prosthodontics term. 8th ed. J prosthet Dent.2005;94(1):10-92.
- [7] Curtis DA, Wagnild GW, Finzen FC. Incidence of various classes of removable partial dentures. J Prosthet Dent.2002;67(5):664-7.
- [8] Shahmiri R, Aarts JM, Bennani V, Das R, Swain MV. Strain distribution in a Kennedy class II implant assisted removable partial denture under various loading conditions. Int J Dent.2013;35:1279.
- [9] Davenport JC, RM Basker, JR Heath, JP Ralph, PO Glantz. The removable partial denture equation. J Br Dent.2000;189(8):414-424.
- [10] Patrnogic V, Todorovic A, Scepanovic M, Radovic K, Vesnic J, Grbovic A. Free end saddle length influence on stress level in unilateral complex partial denture abutment teeth and retentions elements. Vojnosanit Pregl.2013;70(11):1015-22.
- [11] Zarb GA, Bolender CL, Carlsson GE. Biomechanics of edentulous prosthodontics treatment for edentulous patients. St Louis Mosby.2004.
- [12] Hill PA, Tumber A, Papaioannou S, Meikle MC. The cellular actions of interleukin-II on bone resorption in vitro. Endocrinol.1998;(139):1564-72. In: Kingsmill VJ. Post-extraction remodelling of the adult mandible. Crit Rev Oral Biol Med.1999;10(3): 384-404.
- [13] Ishihara Y, Nishihara T, Kuroyanagi T, Shirozu N, Yamagishi E, Ohguchi M, et al. Gingival crevicular interleukin 1 and interleukin 1 receptor antagonist levels in periodontally healthy and diseased sites.
- [14] Abbas AK, Lichtman AH, Pillai S. Cellular and Molecular Immunology. Saunders Elsevier 6th ed.2007.
- [15] Proff P, Romer P. The molecular mechanism behind bone remodelling: a review. Clin Oral Invest. 2009;13(4): 355-62.
- [16] Carranza FA, Henry HT, Michael GN. Clinical periodontology .9th ed. WB. Saunders Co. Philadelphia.2012.
- [17] Grieve WG, Reinhardt RA, DuBois LM. Prostaglandin E and interleukin 1 beta levels in gingival crevicular fluid during human orthodontic tooth movement. Am J Orthod.1994;369-74.
- [18] Kurtis B, Tuter G, Korkmaz T, Yucel A, Serdar M, Ozcan G. Clinical examination and interleukin 1 beta levels in gingival crevicular fluid in patients treated with removable partial dentures. J Prosthet Dent.2009; 102(1):62-70.
- [19] Seifi J. The effect of prostaglandin E2 and calcium gluconate on orthodontic tooth movement and root resorption in rats. J. Medical Sciences.2003;5-18.
- [20] Rodan GA, Yeh CK, Thompson DD. Prostaglandins and Bone. The biology of orthodontic tooth movement. CRC Press. Boca Raton.1989;263-8.