

Nephrotoxic effects of Nitrogen Dioxide Gas Exposure in Albino Rats

Shakti Mishra¹, Asha Agarwal², Rafat Khan³

Department of Zoology, School of Life Sciences, Khandari Campus, Dr. B.R. Ambedkar University, Agra 282002, India

Abstract: In the present study albino rats *Rattus norvegicus* (Berkenhout) were exposed to 40 ppm and 80 ppm nitrogen dioxide gas for 30 and 60 days for one hour per day. A decrease in serum creatinine, urea, potassium (K⁺) ion, protein and increase in serum sodium (Na⁺) ion levels after prolonged exposure to nitrogen dioxide gas. This study indicates that toxic effects of nitrogen dioxide gas exposure causes nephrotoxicity in albino rats.

Keywords: Nitrogen dioxide, serum protein serum creatinine, serum urea, serum sodium (Na⁺) ion and serum potassium (K⁺) ion

1. Introduction

Nitrogen dioxide considered as a serious air pollutant. Toxic effects of nitrogen dioxide usually occur after the inhalation of the gas beyond the threshold limit value (update of WHO, 2008) Nitrogen oxides are released to the air from the exhaust of motor vehicles, the burning of coal, oil or natural gas and during processes such as welding, electroplating, engraving and dynamite blasting, burning a lot of wood or use of kerosene, heater and gas stoves. Nitrogen dioxide causes oxidative stress in the body by inducing the generation of free radicals. Blood parameters are an asset in diagnosis the structure and functional status of body organs to toxicants (Bansal et al., 1979). The kidney functions may be assessed from the level of serum potassium (K⁺) ion, sodium (Na⁺) ion, urea, creatinine and protein in the serum (Nwankwo et al., 2006 and crook, 2007).

2. Materials and Methods

The albino rat, *Rattus norvegicus* (Berkinhout) of both the sex has been selected for the present investigation. The colony of the albino rats was inbred at the animal house of zoology Department .60 healthy and adult rats ranging in weight from 100 to 120g were kept in polypropylene cages at temperature 25°C ± 5°C, relative humidity 60± 5% and photoperiod 12hrs/day. The rats were fed on commercial food pellets (Golden feed, New Delhi) and water ad libitum. The Nitrogen dioxide gas was prepared by saltzman method (saltzman, 1954) and modified by levaggi et al, 1972. Nitrogen dioxide generator was used for the generation of nitrogen dioxide gas and rats were exposed in fumigation chamber (model AP-07, SFC-120) manufactured by standard Appliances, Varanasi.

The experimental rats were grouped in five sets. One control set 'A' and four experimental sets 'B', 'C', 'D' and 'E' of twelve rats each. Control set 'A' while experimental set 'B' was exposed to 40 ppm nitrogen dioxide gas and experiment set 'C' was exposed to 40 ppm nitrogen dioxide gas with supplementation of antioxidant vitamin C (5mg/rat/day) for 30 and 60 days for one hour per day in the fumigation chamber. Five rats were taken after 40 days and remaining five after 60 day from control (A) and experimental sets (B, C, D and E) the rats were sacrificed.

3. Result and Discussion

Alteration in the level of serum protein, urea, creatinine, sodium (Na⁺) and potassium (K⁺) ions is the indication of nephrotoxicity due to nitrogen dioxide gas exposure in albino rats. The results of this study are shown in table:1.

A decrease in serum protein, urea and creatinine is the result of inflammatory action of the nitrogen dioxide gas, causes influx of protein from serum to the site of oxidant injury in albino rats (shimizu et al, 1986; Muller et al, 2001; Agarwal et al, 2006). Serum urea and creatinine are the end products of protein metabolism which is processed in the liver, so the concentration of urea, creatinine depends mainly on the metabolism of protein and amino acids (Varley et al, 1980; Pant, 1999). An increase in level of serum sodium (Na⁺) ion with decrease in potassium (K⁺) ion in rats is the result of toxic action of nitrogen dioxide gas. Kidney maintain the water electrolytes and acid base balance (Nicopon and Hejlasz, 1991) Hypokalemia condition due to alkalosis is caused by the kidney response to alteration in serum potassium and sodium ion (Suga et al., 2001).

Another explanation regarding ionic disorder is the respiratory distress due to the air pollution which stimulates respiratory centre and disturbs the acid base balance in the body and respiratory alkalosis develop due to air pollutants (Oehme et al., 1996).

References

- [1] Bansal, S.K., Verma, S.R., Gupta, A.K. and Dalela, R.C., 1979, physiological dysfunction of the haemopoietic system in a fresh water teleost. Labeo rohita, following chronic chlordane exposure. Part 1. Alteration in certain haematological parameter. Bull Environ. Contam. Toxicol. 22:666-673.
- [2] Nwankwo, E.R., Nwankwo, B., Mubi, B., 2006. Prevalence of impaired kidney in hospitalised hypertensive patients in maiduguri Nig. internet J. Int. Med., 6(1).
- [3] Crook, M.A., 2007. The kidney In: Clinical chemistry and metabolic medicine, 7th Edi. Book power Britain pp. 36-57.

[4] Saltzman, B.E., 1954. colorimetric micro alteration of nitrogen dioxide in the atmosphere. Anal chem., 26: 1949-1950.

[5] Levaggi, D.A., S. Wayman and M. Feldstein, 1972. Method for the production of nitric oxide. Environ. Sci. Technol., 6: 250.

[6] Agarwal, A., K. Sharma and S. Mishra 2006. Effect of nitrogen dioxide inhalation on Lung total protein and serum total protein of albino rat. Asian. Jr. of Microbial Biotech. Env. Sci., 8(1): 143-145.

[7] Muller, B., M. Oske. R. Mochscheid, C. Seifer. P.J. Barth. H. Garh and P. Von wichest, 2001. Effect of N-acetylcysteine treatment on NO₂ impaired type 11 pneumocyte surfactant metabolism. European journal of clinical investigation 31 : 179-188.

[8] Shimizu, T., H. Sotokawa, M., Hatama, M., Izumiyama, H. Otoma and K. Kogure, 1986. Effect of NO₂ inhalation on protein and lipid accumulation in lung. Environ Res., 26: 422-437.

[9] Varley, H., A.H. Gowenlock and B. Mausice, 1980. Practical clinical Biochemistry 5th Edition William Hein Mann medical books Ltd. London, 1277 pp.

[10] Pant, M.C., 1999. Essential of biochemistry, Kedar Nath Ram Nath and Co. Meerut, 8th Edition pp 581.

[11] Suga, M1. Phillips, P.E., Ray, J.A., Raleigh, K.L., Gordon and R.J. Johnson, 2001. Hypohalemia induces renal injury and alternations in vasoactive mediators that favor salt sensitivity. Am. J. Physiol. Renal Physical., 28 : 620-629.

[12] Nicpon, J. and Z. Hejlasz. 1991. The effect of metabolic alkalosis on colostrum and milk quality of cows and on the health status on their newborns. DTW Dtsch. Tier arztl. Wochenschs 98 (6) : 207-9

[13] Oehme, F.W., R.W. Coppock, M.S. Mostrom and A.A. Khan, 1996. A review of the toxicology of air pollutants. Toxicology of chemical mixtures. Vet. Human. Toxicol., 38 (5) : 371-337.

[14] Update of WHO air quality guidelines on Cancer. June 2008 Retrieved August 1, 2015.

Table 1: Serum potassium (K⁺) ion (meq/l), serum sodium (Na⁺) ion (meq/l), serum creatinin, serum urea, serum protein in albino rats after exposure to nitrogen to nitrogen dioxide gas

Parameters	Exposure days	Control sets (S)	Experimental sets (S)	
		Control Sets	Experimental Sets(Concentrations)	
		Range	40 ppm Range	80 ppm Range
Serum Potassium (K ⁺)		Mean ± S.Em	Mean ± S.Em	Mean ± S.Em
	30	(6.0 - 7.5) 6.90 ± 0.29	(5.6 - 6.5) 6.04 ± 0.15 ^S ↓	(5.2 - 6.3) 5.92 ± 0.19 ^S ↓
	60	(6.5 - 8.0) 7.18 ± 0.26	(5.0 - 6.0) 5.68 ± 0.19 ^{H.S} ↓	(5.9 - 6.0) 5.26 ± 0.19 ^{V.H.S.} ↓
Serum Sodium (Na ⁺)	30	(110 - 121) 117 ± 2.03	(120 - 128) 122.60 ± 3.29 ^S ↑	(122 - 126) 124.00 ± 0.70 ^{H.S} ↑
	60	(110 - 116) 113.40 ± 1.08	(166 - 119) 117.40 ± 0.51 ^{H.S} ↑	(116 - 120) 118.00 ± 1.58 ^{V.H.S.} ↑
		(32 - 40) 37.20 ± 1.60	(22 - 36) 31.20 ± 2.90 ^{N.S} ↓	(22 - 35) 29.40 ± 2.10 ^S ↓
Serum Urea	30	(32 - 40) 36.40 ± 1.44	(22 - 36) 27.00 ± 2.60 ^{H.S} ↓	(22 - 36) 26.80 ± 2.60 ^{H.S.} ↓
	60	(1.0 - 1.5) 1.22 ± 0.12	(0.5 - 1.2) 0.72 ± 0.14 ^S ↓	(0.4 - 1.1) 0.72 ± 0.13 ^S ↓
		(1.0 - 1.5) 1.24 ± 0.10	(0.5 - 1.0) 0.76 ± 0.10 ^{V.H.S} ↓	(0.4 - 0.8) 0.60 ± 0.07 ^{V.H.S.} ↓
Serum Creatinine	30	(6.5 - 7.3) 7.04 ± 0.17	(5.9 - 6.9) 6.26 ± 0.19 ^{H.S} ↓	(5.8 - 6.5) 6.04 ± 0.12 ^{V.H.S.} ↓
	60	(6.9 - 7.6) 7.12 ± 0.13	(5.9 - 6.9) 6.18 ± 0.18 ^{V.H.S} ↓	(6.0 - 6.9) 6.18 ± 0.19 ^{V.H.S.} ↓