

Processing of Thalipeeth Flour-Blend from Various Cereal-Pulse Malt Flour and Its Quality Evaluation during Storage

Ranjeet Chunilal Kokani¹, T. V. Ranganathan²

M.Tech (Food processing & Engg), Karunya Institute of Technology & Science, Karunya University Coimbatore-641114

Guide, Karunya Institute of Technology & Science, Karunya University Coimbatore-641114

Abstract: *Thalipeeth is Maharashtrian ethnic food which is usually consumed during snack with curd or ghee. It is made by combination of pulses and cereals which provide nutritionally balance amount of nutrient and protein. Malting is process which improves the minerals and vitamins such as B₁, B₂, B₆, folic acid and vitamin C and reduces anti nutritional factors like phytate, tannins and oxalates. In this study flours of wheat, rice, Bengal gram dhal, black gram dhal sorghum, soybean and ragi (malted) was taken in equal proportion(i.e. 1:1:1:1:1:1), thalipeeth was made with the traditional method and packed in to standy pouches and kept under ambient condition i.e. at 25 °C+/-2 and at frozen condition i.e. - 4°C+/-2. The storage study was conducted for 60 days. From the study, it is concluded that the quality of ready to eat thalipeeth at ambient as well as at frozen condition were found to be remained in good condition till the last day of analysis (60th day) with respect to the quality characteristics like sensory, microbial and proximate parameters. However, the frozen product was highly superior in terms of high moisture retention and other quality parameters as compared to ambient stored sample. There were no significant difference observed during 60 days of storage study and therefore, this can be serving as good RTE product for people and it can be consumed at any time either after heating or as such after opening the packet.*

Keywords: Thalipeeth, Malting, combination of pulses and cereals, ambient condition, frozen condition, standy pouches

1. Introduction

In India majority of food consumption is still at home, nevertheless out of home food consumption is increasing due to urbanization, breaking up of traditional joint family system, desire for quality, time which translate into a increased need for convenience, increased number of working women, rise in per capita income, changing lifestyle and increasing level of affluence in middle income group has brought about changes in food habits. In India, consumption of ready to eat and ready to cook food has been significantly increase amongst the age group of 25-40 years (comprises about 60% of population irrespective of affordable price). In India, consumption of cereals and pulses is very high and cereals and pulses contain some antinutritional factors which could affect the digestibility of the food and therefore malting is one of the method used for cereals and pulses to reduce antinutritional factors and it improves digestibility, increases nutrient content. Malting is controlled germination process which activates the enzymes of the resting grain leads to conversion of cereal starch to fermentable sugar and partial hydrolysis of cereal protein.

Thalipeeth is made by using Cereals and pulses in combination. Cereals and pulses are the most important staple diet of the population all over the world. Cereals are cheapest source calories and provide adequate nutrient while pulses provide adequate amount of protein and other nutrients therefore, combination of cereals pulse provide balance diet. Thalipeeth is a Maharashtrian ethnic product and consume during snacks. Thalipeeth is multigrain thin unleavened pancake generally consume along with curd or chutney. Generally a mixture of wheat, rice, Bengal gram dal, black gram dal, ragi, sorghum used for flour of

thalipeeth. During making of thalipeeth, Mixture of grains first clean and then roasted for 10-15 min to obtained desirable pleasant flavour. Resultant flour is mixed with fresh spices like chopped onion, garlic, and coriander, spices powder like ajwain, cumin, chilli powder, turmeric, and asafoetida as per the taste, further water is added in the mixture and then it is kneaded into a smooth sticky dough unlike bhakari, a small dough piece is rolled on wet piece of cloth(thickness of 2-3mm in round shape) and transferred to hot pan seamer with oil, then it is the covered for 3-5 min and few holes are punched to facilitate the steam to escape out.

2. Materials and Methods

The raw materials for the present study were chosen to be wheat, ragi, jowar, Bengal gram dhal, black gram dhal; rice, soybean, and oil, masala powder (dry chilli powder, turmeric powder, coriander powder, cumin powder, salt, and ajwain).The thalipeeth was prepared by standard traditional method. The basic preparation of thalipeeth includes dough preparation. For dough preparation raw ingredients (wheat, soybean, rice, Bengal gram dhal, black gram dhal, jowar, ragi was malted and used) was first soaked for overnight (24hrs) and then dried in the cabinet drier for 8hrs at 55°C and roasted till brown colour develop and ground into fine flour. By adding specific quantity of water and curd dough was kneaded with the hand. Dough kept for few minutes to improve its texture of final product. Dough was rolled into thickness of 2-3mm in round shape and transferred to hot iron pan (full flame) seamer with oil and then it was covered for 3-5 min and few holes are punched to facilitate the steam to escape out. Prepared product was stored in different packaging materials(retort pouch, LDPE and PP bags) and

stored at room temperature (25°C+/-2, RH 80%) and frozen temperature (-4°C+/-2, RH 90%). The preservatives used in present study were Sorbic acid- for the present study Sorbic acid used in dough and flour blend in concentration of 0.02%. Calcium propionate – sodium and calcium propionates are used mostly in the prevention of mold growth and rope development in baked goods. It was used in concentration of 0.05% in dough, Acetic acid – Sorbic acid is effective at lower P^H, so to improve activity of Sorbic acid added in thalipeeth, the P^H of the product was lowered by adding acetic acid and it was added in concentration of 0.05% in dough of thalipeeth as well as in flour blend. BHA additives –this antioxidant was used on the basis of fat content of the product .it is basically used in oil which is used for roasting in concentration of 0.001%. For the study various packaging material used like LDP, standy pouches, HDPE.

Treatment Details

- T₁ –Control -Wheat, Bengal gram dhal, rice, jowar , ragi, soybean (roasted) (as per traditional method)
- T₂ – Wheat + Bengal gram dhal + ragi + rice + jowar + black gram dhal(soaked)+ spices mixture
- T₃- Wheat (soaked) + rice (soaked)+jowar(soaked) +Bengal gram dhal (soaked)+ black gram dhal(soaked)+ soybean (soaked)+ malted ragi+ spices
- T₄ -Wheat flour (soaked) + rice (soaked) +jowar (soaked) + ragi malted +chana flour (soaked) + black gram dhal + soybean (soaked) +roasted + spices added.
- T₅- preparation of Thalipeeth from above flour blend
- T₆- As above soy defatted replace by soy flour +spices+ preparation of Thalipeeth.
- T₇- As above T₅ + addition of preservative + sorbate(0.2%)+calcium propionate+ acetic acid packaging and storage, quality evaluation

Standardization of proportion of flour in mixture

Treatments		
T₁ (equal amount of flour) wheat 20:rice 20: black gramdhal20: chana dhal20: ragi20: sorghum20: soy flour 20	T₂ (different proportion of flour) flour blend of wheat 30: rice 30: chana dhal 20: black gram dhal 20	T₃ (different proportion of flour) Flour blend of wheat 50: ragi 30: chana dhal 20

T₁, T₂, T₃ flour blends treatments used for making thalipeeth with addition of onion, chillies coriander and salt and compared with the readymade thalipeeth flour. Out of 3 trials taken first trial was most acceptable.

Standardization of amount of water and type of water to be added in the Flour

Treatments			
T ₁	T ₂	T ₃	
50 gm of flour blend	50 gm of flour blend	50 gm of flour blend	50 gm of flour blend
10gm of onion	10 gm of onion	10gm of onion	10 gm of onion
5gm of chillies	5gm of chillies	5gm of chillies	5gm of chillies
5 gm of coriander	5gm of coriander	5 gm of coriander	5gm of coriander
1tsp salt	1tsp of salt	1tsp salt	1tsp of salt
25ml of water	35 ml of water	25ml of water	35 ml of water
		Hot water	Tap water

From above treatments for amount of water T₁ was finalised while for type of water hot water was most acceptable.

Standardization of Various Ingredient of Making Thalipeeth

Oil (ml)	Chillies (gm)	Turmeric	Chilli powder	Coriander powder	Cumin powder	Curd	water	Salt (%)	Onions (gm)	Sesame seeds
2	5	1tsp	1 ½ tsp	1tsp	1tsp	10	25	1	10	1tsp
5	10	¼ tsp	2tsp	1 ½ tsp	1 ½ tsp	20	35	2	15	1 ½ tsp
8	15	1/8 tsp	¼ tsp	2tsp	2 tsp	30	45	3	25	2 tsp

For initial trials onions and chillies were used but for final bulk except onion and chillies other standardise ingredient used. And highlighted treatments were finalised.

Standardization of Dough Making

Treatments			
T ₁	T ₂	T ₃	T ₄
50 gm of flour blend	50gm of flour blend	50gm of flour blend	50gm of flour blend
25gm of onion	25gm of onion	25gm of onion	-
5gm of chillies	5gm of chillies	5gm of chillies	-
5gm of coriander	5gm of coriander	5gm of coriander	-
1 ½ tsp chilli powder	1 ½ tsp chilli powder	1 ½ tsp chilli powder	1 ½ tsp of chilli powder
¼ tsp of turmeric powder	¼ tsp of turmeric powder	¼ tsp of turmeric powder	¼ tsp of turmeric powder
1tsp of coriander powder	1tsp of coriander powder	1tsp of coriander powder	1 tsp of coriander powder
1tsp of cumin powder	1tsp of cumin powder	1tsp of cumin powder	1tsp of cumin powder
1tsp of salt	1tsp of salt	1tsp of salt	1 tsp of salt
35 ml of hot water	35 gm of curd	20gm of curd+25ml of hot water	20gm of curd + 25ml of water

Out of four trials taken for dough making T₄ was finalised for bulk preparation. Use of curd and hot water gives smooth and wet texture to the thalipeeth. Though 3rd trial turns out good but due to onions and other fresh materials there are chances of getting spoiled.

Standardized Recipe of Thalipeeth

Ingredients	Amount
Flour blend (equal proportion)	140gm
chilli powder	1 ½ tsp
Turmeric powder	¼ tsp
Coriander powder	1tsp
Cumin powder	1tsp
Salt	1tsp
Sesame seeds	1tsp
Curd	30gm
Water	25ml

3. Results and Discussion

For selecting the most suitable method for the bulk preparation of ready to eat thalipeeth screening of the data obtained by the sensory evaluation of 25 panel members was done. Highest scores obtained to recipe was finalised depend on its sensorial characteristics. For bulk preparation of ready to eat thalipeeth various trials were taken for selecting most suitable method of preparation. Traditional method was used with the some modifications in it. Pre treatment given to the grains for making flour blend like soaking, roasting, drying

and malting in case of ragi. Out of these treatments soaking, drying and roasting was finalised for wheat, rice, Bengal gram, black gram and soybean while malting was selected for ragi. Three trials were taken for standardization of flour mixture. Out of those three trials T₁ i.e equal proportions of flour was finalised for further study. Three trails were taken and type of water and amount was finalised accordingly, in case of type of water T₃ was finalised and T₁ was finalised for amount of water. Various ingredients like oil salt, chilli powder, turmeric powder, curd, cumin and coriander powder and water percent were finalised for final preparation of thalipeeth. 5% oil, ¼ turmeric powder, 1 ½ chilli powder 1tsp cumin- coriander powder, 20 gm curd, 35 ml water and 2 % salt was finalised for preparation of thalipeeth. Four trails were taken for dough making and out of four trails T₄ was finalised i.e. addition of curd and water which gives smooth and wet texture to the thalipeeth.

Raw Material Analysis

The raw material used for making thalipeeth were grains of wheat, rice, sorghum, black gram dhal, Bengal gram dhal and soybean which were subsequently roasted in laboratory and converted to fine flour using lab flour mill and pass through 60 mesh sieve. The flours were blended at various proportions (treatment) and most acceptable treatment was selected by using sensory panel of judges and then further used same treatment for bulk preparation and storage studies. The results of proximate analysis of the raw material are given in following tables.

Proximate analysis							
Parameter Ingredient	Moisture (gm)	Protein (gm)	Carbohydrate (gm)	Fat (gm)	Ash (gm)	Crude fibre (gm)	Energy (Kcal)
Wheat	12.20	12.10	72.00	1.70	2.70	1.90	351.30
Rice	13.70	6.80	78.50	0.50	0.60	0.20	346.00
Ragi	13.10	8.20	75.10	1.30	2.70	3.60	344.70
Sorghum	13.10	12.10	67.50	5.02	2.30	1.20	363.58
Bengal gram dhal	9.90	20.80	60.70	5.60	3.00	3.90	376.40
Black gram dhal	10.90	24.00	60.20	1.50	3.20	0.90	350.30
Soybean	8.50	43.70	24.10	19.50	4.60	3.70	315.60

Proximate analysis of different grains used in flour blend (on dry basis) per 100gm

For making thalipeeth raw material was procured from local market, ingredient require for making thalipeeth are wheat, rice, ragi, Bengal gram dhal, black gram dhal, sorghum and soybean which are of good quality .The wheat exhibited moisture 12.20 % protein 12.10 % carbohydrate 72% , fat 1.7% , ash 2.7 % while crude fibre 1.9% and energy 351.30Kcal. Rice exhibited moisture 13.7% protein 6.80 % ,carbohydrate 78.5%, fat 0.5%, ash 0.6% while crude fibre 0.2% and energy 346 Kcal. Ragi contain moisture 13.10% protein 8.20% carbohydrates 75.1%, fat 1.3% ash 2.7%

while crude fibre 3.6% and energy 344.70 Kcal. The sorghum exhibited moisture 13.10%, protein 12.10% carbohydrates 67.5%, fat 5.0%, ash 2.7 % while crude fibre 1.2% and energy 363.58%.Bengal gram dhal exhibited moisture 9.9%, protein 20.8% carbohydrates 60.7%, fat 5.6 % , ash 3% while crude fibre 3.9 % and energy 376.40 Kcal. black gram dhal exhibited moisture 10.9% protein 24%, carbohydrates 60.20% , fat 1.5%, ash 3.2% while crude fibre 0.9% and energy 350.3 Kcal.and soybean exhibited moisture 8.50 % , protein 43.7%, carbohydrates 24.1% , fat 19.5%, ash 3.2% while crude fibre 3.7% and energy 315.6 Kcal.

Proximate composition of malted ragi (on dry basis) per 100gm

Proximate analysis							
Parameters Ingredient	Moisture (gm)	Protein (gm)	Carbohydrate (gm)	Fat (gm)	Ash (gm)	Crude fibre (gm)	Energy (Kcal)
Malted ragi	12.67	10.30	74.10	1.14	2.10	3.18	347.86

From the above table it is observed that malted ragi exhibited moisture 12.67%, protein 10.3%, carbohydrate74.1%, fat 1.14%, ash 2.1% while crude fibre 3.18 % and energy 347.86 Kcal.

Physical processing characteristics of malted ragi flour

Sr. No	Characteristics	Finger millet malt flour
1	Germination characteristic	
a	Weight of raw finger millet	500 gm
b	Weight of soaked finger millet	835 gm
c	Washing grains	2-3 time
d	Soaking time	24 hours
e	Germination period	24-30 hours
f	Germination temperature	25-30°C
g	Drying temperature (cabinet dryer)	55°C
h	Drying period	8-10 hours
i	Weight of grains after germination	890gm
2	Milling characteristics	
a	Flour obtained	300gm (60%)
b	Bran portion (using 60 mesh sieve)	200gm (40%)
3	Flour characteristics	
a	Colour	Off white
b	Texture	Very fine
c	Appearance	Powdery

Malting is important process which improves the nutritive value of all ingredients and bioavailability of nutrient significantly. Ragi has been considered and acceptable malt flour use in various food preparations. During malting seeds are germinated in the controlled conditions and dehydrated and finally converted into fine flour which pass through 60 mesh sieve during this process of malting starch is converted into simple digestible sugar, stored proteins synthesized into amino acids, fatty acid in seeds is converted into essential fatty acids, mineral and vitamins synthesis, flavour compounds also developed and enzyme diastase become more predominant and leads to easy digestion of starch and carbohydrates. Thus as whole nutritive values improve due to malting and hence it is used in various food preparations.

Proximate analysis of defatted soybean flour (on dry basis) per 100 gm

Parameters	Moisture (gm)	Protein (gm)	Fat (gm)	Carbohydrate (gm)	Ash (gm)	Energy (Kcal)
Ingredients						
Defatted soybean flour	9.1	52.10	2.79	33.05	3.05	365.71

From the above table it is observed that defatted soybean flour exhibited the proximate composition i.e. moisture

9.1%, protein 52.10%, carbohydrate 33.05% mineral content was 3.05% while energy 365.71 Kcal.

Proximate analysis							
Parameters	Moisture (gm)	Protein (gm)	Fat (gm)	Carbohydrate (gm)	Ash (gm)	Crude fibre (gm)	Energy (Kcal)
Ingredients							
Flour blend	10.20	13.50	3.57	72.21	0.69	3.50	374.97

Proximate composition of most acceptable flour blend (on dry basis) per 100 gm* (wheat 20: rice 20: sorghum 20: ragi malted 20: defatted soybean 20: Bengal gram dhal 20: black gram dhal 20). Proximate composition of flour blend is given in 4.1.d, from the above table it is observed that flour blend exhibited moisture 10.20 %, protein 13.50%, fat 3.57%, carbohydrate 72.21%, ash 3.50 % while crude fibre 3.50% and energy 374.97 Kcal.

Proximate analysis of finished product:

Analysis of thalipeeth and flour blend was done on the basis of proximate analysis, microbial analysis and sensory quality analysis. 20 panel members has selected for the sensory analysis of the product stored for 60 days.

Changes in proximate composition of ready to eat thalipeeth stored at ambient condition*(25°C +/- 2)

Parameters	Day	0 th day	10 th day	20 th day	30 th day	40 th day	50 th day	60 th day	SE +/-	CD at 5%
Moisture (gm)		18.58	18.55	18.50	18.40	18.32	18.26	18.10	0.30	0.90
Protein (gm)		19.25	19.27	19.28	19.32	19.36	19.42	19.48	0.28	0.84
Carbohydrate (gm)		48.47	48.49	48.50	48.55	48.57	48.59	48.65	0.18	0.54
Fat (gm)		10.90	10.90	10.86	10.80	10.76	10.76	10.72	0.23	0.69
Ash (gm)		3.80	3.80	3.77	3.74	3.70	3.67	3.65	0.15	0.48
Crude fibre(gm)		3.10	3.05	3.60	3.50	3.40	3.10	3.40	0.22	0.66
Energy (Kcal)		368.10	314.60	314.76	368.32	368.88	368.88	369.00	0.9	2.7

The data on changes in proximate composition of thalipeeth (RTE) stored at ambient condition is presented in above table from the data it was revealed that proximate constituent like moisture was found to be 18.58 % on 0th day and that of on 60th day 18.10 %. In case of protein the values on 0th day was 19.25 % and that of on 60th day 19.48 %. Fat percent on 0th day was 10.90 % and on 60th day 10.72 %. Carbohydrates on 0th day 48.47 % while on 60th day it was

48.65%. Crude fibre on 0th day was 3.10 % and on 60th day 3.40 %. While energy on 0th day was found to be 368.10 kcal while on 60th day it was found to be 369.00Kcal.The deference in all above parameters i.e. on 0th day and 60th day were statistically found to be insignificant thus during the storage of thalipeeth (RTE) at ambient condition there was no significant changes were observed.

Changes in proximate composition of ready to eat thalipeeth stored at frozen condition.*(-4°C+/-2)

Parameters	Day	0 th day	10 th day	20 th day	30 th day	40 th day	50 th day	60 th day	SE +/-	CD at 5%
Moisture (gm)		18.58	18.55	18.51	18.47	18.43	18.39	18.35	0.23	0.69
Protein (gm)		19.25	19.28	19.31	19.35	19.37	19.39	19.42	0.22	0.66
Carbohydrate(gm)		47.47	47.48	47.50	47.55	47.59	47.62	47.67	0.20	0.60
Fat (gm)		10.90	10.92	10.94	10.97	10.99	10.99	11.00	0.11	0.33
Ash (gm)		3.80	3.82	3.85	3.87	3.89	3.91	3.93	0.10	0.30
Crude fibre (gm)		3.10	3.12	3.15	3.17	3.18	3.20	3.22	0.06	0.18
Energy (Kcal)		366.88	365.32	366.13	366.33	366.87	366.95	367.36	2.04	6.12

The data on changes in proximate composition of RTE thalipeeth stored at frozen condition is presented in table 4.2.b, from the data it can be revealed that proximate constituent like moisture was found to be 18.58% on 0th day while on 60th day it was 18.35%. In case of protein the value at 0th day was 19.25% and that of on 60th day was 19.42%. A carbohydrate on 0th day was 47.47% and that of on 60th day was 47.67%. Fat on 0th day was 10.90% while on 60th day was 11.00%. In case of ash on 0th day was 3.80% and that of on 60th day was 3.93%. Crude fibre on 0th day was 3.10% and on 60th day was 3.22% while energy on 0th day was 366.88 Kcal and on 60th day was 367.36Kcal.

The deference in all above parameters of RTE thalipeeth on 0th day and 60th day were statistically found to be insignificant. Thus, RTE thalipeeth on frozen condition were found to be in good condition and there were no significant changes under frozen condition storage.

However, in case of Jowar bhakari (M35-1) no significant difference was notice in moisture and other parameters after baking (Naikare and Kadam, 1982). In another studies carried out on Puran Poli storage studies at refrigerated condition, Kanitkar (2009)?? Observed that there were no significant changes in moisture as well as other components in product when it was stored under refrigerated condition

Microbial analysis

Microbial analysis of finished product: The microbial examination was conducted as per the standard procedures for the fresh product sample and stored sample of Thalipeeth.

Microbial analysis of ready to eat Thalipeeth stored at ambient condition*(25°C +/- 2)

Dilutions	Days	0 th day	60 th day
10 ⁻¹		2×10 ⁻¹	12×10 ⁻¹
10 ⁻²		3×10 ⁻²	13×10 ⁻²
10 ⁻³		3×10 ⁻³	10×10 ⁻³
10 ⁻⁴		4×10 ⁻⁴	12×10 ⁻⁴
10 ⁻⁵		4×10 ⁻⁵	13×10 ⁻⁵
10 ⁻⁶		5×10 ⁻⁶	15×10 ⁻⁶
10 ⁻⁷		6×10 ⁻⁷	12×10 ⁻⁷
10 ⁻⁸		5×10 ⁻⁸	10×10 ⁻⁸
10 ⁻⁹		6×10 ⁻⁹	16×10 ⁻⁹
10 ⁻¹⁰		7×10 ⁻¹⁰	15×10 ⁻¹⁰

*mean of three observations

Data on the microbial analysis of the prodcut was given in table 4.3.a. From the table 4.3.a, it is observed that the total plate count of RTE thalipeeth was on 0th day was 2×10⁻² Cfu/ml while on 60th day it was 15×10⁻¹⁰ Cfu/ml.

It means that the values indicate increasing trend of Cfu count in the product as the storage period advances, however the difference observed are statistically insignificant. Hence no significant change was observed in the product
 Cfu: Colony forming units

Microbial analysis of ready to eat thalipeeth stored at refrigerated condition.*(-4°C+/-2)

Dilutions	Days	0 th day	60 th day
10 ⁻¹		1×10 ⁻¹	10×10 ⁻¹
10 ⁻²		2×10 ⁻²	11×10 ⁻²
10 ⁻³		3×10 ⁻³	12×10 ⁻³
10 ⁻⁴		4×10 ⁻⁴	11×10 ⁻⁴
10 ⁻⁵		5×10 ⁻⁵	13×10 ⁻⁵
10 ⁻⁶		6×10 ⁻⁶	15×10 ⁻⁶
10 ⁻⁷		5×10 ⁻⁷	12×10 ⁻⁷
10 ⁻⁸		6×10 ⁻⁸	10×10 ⁻⁸
10 ⁻⁹		6×10 ⁻⁹	16×10 ⁻⁹
10 ⁻¹⁰		7×10 ⁻¹⁰	15×10 ⁻¹⁰

*mean of three observation

Data on the microbial analysis of the prodcut was given in table 4.3.b. From the table no 4.3.b it is observed that the total plate count of RTE Thalipeeth on 0th day was 1×10⁻¹ Cfu/ml while on 60th day 15×10⁻¹⁰ Cfu/ml. It means that the values indicate increasing trend of Cfu/ml count in the product as the storage period advances, however the difference observed are statistically insignificant. Hence no significant change was observed in the product
 Cfu: Colony forming units

Sensory analysis: Sensory analysis of the thalipeeth was done by using 5 point hedonic scale. 20 panel members were employed for this sensory evaluation. The product is evaluated on the basis of its parameters as colour, texture, flavour, apperance and overall acceptability

Sensory analysis of ready to eat thalipeeth at ambient conditon *(25°C +/- 2)

Parameters	Storage Days	0 day	10 day	20 day	30 day	40 day	50 day	60 day
Colour (5)		4.23	4.50	4.18	4.02	4.19	4.20	4.20
Texture (5)		4.23	4.30	4.17	4.14	4.23	4.12	4.12
Flavour (5)		4.17	4.16	4.18	4.56	4.29	4.20	4.20
Appearance (5)		4.11	4.21	4.32	4.50	4.05	4.05	4.05
Overall acceptability (5)		4.59	4.21	4.21	4.26	4.3	4.3	4.3
Means of mean		4.27	4.28	4.21	4.30	4.21	4.17	4.17
Rank		1	1	1	1	1	1	1

*Means of three observations obtained by 20 panel judges using 5point hedonic scale
 Keys: 5-excellent, 4-very good, 3- good, 2-modarate, 1-poor

Changes in sensory analysis of ready to eat thalipeeth at refrigerated condition *(-4°C+/-2)

Storage Days	0 th day	10 th day	20 th day	30 th day	40 th day	50 th day	60 th day
Colour (5)	4.25	4.55	4.18	4.22	4.19	4.55	4.5
Texture (5)	4.23	4.3	4.17	4.14	4.23	4.2	4.24
Flavour (5)	4.17	4.16	4.18	4.56	4.29	4.18	4.20
Appearance (5)	4.11	4.21	4.32	4.5	4.05	4.9	4.18
overall acceptability (5)	4.59	4.21	4.21	4.26	4.3	4.21	4.6
Mean	4.27	4.28	4.21	4.33	4.21	4.40	4.34
Rank	1	1	1	1	1	1	1

*means of three observations obtained by 20 panel judges using 5point hedonic scale

Keys: 5-excellent, 4-very good, 3- good, 2-moderate, 1-poor

Costing of raw material

Ingredient	Amount	Rate / Kg (ruppe)	Price (rupee)	Manufactured by
Wheat	120	22	3	Local market
Rice	120	36	4.3	Local market
Black gram dhal	120	55	7	Local market
Bengal gram dhal	120	61	7.3	Local market
Soybean	120	56	7	Local market
Ragi	120	28	3.36	Local market
Sorghum	120	41	5	Local market
Chilli powder	10	120	1.2	Local market
Turmeric powder	5	190	0.95	Local market
Cumin powder	10	170	1.7	Local market
Curd	500	100	50	Local market
Coriander powder	10	430	4.3	Local market
Oil	30	80	2.4	Local market
Salt	10	20	0.2	Local market
BHA	0.002	1,600	0.003	Local market
Sorbic acid	0.4	4300	1.72	Local market
Acetic acid	0.1	650	26	Local market
Calcium propionate	0.1	2750	0.27	Local market
Labelling cost	1	3	3	Local market
Standby pouch	10	160	16	Local market
Total			145	

Note: The above table shows the costing for final bulk preparation of ready to eat Thalipeeth i.e. for 1.3 Kg which had a total cost of Rs. 145.00

Calculation:

a. 1.3 kg of total preparation cost for Rs.145.00
 Therefore 150 gm of net product cost is Rs. 17 which a cost price without addition of any profit

Addition of over head charges (40%)	17 × 40/100	7.0
Addition of profit (20%)	17 × 20/100	3.4
Addition of taxes (if any 12.5%)	17 × 12.5/100	2.1
Addition of returns (5%)	17 × 5/100	0.8
Addition of profit (15%) (dealers concession + distribution cost)	17 × 15/100	2.5

Addition of above charges to the cost price we get a MRP price of,

$$15.8 + 16 = 32.8 \text{ (approx Rs 33)}$$

$$\text{MRP} = 33/-$$

4. Summary and Conclusion

Thalipeeth is Maharashtrian ethnic product and made up of cereals- pulses combination. In India cereals and pulses are the staple diet of the population. In today's world everyone

is busy and therefore there is need of ready to eat or instant food because of increase need of convenience.

The main objective of the study was to improve the shelf life of the product. This snack is widely available in market but no attempts had been taken to development of thalipeeth or extending its shelf life. Thalipeeth is prepared with combination of cereals and pulses which provide balance amount of protein and other nutrients. Malting is another method used to improve the nutritive value of the product. In this study malting is done for ragi, because it's most acceptable and palatable after malting. Malting decreases anti nutritional factors such as phytate, tannins and oxalates at the same time increases various minerals and vitamins such as B₁, B₂, B₆, folic acid and vitamin C and amylases. Therefore malting is used in case of ragi.

Thalipeeth is multigrain thin unleavened pancake generally consume along with curd or chutney. Generally blend of flours like wheat, rice, bengal gram dhal, balck gram dhal, ragi, sorghum used for making thalipeeth by traditional method. Mixture of grains first clean and then roasted for 10-15 min to obtained desirable pleasant flavour. After grinding resultant flour is mixed with fresh spices like chopped onion, garlic, and coriander, spices powder like ajwain, cumin, chilli powder, turmeric, and asafoetida as per the taste, further water is added in the mixture and then it is kneaded into a smooth sticky dough unlike bhakari, a small dough piece is rolled on wet piece of cloth(thickness of 2-3mm in round shape) and transferred to hot pan seamer with oil, then it is the covered for 3-5 min and few holes are punched to facilitate the steam to escape out. In processing of thalipeeth various operation involved are cleaning grains soaking, drying, malting and finally roasting of grains, grind into fine flour. Flour mixture was blend in desirable proportion for preparation of thalipeeth by traditional method. Packaging of finished product in standby pouches which filling under laminar flow. Various treatments were planed with variation in flour blend (i.e four trials were taken), the thalipeeth was prepared from each flour blend (recipe) and after proper baking and cooling served to a panel of 20 judges for sensory evaluation . from the score card sheet most acceptable treatment was selected and further used in bulk preparation and same was packed and kept for storage study (at ambient and frozen conditions) thalipeeth preparation used as explained above.

References

[1] Arya S S,1984, Traditional Indian Foods- some recent developments, Def Sci J, Vol 34, No2: 173-182

- [2] Austin A and Ram A, 1971, studies on chapatti making quality of wheat, ICAR tech Bull (Agric) no 31, new Delhi
- [3] Antony U, Chandra, T S, 1998, Antinutrient reduction and enhancement in protein, starch and mineral availability in fermented flour of finger millet (*Eleusine coracana*) Journal of Agriculture and food Chemistry;46:2578-82
- [4] American soy association, Homepage: www.Asaasc.com
- [5] Arditi T, Meredith T and Flowerman P, 2000 renewed interest in soy isoflavons and saponins, cereal food world 45(9):414-417
- [6] Aykord W R and Doughty J, 1977, legumes in human nutrition, FAO food nutri ser, 12 V-138
- [7] Belitz H D and Grosch W, Schieberle, 2004, Food chemistry 3rd edn, Germany Springer :1070
- [8] Basu D, Kulirani F B, Ray D B, 2006, Agriculture, food security, nutrition and health in north-East India. Mittal Publications.
- [9] Brennan J G, 2006, Hand book of Food Processing, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany : 1-567
- [10] Borgstrom G, 1968, Principles of food science- food technology, Vol 1, Macmillan company, New York.
- [11] Chopra.S and Sankhala A ,2004, effect of soaking and germination on tannin, phytate and in vitro iron in underutilised legumes Horse gram and Moth beans, Journal of Food Sci Technol, 41, 5:547-550
- [12] Desai A D, Kulkarni S S, Sahoo A K, Ranveer R C and Dandge P B, 2010, Effect of supplementation of malted ragi flour on nutritional and sensorial quality characteristic of cake, Advance Journal of Food Science and Technology, 2(1):67-71
- [13] Dennis Edward Briggs, 1998, A book on Malts and Malting, Blackie and Professional pub, UK: 805.
- [14] Deak T and Novak E K, 1972, Assimilation of sorbic acid by yeasts, Acta alimentaria: 87
- [15] Fennema O R, 2008, Introduction to Food Chemistry, 3rd Edn, Marcel Dekker pub, New York :1067.
- [16] Fellows P J, 2009, Food Processing Technology principle and practice, wood head publishing LTD and CRC Press:
- [17] Frazier C W and Dennis C W, 2008, A book on Food Microbiology, 4th Edn, Tata McGraw- Hill pub, New Delhi: 513.
- [18] Friedman M and Brandon D L, 2001, Nutritional and health benefits of soy protein, Journal Agric food chem., 49, 3:1069-1086
- [19] Fukutake M, Takashashi M, Ishida K, Kawamura H, Sugimura T, Wakabayashi K, 1996, quantification of genistein and daidzein in soybean and soybean products, food and chemical toxicology ,34:457-61
- [20] Ganesh kumar K, Venkataraman L. V, Java T V, Krishnamurthy K S (1978) Changes in phytin, Ca^{++} , Mg^{++} and pectin, Journal of Food Science, 43(3):85-88
- [21] Glew R S, Lu Te Chuang, Roberts J L, Glew R H, 2008 Amino acid, fatty acid and mineral content of black finger millet, cultivated on the Jos Plateau of Nigeria, Food Chemistry, 2:115-18
- [22] Gopalan C, Sastri V B R and Balasubramanian S C, , 2004, Nutritive value of Indian Foods, National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India:161
- [23] Gorpade V M, Kadam S S and Salunkhe D K, 1986, Trypsin inhibitors in moth beans and horse gram, thermal stability and changes during germination and cooking, Journal of Food Science and Technology, in press
- [24] Geervani P and Theophilus F 1980, effect of home processing on the protein quality of selected legumes journal of food science ,45,3: 707-710
- [25] Hemamalini G, Umapathi K P, Rao T R , Sarawathi G, 1980, nutritional evaluation of sprouted ragi, Nutr Rep Int, 22:271-277
- [26] Jaya, T V, Krishnamurthy K S , Venkataraman, L V (1976), Effect of germination and cooking on the protein efficiency ratio of some legumes, Nutrition reports international 12(3): 175-183
- [27] Jood S, Mehta U, Singh R and Bhat C M, 1985, Effect of processing on flatus producing factor in legume, J Agri. food chem., 33: 266-271
- [28] Lestienne I, Icard-Vernière C, Mouquet C, Picq C, Trèche S, 2005, Effects of soaking whole cereal and legume seeds on iron, zinc and phytate contents, Journal of Food Chemistry ,89:3:421-425
- [29] Mella. O (Graduate Student), Department of Food Science and Technology and Researcher, Tanzania Food and Nutrition Centre Ministry of Health and Social Welfare EFFECTS (http://intsormil.org/smscientificpresents/2011ZambiaWorkshop/mella.pdf)
- [30] Malleshi N G and Deskiachar H S R, 1986, nutritive value of malted millet flour, Plant Food Hum Nutri 36: 191-196
- [31] Malleshi N G and Deskiachar H S R, 1982, formulation of weaning food with hot paste viscosity based on millet ragi and green gram, J Food Sci Nutr 44:193-197
- [32] Manay S N and Shadaksharaswamy M, 2008, Foods Facts and Principles, New Age International Pub, New Delhi:513
- [33] Messina M 1999, Legumes and soybeans: overview of their nutritional profiles and health effects, American journal of clinical nutrition, 70, 3: 4395-4505
- [34] Murthy H S , Patil D S and House L R, 1982, Studies on processing and cooking quality characteristics in sorghum , symp, on nutritional and processing quality of sorghum, All India Coordinated sorghum Improvement project workshop, May 17-19 Pune.
- [35] Naikare S M, Garad G N, Aher V P, Danagt S S and More H M, 2003, processing of sorghum into value added products, recent trends in millet processing and utilization: 19-27
- [36] **Nirmala M, Rao S and Muralikrishna G, 2000, Carbohydrates and their degrading enzymes from native and malted finger millet (Ragi, *Eleusine coracana*, Indaf-15), J Fd. Chem, 71 – (2) : 293**
- [37] Nirmala, M. and Muralikrishna, G, 2003, Three alpha amylases from malted finger millet (Ragi, *Eleusine coracana*, Indaf-15) purification and partial characterization, Journal of Phytochemistry, 62(1):21-23
- [38] Rao P U and Belavady B, 1978, Oligosaccharides in pulses varietal difference and effects of cooking and germination, Journal of Agriculture and Food Chemistry, 26:316-19

- [39] Ochanda, Oduor S, Akoth O Mwanjallah M, Joy O, Kagwiria and Mutiso.M ,2010,Effects of malting and fermentation treatments on group B-vitamins of red sorghum, white sorghum and pearl millets in Kenya, *Journal of applied biosciences*,34:2128-2134.
- [40] Pedersen B, Knudsen K E B, Eggnum B O, 1989, nutritive value of cereal products with emphasis on the effect of milling, *Wld Rev Nutr Dietet*, 60:1-91
- [41] Pawar, P A and Dhanvijay, V P , 2007, weaning foods An overview, *Beverage food world*,34:27-33
- [42] Pan. Z and Tangratanavalee W 2003, Characteristics of soybeans as affected by soaking conditions, *journal of Food Science and technology*,36,1: 143-151
- [43] Poter W L, 1980, Recent trends in food applications of antioxidants Auto oxidation in food and Biological System, Plenum Press, New York:295
- [44] Rai M, 2000, productivity grain focus. Survey of Indian agriculture. In: *the Hindu*: 53-55
- [45] Rehman Z and Shah W H (2004), Domestic processing effects on some insoluble dietary fibre components of various food legumes, *Journal of Food Chemisrty* ,78(4):613-617
- [46] Reddy N R,Balakrishnan C V and Salunkhe D K, 1978,phytate phosphorous and mineral changes during germination and cooking of black gram seeds,*J food Sci*;43,540
- [47] Reid D S, 1996, basic Physical Phenomena in the freezing and thawing of palnt and animal tissue in frozen food technology, C P Mallet London, Blackie academics and professionals : 1-19
- [48] Sankara Rao D S, Deosthale G Y ,1983, Mineral composition, ionisable iron and soluble zinc in malted grains of pearl millet and ragi, *journal of food chemistry*,11:3:217-223.
- [49] Sagar V. R and Suresh Kumar P, 2010, Recent advances in drying and dehydration of fruits and vegetables: a review, *J Food Sci Technol* 47(1):15–26
- [50] Scheneman J M, and Costilow R N, 1955, sorbic acid as preservative for sweet, *Appl. Microbiol*: 186
- [51] Salunkhe D K, Kadam S S and Chavan J K,1985, Postharvest Biotechnology of food legumes, CRS press,Inc :1-136
- [52] Saravacos G D and Kostaropoulos A E, 2006, Hand book of Food Processing Equipment, kulwar academic publisher: 1-698
- [53] Shewry P P, 2007, improving protein content and composition of cereal grains, *journal of cereal science* ,46:3:239-250
- [54] Singh M K, 2007, Food Preservation, discovery publishing house, New Delhi.
- [55] Slavin L J, Jacobs D and Marquart L, 2000, Grain Processing and Nutrition, *journal of food science and nutrition*,40:4:309-326
- [56] Stuckey B N, 1972, Antioxidants as food stabilizer , CRC Hand book of Food Additives, 2nd edition, the chemical rubber Co.CLeaveland Ohio: 185
- [57] Srilakshmi B, 2007,food science, new age international pub, New Delhi : 438
- [58] Thakare S K ,1982, nutritional and protein quality of wani sorghum, sym Proc.on Nutritional and processing quality of sorghum on 18th may 1982, Annual workshop at Pune.