

# Seasonal Variation in the Abundance of *Noctiluca scintillans* in the Coast of Arthunkal, Southwest Coast of India

Lekshmi, S<sup>1</sup>, Miranda, M.T.P.<sup>2</sup>

Department of Zoology, Fatima Mata National College, Kollam – 01, Kerala, India

**Abstract:** The seasonal variation in abundance of *Noctiluca scintillans* was investigated for one year during January – December 2015. A plankton net was used for collection and the collected samples were fixed in 4% neutral formaldehyde and placed in polythene vials for later sorting and identification. During the sampling period, the temperature and dissolved oxygen were recorded. The samples were examined under the microscope and *Noctiluca* cells were counted. High cell densities were recorded during September. No green or red water discoloration was observed. Increased number of *N. scintillans* cells could disrupt the traditional diatom-sustained food chain and it affects the health of an ecosystem.

**Keywords:** *Noctiluca scintillans*, seasonal variation, abundance

## 1. Introduction

*Noctiluca scintillans* is a non-photosynthetic, non-toxic and bioluminescent species of dinoflagellates. It is a bloom-forming marine species, commonly found in coastal and eutrophic areas of the Arabian Sea.

The incidence of *N. scintillans* blooms and discolorations of water in the Indian seas have been reported by Prasad (1953) in Palk Bay. The increased frequency of appearance of blooms throughout the world in the recent past, is a result of a change in the marine planktonic ecosystems, mostly due to the anthropogenic activities in the coastal zone. *N. scintillans* is also associated with fish kills due to enhanced levels of ammonia in the sea water (Elbrachter & Qi, 1998). *Noctiluca scintillans* is also a phagotrophic species feeding on bacteria, detritus, protozoans, copepods, eggs of copepods and fish as well as phytoplankton (Strom, 2001). This paper addresses the abundance and distribution of *N. scintillans* along the Coast of Arthunkal.

## 2. Materials and Methods

**Study Site:** Arthunkal coast (9°39'19" N; 76°17'23"E)



Figure 1: Study site

### Field sampling

The study was carried out along the coast of Arthunkal in Kerala, South West Coast of India. Sampling was carried out seasonally (January - December 2015) using a plankton net of mesh size 10 $\mu$ . Samples were taken from two stations (Station I and Station II) in the euphotic zone representing the nearshore region and 1km distance from the shore region respectively. The samples were fixed in 4% neutral formaldehyde and placed in polythene vials. They were examined under the microscope and the cells were counted. Subsurface water samples were collected in plastic bottles for subsequent chemical analysis. Dissolved oxygen and temperature were detected at the site itself. Dissolved oxygen was determined by Winkler's method and the temperature was recorded with a thermometer.

### 3. Results and Discussion

*N. scintillans* (Fig. 2) was observed with maximum cell densities in September. High abundance occurred at Station I and low abundance at Station II. The high cell concentration was found during monsoon (1200 cells/L) at station I and low concentration occurred during pre-monsoon. Seasonal variation in abundance of *N. scintillans* in Station I and Station II is shown in fig.3. No green or red water discoloration was observed. At station I, temperature recorded the highest value during pre-monsoon (30.5°C) and low value in monsoon (25.5°C). In the case of dissolved oxygen the maximum value was obtained at station I (5.8 mg/l) and minimum at station II (4.2mg/l). Seasonal variation of dissolved oxygen and temperature at Station I and II are given in Fig. 4 & 5. Increased number of *N. scintillans* cells could disrupt the traditional diatom-sustained food chain and it affects the health of an ecosystem.



Figure 2: *Noctiluca scintillans*




Figure 3: Seasonal variation in abundance of *N. scintillans* at Station I and II




Figure 4: Seasonal variation of dissolved oxygen and temperature at Station I




Figure 5: Seasonal variation of dissolved oxygen and temperature at Station II

### 4. Acknowledgements

We are extremely thankful to the Kerala State Council for Science, Technology and Environment for the award of a major research project under which this work was carried out.

### References

- [1] Prasad, R.R. (1953) Swarming of *Noctiluca* in the Palk bay and its effect on the Choodai fishery with a note on the possible use of *Noctiluca* as an indicator species. Proc. Indian Acad. Sci. 38 B (1): 40-47
- [2] Elbrachter, M. and Y.Z. Qi. 1998. Aspects of *Noctiluca* (Dinophyceae) population dynamics, In: Physiological ecology of harmful algal blooms. (Eds.): D.M. Anderson, A.D.Cambella and G.M. Hallegraeff. Springer, London, 662.
- [3] Strom, S. L. (2001) Light-aided digestion, grazing and growth in herbivorous protists. *Aquat. Microb. Ecol.*, 23 253–261.