

Added Effect of Graded Motor Imagery on Upper Limb Function in Stroke Patients

Rutu Parikh¹, Dr Sumitra Sakhawalkar², Dr Sayli Paldhikar³, Dr Snehal Ghodey⁴

¹Bachelor in Physiotherapy, Maeers Physiotherapy College, Talegaon Dabhade, Pune, Maharashtra. India

²Assistant Professor, Neurophysiotherapy Department / Maeers Physiotherapy College, Talegaon Dabhade, Pune, Maharashtra. India

³Professor, HODN Europhysiotherapy Department / Maeers Physiotherapy College, Talegaon Dabhade, Pune, Maharashtra. India

⁴Principal, Maeers Physiotherapy College, Talegaon Dabhade, Pune, Maharashtra. India

Abstract: GMI is a dynamic state during which the representation of a specific motor action is internally reactivated within working memory without any overt motor output. Imagery of movement activates largely the same brain areas that are activated when movement are actually performed. 10 subjects consisting of 7 males and 3 females were randomly allocated to GMI (n=5) (group A) and conventional (n=5) (group B) groups by chit method. Conventional group (group A) received conventional treatment consisting of range of motion exercises, task oriented exercises, balance and gait training etc. The GMI group (group B) received same treatment along with GMI training. Which included Implicit and explicit motor imagery and mirror therapy. Intervention was of 4 weeks, 5 times a week, 90 minutes per session. Outcome measures were- 1) Fugl Meyer assessment of sensorimotor function after stroke (FMA) 2) The Chedoke arm and hand activity inventory (CAHAI). Intra-group statistical analysis was done by paired t test which determined significant difference between the scores on both scales of both groups. Inter-group statistical analysis was done by unpaired t test which revealed significant difference in experimental group on FMA scale as compared to control group but no significant difference on CAHAI scale between both groups. **Conclusion-** upper extremity function shows significant improvement after graded motor imagery along with conventional therapy on FMA Scale whereas improvement in CAHAI scale is not significant.

Keywords: Graded motor imagery (GMI), Stroke, mirror therapy (MT), Upper limb function

1. Introduction

A stroke also called as cerebrovascular accident (CVA) is a rapidly developing loss of brain function due to disturbance in blood supply of brain caused by blocked or burst blood vessel. (1) WHO defines stroke as a neurological deficit of a cerebrovascular cause that persists beyond 24 hours. (2) There are two types of stroke are ischemic and hemorrhagic strokes. (1) Ischemic stroke is the most common type, affecting about 80 percent of individuals with stroke, and results when a clot blocks or impairs blood flow, depriving the brain of essential oxygen and nutrients. Haemorrhagic stroke occurs when blood vessels rupture, causing leakage of blood in or around the brain. (1)

Stroke is one of the leading causes of death and disability in India. (3) Stroke is no longer a disease of the developed world. Low and middle income countries like India account for 85.5% of total stroke deaths worldwide and number of disability adjusted life years in these countries was approximately seven times than the high income countries. (4)

Motor impairments frequently occur after stroke. It is estimated that after acute stroke approximately 80% of the patients have some form of motor impairment. About 20% of these patients regain at least part of their lost motor functions in the subsequent months; thus leaving 50-60% patients with chronic motor disorders. These disorders are often related to balance, timing and co-ordination, and to loss of strength and/or spasticity in the affected limbs. These motor impairments may substantially compromise quality of life after stroke. (5)

The most common deficit after stroke is hemiparesis of contralateral upper limb, with more than 80% of the stroke patients experiencing this condition. (6) The common manifestations of upper extremity motor impairment include paresis, muscle weakness, changes in muscle tone, joint laxity and impaired motor control. (7) These impairments lead to loss of upper limb function leading to activity limitation and participation restriction causing a decrease in quality of life. (8) Therefore, much therapeutic effort is invested in functional recovery of motor skills after stroke. This functional recovery is in the form of neurological recovery which is defined as recovery of neurological impairment and is often the result of brain recovery and reorganization. It has been increasingly recognized as being influenced by rehabilitation. Most spontaneous recovery occurs during the first 3 to 6 months of stroke. (9)

This brain reorganization occurs by a phenomenon called as neuroplasticity. Neuroplasticity is also known as brain plasticity or brain malleability is the ability of the brain to reorganize itself by forming new neural connections throughout the life. It allows the neurons in brain to compensate for injury and disease and to adjust the activities in response to new situations or changes in environment. (10)

Graded motor imagery (GMI) is a form of rehabilitation technique which uses the principles of motor recovery and neuroplasticity to promote graded cortical brain activation. (11) GMI is a dynamic state during which the representation of a specific motor action is internally reactivated within working memory without any overt motor output. (12) Imagery of movement activates largely the same brain areas

that are activated when movement are actually performed.

⁽¹³⁾ GMI consists of three steps-

- 1) Implicit motor imagery(IMI)-also called as left right discrimination. Consists of identifying whether a limb is right of left.
- 2) Explicit motor imagery(EMI)-consists of imagining a movement without actually performing it.
- 3) Mirror therapy(MT)-consists of a technique that uses visual feedback about motor performance using a mirror box. ⁽¹³⁾

GMI has been successfully used in treatment of persistent and complex pain states like-complex regional pain syndrome and phantom limb syndrome. ^{(14) (15)} abundant evidence is also available regarding practice of motor imagery in athletes and healthy individuals. ⁽¹⁶⁾ GMI is also helpful in the treatment of conditions like carpal tunnel syndrome, osteoarthritis, chronic neck and back pain. ⁽¹⁶⁾

In neurological conditions like stroke, GMI is considered as a “backdoor” to accessing the motor system and rehabilitation at all stages of stroke recovery because “it is not dependent on residual functions yet still incorporates voluntary drive.” ⁽¹⁷⁾

De vries and Mulder gave the concept that in stroke patients motor system can also be activated “offline” by motor imagery or observing movements. ⁽⁵⁾ The presence of mirror neurons goes hand in hand with this concept. These are neurons which fire both when we perceive (observe and imagine) the action as well as perform the action. ⁽¹⁸⁾

The above-mentioned concept shows that motor imagery, observation and execution are closely related phenomena sharing neural control processes. ⁽⁵⁾

Motor impairment after stroke is a major cause of permanent disability. Recovery of upper extremity is crucial in order to perform activities of daily living but is often variable and incomplete. It is proven that prolonged passive movement therapy in stroke patients neither improves performance nor induces cortical plasticity. And an early initiation of active movement becomes difficult due to lack of motor performance. Graded motor imagery(GMI) can be used to bridge this gap between passive therapy and active therapy. ⁽¹⁷⁾

Unlike active and passive motor therapies, graded motor imagery, in principle, is not dependent on residual function but still incorporates voluntary drive. Importantly, in the primate, primary motor cortex(M1) is directly involved in motor imagery as suggested by direct cellular recordings. In patients with stroke, motor imagery may therefore provide a substitute for executed movement as a means to activate the motor network. ⁽¹⁹⁾

.Many studies have been performed using motor imagery, mirror therapy and hand laterality recognition separately, but a few studies have been performed using all three together.

2. Review of Literature

1) Graded motor imagery for patients with stroke: a non-randomized controlled trial of a new approach.

PMID-27442717 Polli A et al European Journal of physical rehabilitation medicine 2017

This study included a total of 28 patients-14 in experimental group and 14 in control group with first ever unilateral stroke.

Duration was of 4week, 4 sessions per week of one hour each. Thus it included a total of 16 sessions.

Primary outcome measure-

- a) Wolf motor function test
- b) Fugl Meyer assessment

Conclusion: GMI is a feasible treatment for stroke patients with better outcome than conventional therapy.

2) Effect of graded motor imagery on upper limb motor functions and quality of life in patients with stroke: a randomized clinical trial.

Manisha Uttam, Divya Midha, Narkeesh Arumugam.

International Journal of therapies and Rehabilitation Research.

This study included a total of 26 subjects 13 in experimental and 13 in the control group with unilateral stroke within 1 to 6 months of onset.

Duration was of 5 weeks with 6 sessions per week of 1 hour each. Thus it included 30 sessions in total.

Primary outcome measure

- a) Fugl Meyer assessment scale (FMA)
- b) Chedoke arm and hand activity inventory scale

Secondary outcome measure- Stroke specific quality of life (SS-QOL)

Conclusion: GMI group along with conventional treatment shows more significant improvement than conventional group alone in improving upper limb motor functions and quality of life in patients with Stroke.

3) Mental practice with motor imagery in stroke recovery: randomized control trial of efficacy

Magdalena letsvaart, Marie Johnston, Brain-a journal of neurology. This study included 102 participants with 51 patients in experimental group and 51 in control group. Duration was of 4 weeks. 3 days of one to one supervised session(45min) plus additional 2 days’ home program(30min).

First 30 min were reserved for mental practice of actively imagining a variety of elementary movements and goal directed movements. Further 10 min was reserved for active motor imagery using mirror and video. Final 5 min for mentally rotating visual depiction of hands.

Primary outcome measure-action research arm test

Conclusion: No evidence of benefit of motor imagery in stroke no enhanced improvement as a result of motor imagery with mental practice.

4) Motor imagery training improves upper extremity performance in stroke patients

Seong –sik Kim, pt, PhD and byoung-hee-lee, pt, PhD. The journal of physical therapy science. It included 24 participants 12 in experimental group and 12 in control group.

Duration was of 4 weeks with 3 sessions per week of 30 minutes. Thus there were a total of 12 sessions total. Plus, conventional therapy 5 times a week

Primary outcome measure- Fugl Meyer assessment of upper limb function

Wolf motor function test

Conclusion: Motor imaging has a positive influence on upper extremity function.

5) Meta-analysis on the effect of mental imagery on motor recovery of hemiplegic upper limb function

Total of 6 trials were taken in study

Primary outcome measure-action research arm test.

Conclusion: 5 out of 6 studies yielded a positive finding in favour of mental imagery.

3. Methodology

Materials

- Laptop
- Mouse attached to laptop
- Mirror box
- Pen paper
- Fugl Meyer scale [FMAS]
- Chedoke arm and hand activity inventory. [CAAHAI]
- Movement imagery questionnaire [MIQ-R]
- And materials required in performing the scale

Method

Study design-experimental study

Sampling technique-purposeful simple random sampling

Sample size-10

Sampling population- patients with stroke within 6 months residing in Talegaon Dabhade.

Inclusion Criteria-Subjects between age 35 to 65.

Who have experienced one episode of unilateral stroke only.

Gender -Both males and females.

Type-Both ischemic and haemorrhagic strokes

Duration from episode-between 1 to 6 months

Mini mental state examination score >23

A score of 25 and above on Movement imagery questionnaire [MIQ-R]

Exclusion Criteria

Individuals with musculoskeletal disorder

Neurological disorder other than stroke

Visual impairment

Systemic diseases

Patient who is non cooperative

Patients with psychological problems

Outcome measures

1) FuglMeyer scale [FMAS]-Time frame-before and after 4 weeks. The upper extremity section of this scale was used. ⁽²⁴⁾

2) Chedoke arm and hand activity inventory. [CAAHAI] ⁽²⁵⁾
Time frame-before and after 4 weeks

Methodology-20 patients with acute stroke were screened for the inclusion criteria. 10 eligible patients who fulfilled all the inclusion criteria were included in the study. 5 patients allotted to experimental group and 5 patients allotted to control group randomly by chit method. Groups named A and B respectively.

Intervention-⁽²⁰⁾

Duration-

a) Total number of weeks-4 weeks

b) Sessions per week-5 sessions

c) Total number of sessions-20 sessions

Time allotted to each session-90min-60min of GMI 30min for conventional therapy

Conventional therapy for both group A and B-Passive range of motion exercises for upper limb, weight bearing exercises for upper limb, task oriented exercises for upper limb, positioning of upper limb, lower extremity conventional exercises, functional mobility training, balance and gait training.

Conventional therapy-





Graded motor imagery for experimental group (group A)

Graded motor imagery(GMI) has 3 steps (4)

- 1) Implicit motor imagery(IMI)
- 2) Explicit motor imagery(EMI)
- 3) Mirror therapy(MT)

Each step was introduced to the patients in a graded manner. Each step was progressed from easy to difficult level gradually. Steps 2 and 3 were introduced to the patient in the last two sessions of the preceding step.

a) IMI included a training based on hand discrimination task. An online software was used called as "orientate". It is a laterality recognition program which displays pictures of left and right hands randomly. It calculated the time consumed by the patient to guess the answer as well as recorded the progress of the patient over time.

60 pictures were displayed on a laptop screen and the patient had to identify whether the picture is of the right and or left hand.

A total of 7 sessions were conducted of which 5 were purely of IMI last two consisted of IMI and EMI.

Following target was aimed to be achieved at the end of this step-

- 1) Accuracy of 80% and above.
- 2) Similar results of left and right hands.
- 3) Response time of 2 seconds +/- 0.5 seconds.

b) EMI Training consisted of imagining movements without actually performing it. It was introduced in last two sessions of IMI.

- Standardized scripts were used to explain the task to the patient in detail.
- The scripts were translated to the language the patient was comfortable with by a professional.
- Initially easy tasks were asked to be imagined and then gradually the complexity of skill was enhanced.
- List of tasks to be imagined-
- To lift the arm
- To open a jar
- To pour a glass of water
- Draw a line with a ruler
- To button up a shirt
- Carry a bag up the stairs

Mirror therapy MT-using mirror box⁽¹³⁾


Is a technique that uses visual feedback about motor performance to improve rehabilitation outcomes.

It was introduced in the last two sessions of EMI.

It Involved the patient placing their affected hand and forearm inside the mirror box and their unaffected arm and forearm in front of the mirror

The patients were then instructed to perform a movement with their unaffected arm and to simultaneously attempt to copy the movement with their hidden affected arm^(8, 10)






4. Results


Graph 1 shows the pre and post mean scores of group A and group B on FMA with a mean difference of 11.6 and 5.2 respectively.

Graph 2 shows the pre and post mean scores of group A and group B on CAHAI with a mean difference of 8.8 and 5.8 respectively.



Graph 1


Group A-experimental group
 Group B-control group



Graph 2

Group A-experimental group
 Group B-control group

Graph 3 shows the comparison between group A and group B of the difference in means



Graph 3

Group A-experimental group

Group B-control group

Paired t test was used for intra-group statistical analysis of group A and group B. As shown in table 1 the p-value of group A for FMA was 0.0334 and for CAHAI was 0.0060. While that of group B for FMA was 0.0113 and for CAHAI was 0.0013. This shows that there was a statistically significant improvement in both groups on FMA and CAHAI after the treatment compared to the pre-treatment value ($p < 0.05$ for both groups at 95% confidence limit)

Paired t test			
Group	Scale	P Value	Significance
Experimental(Group A)	FMA	0.0334	Significant
Experimental(Group A)	CAAHAI	0.0060	Significant
Control(Group B)	FMA	0.0113	Significant
Control(Group B)	CAAHAI	0.0013	Significant

Unpaired t test was used for inter-group statistical analysis of group A and group B.

The mean difference in pre and post scores of experimental group for FMA and was 11.6 that for CAHAI was 8.8.

The mean difference in pre and post scores of control group for FMA and was 5.2 that for CAHAI was 5.8.

The post-treatment FMA score was statistically significantly higher in the experimental group than in the control group ($p = 0.0364$).

However, there was no statistically significant difference between the groups for the CAHAI score after treatment ($p > 0.05$), but there was a significant improvement compared to pre-treatment scores in both groups ($p < 0.05$ and $p = 0.001$, respectively).

Unpaired t test				
	FMA		CAHAI	
	Expt	Control	Expt	Control
Mean Difference	11.6	5.2	8.8	5.8
S.D	5.4589	1.6431	3.7013	1.6431
T Test Unpaired	2.5103		1.6565	
P Value	0.0364 Significant		0.1362 Not-Significant	

5. Discussion

The study was designed to see the added effect graded motor imagery on upper limb function in stroke patients. A sample of 20 was screened out of which 10 meeting the inclusion criteria were included in the study. These 10 subjects were divided randomly by chit method into two groups of 5 and were named as experimental group (group A) and control group (group B) respectively. Both groups were assessed with FMA and CAHAI before and after treatment. Both groups were given the same conventional treatment, additionally the experimental group (group A) was given graded motor imagery.

Based on the statistical analysis mentioned in the result above, current study showed statistically significant improvement in upper limb function on FMA and CAHAI after the treatment compared to the pre-treatment value in both the groups.

The results mentioned above revealed a statically significant improvement on the FMA scale for the experimental group as compared to the control group. However, there was no statistically significant difference between the groups for the CAHAI score after treatment.

The results obtained are similar to that of a study conducted by By Decety and colleagues. They revealed that there is involvement of the premotor area, supplementary motor area, cingulate area and parietal cortical area as well as basal ganglia and cerebellum not only during actual movement but also during imagination of movement. They proposed that graded motor imagery shares neural mechanism with processes used in motor control. They emphasized the importance of activation of the pre-frontal cortex during GMI and stated that this helps in maintaining dynamic motor representations in working memory. Thus they gave a general idea that prefrontal cortex is responsible for the creation and maintenance of explicit representations that guides action. This might have been one of the reasons for the improvement in upper limb function seen in our study.⁽¹³⁾

According to a systemic review conducted by Katleho Limakasto and colleagues in 2016 in south Africa Laterality recognition/implicit motor imagery tasks activate premotor and supplementary motor areas, with an exception of the primary motor cortex(M1) cortex. IMI is therefore fundamental in the preparation for subsequent phases of GMI programme. Thus this could have been one of the mechanisms how IMI contributed for the improvement of upper limb function in our study.⁽¹⁵⁾

In a study on motor imagery by Devries and Mulder in 1232 it was explained that Explicit motor imagery activates the somato sensory(S1), premotor and M1 cortices contralateral to the paretic extremity. Over the past decade, neuroimaging and psychophysical research on motor control has shown that there are striking similarities between real and imagined movements. These findings have led to a theoretical term called” the simulation hypothesis”. This hypothesis states that overt movement and motor imagery (covert movement) are essentially based on the same processes. Movement

execution, motor imagery and action observation are all driven by the same basic mechanism. Motor imagery and action observation are conceived as "offline" operations of the motor areas in the brain. The simulation hypothesis is based on 2 different lines of evidence.

First, it has been shown that there are similarities in the behavioural domain. For instance, the time to complete an imagined movement is known to be similar to the time needed for actual execution of that movement; this phenomenon is known as mental isochrony. Parsons showed that the time needed to judge whether a rotated picture of a hand is a left or a right hand is related to the degree of the rotation of that picture.

A second line of evidence for the simulation hypothesis shows that the neural system, used for action control is, indeed, activated during imagination of these actions. An increasing number of brain imaging studies have shown this similarity at the neural level.⁽⁵⁾

Moseley in the year 2004 conducted a study using GMI on patients with chronic CRPS1 and concluded that GMI reduced pain by 20 points on Numerical pain rating scale in the chronic CRPS1 population. Moseley conclude that patients in GMI group did better than patients in other group and the treatment components were only effective when they followed the sequential pattern. Moseley stated that CRPS1 involves cortical abnormalities similar to those observed in phantom limb pain and that after stroke. He thought that the possible explanations for the results obtained were sequential activation of cortical pre-motor and motor networks or sustained and focused attention on the affected limb, or both. Moseley made major contribution towards the development of GMI intervention strategy. There are many studies that look at the provision of one component of GMI process, but limited research is there looking at the whole process.⁽¹⁴⁾

On similar lines Manisha Uttam and colleagues hypothesised that even though CRPS1, Phantom limb pain and stroke are different conditions originating from distinct mechanisms of peripheral trauma, deafferentation and cortical damage respectively, they share identical aspects of symptomatic presentation and pattern of cortical reorganisation. Thus, based on these common findings the established mechanisms of GMI on CRPS1 and phantom limb pain can also be applicable to Stroke.⁽²¹⁾

Ramachandra was the first to study about mirror therapy. According to him mirror therapy works on the principle of mirror visual feedback(MVF). MVF addresses changes in the S1 and M1 cortices. In addition, it provides visual input to the brain, that movement is executed normally without inhibition. The therapeutic effect associated with mirror visual feedback may be due to activation of mirror neurons in the brain hemisphere contralateral to the paretic limb. These mirror neurons have been shown to fire during observation and execution of movement. Mirror neurons accounts for about 20% of all neurons present in a human brain. These neurons are capable of laterality reconstruction i.e., ability to differentiate between the left and right sides. When using a mirror box, it is found that these mirror

neurons get activated and helps in recovery of affected parts. This system is thought to use the observation of movement to stimulate the motor process which would be involved in the movement.⁽²²⁾

According to a study done by Rizzolatti and his colleagues in 2010 using mirror therapy on stroke patients, motor command neurons are found in abundance in the frontal lobes as well as the parietal lobes. These neurons fire to orchestrate a sequence of muscle twitches to produce skilled movement. According to Rizzolatti "mirror neurons" a subset of motor command neurons also fire when a person merely WATCHES another individual perform the same movement or IMAGINES the movement. Mirror neurons necessarily involve interactions between multiple modalities—vision, motor commands, proprioception—which suggest that they might be involved in the efficacy of MVF in stroke. An additional possibility is that lesion is not always complete; there may be a residue of mirror neurons that have survived but are 'dormant' or whose activity is inhibited and does not reach threshold. (And, indeed, motor areas may have become temporarily inactive as a result of the same mechanism as learned.

6. Conclusion

Thus we conclude that GMI along with conventional treatment shows more significant improvement than conventional therapy alone in improving upper limb function on FMA scale for stroke patients but no significant.

7. Scope of Study

- 1) The study can be performed on a large scale using a larger sample size.
- 2) Longer application and its effect can be seen.
- 3) Independent effect of GMI can be studied without combining it with conventional therapy.

8. Clinical Implication

- GMI is non-invasive, extremely feasible and easy intervention, which can be used as an adjunct to conventional therapy.
- It can also be used in any kind of therapeutic set-up and can also be recommended as home program.
- Lastly it can be used by patients irrespective of residual limb function.

References

- [1] Susan B osullivan(ptEdD) Thomas j Schmitz(ptphD) physical rehabilitation 5th edition 705 711 ISBN 81-8061 934-6.
- [2] PSM park 22nd edition.
- [3] Stroke epidemiology and stroke care services in India- jeyaraj Durai Pandian and Paulin Sudhan.
- [4] Stroke in India-Factsheet-Fiona C Taylor, Suresh Kumar K.
- [5] de Vries and Mulder 2017-motor imagery and stroke rehabilitation: a critical discussion.

- [6] Rehabilitation of motor function after stroke: A multiple systemic review focused on techniques to stimulate upper extremity Recovery-Samar M Hatem, Geoffrey Saussez.
- [7] Upper limb motor impairment post stroke-Preeti Raghvan MD.
- [8] Assessment of upper extremity impairment, function and activity following stroke: Foundations for clinical decision making- Catherine E Lang PT PhD.
- [9] Skilbeck CE wade DT Hever RL, wood VA recovery after stroke J neurosurgical psychaitry 1943 ;46:5-8
- [10] Seiz RJ freund HJ plasticity of the human motor cortex adv. Neurol 1997 73 321-333
- [11] Graded motor imagery for patients with stroke: A non-randomized controlled trail of a new approach- Polli A et al.
- [12] A standard motor imagery introduction program for neurological rehabilitation-wondrusch and Schuster.
- [13] Motor imagery, By Decety and colleagues.
- [14] Graded motor imagery is effective for long standing complex regional pain syndrome: A randomized control trail- G L Moseley.
- [15] The effects of Graded motor imagery and its components on phantom limb pain and disability in upper and lower limb amputees: A systemic review- Katleho Limakatso, Lieselotte Corten.
- [16] Motor imagery in physical therapy practice- Ruth Dickstein, Judith E Deutsch.
- [17] Motor imagery- A backdoor to motor system after Stroke-Nikhil Sharma.
- [18] Mirror neurons- how we reflect behaviour.
- [19] Georgopoulos AP LJ, Petrides M, Schwartz AB, Massey JT. Mental rotation of the neuronal population vector. Science. 1989; 243:234–236.
- [20] Moseley GL Butler DS Beams T Giles-the graded motor imagery handbook.
- [21] Effect of graded motor imagery on upper limb motor functions and quality of life in patients with stroke: a randomized clinical trial -Manisha Uttam*, Divya Midha, Narkeesh Arumugam.
- [22] The use of visual feedback, in particular mirror visual feedback, in restoring brain Function-V. S. Ramachandran and Eric L. Altschuler
- [23] The functional role of parieto-frontal mirror circuit: interpretations and misinterpretations-Rizzolatti G, Sinigaglia C
- [24] The Fugl-Meyer assessment of upper extremity:reliability, responsiveness and validity.- Camilla Biering Lundquist.
- [25] Test-retest reliability, validity and sensitivity of the chedoke arm and hand activity inventory-limb function for survivors of stroke. -Barreca SR, et al