A New Four-Parameter Generalized van der Waals Equation of State: Metastable State of Group IV Elements

R. Balasubramanian¹, G. Theertharaman²

^{1, 2}Department of physics, Arignar Anna Govt. Arts College, Namakkal 637 002, Tamilnadu, India

Abstract: A new four- parameter generalized van der Waals equation of state has been proposed and employed to calculate the spinodal (Metastable State) and the thermodynamic limit of superheat of the of silicon, germanium, lead and tin. It is established that silicon, germanium, tin and lead obey the single parameter law of corresponding states. It is established that the new parameter introduced in the attractive term of the equation of state is a thermodynamic similarity parameter. It has been established that silicon, germanium, tin and lead can be superheated, under rapid heating, up to temperatures 4481K, 7975K, 7161K, and 4333K, respectively. Above these temperatures, silicon, germanium, tin and lead undergo explosive boiling by virtue of homogeneous nucleation. This fact is to be taken into account when silicon, germanium, tin and lead are subjected to rapid heating.

Keywords: Equation of state, Law of corresponding states, Germanium, Lead, Silicon, Tin, Spinodal, Superheating

1. Introduction

The study of the thermodynamic properties of the silicon, germanium, tin, and lead is of scientific and technological significance. The experimental studies on the thermodynamic properties of the silicon, germanium, tin, lead in the metastable region, encounter severe difficulties. Thus, arises a need for theoretical studies on their thermodynamic properties. In recent years, several studies have been made [1-7] on the thermodynamic properties of silicon, germanium, tin and lead. This fact manifests the relevance of the study of the thermodynamic properties of the silicon, germanium, tin, lead. One of the Statisticomechanical and thermodynamical approaches to study the thermodynamic properties of substance is the development of equations of state for substances. To improve the accuracy, the known equations of state are generalized [8-20] by modifying the repulsive and attractive terms.

This work is aimed at developing a new equation of state for silicon, germanium, tin and lead in the metastable state. In this work, the known two–parameter generalized van der Waals equation of state is generalized by modifying its repulsive and attractive terms. The performance characteristics of generalized van der Waals equation of state in describing the properties of silicon, germanium, tin and lead in the metastable state are investigated.

2. Generalized van der Waalsequation of state

The known two – parameter generalized van der Waals equation of state does not precisely describe the thermodynamic properties of fluids. This may be attributed to the inaccurate repulsive and attractive terms in the generalized van der Waals equation of state of state. Hence, in this work, this equation is modified by introducing a parameters c in the repulsive term and n in the attractive term. Such a generalized van der Waals equation of state for one mole of substance has the form

$$P = \frac{RT}{V - b + c} - \frac{a}{V^n} \tag{1}$$

where P – Pressure, V - Molar volume, T – Temperature,R - Universal gas constant, and a,b, c, and n are substance-specific constants.

The vapor-liquid critical point conditions are

$$\left(\frac{\partial P}{\partial V}\right)_{T_c} = 0 \qquad ; \qquad \left(\frac{\partial^2 P}{\partial V^2}\right)_{T_c} = 0 \qquad (2)$$

From Eqs.(1) and (2), we get the critical volume, critical temperature and critical pressure as

$$V_c = N(b-c) \tag{3}$$

(4)

Where, $N \equiv \frac{n+1}{n-1}$

$$T_{c} = \frac{n(N-1)^{2} a}{RN^{n+1}(b-c)^{n-1}}$$

$$P_c = \frac{a}{N^{n+1}(b-c)^n} \tag{5}$$

When Eqs.(3) - (5) are taken into account, we get the critical compressibility factor as

$$Z_c = \frac{P_c V_c}{RT_c} = \frac{n^2 - 1}{4n} \tag{6}$$

The generalized van der Waalsequation of state may be rewritten in terms of the reduced variablesas

$$P_{s}^{*} = \frac{N}{V^{*n}} \left[\frac{n \left(V^{*} - \frac{1}{N} \right)}{V^{*}} - 1 \right]$$
(7)

Volume 7 Issue 4, April 2018 www.ijsr.net

Where,

$$P^* = P/P_c$$
, $V^* = V/V_c$, $T^* = T/T_c$

The reduced equation of state given by Eq.(7) represents the single-parameter law of corresponding states with the thermodynamic similarity parameter n. That is, substances obeying the generalized van der Waalsequation of state, with the same values of parameter n are thermodynamically similar. That is, substances, have similar intermolecular force characteristics.

3. Equation-of-state parameters

The parameters a,b,c and n of the generalized van der Waals equation of state are determined through the critical-point parameters. Eq. (6) is a quadratic equation with respect to the parameter n. the physically meaningful solution (i.e. n>0) of Eq. (6) is

$$n = 2Z_c + \sqrt{4Z_c^2 + 1}$$
(8)

Eq. (4) gives the parameter a of the generalized van der Waalsequation of state as

$$a = \frac{(n+1)^2 R T_c V_c^{n-1}}{4n}$$
(9)

Eqs. (3) gives the parameter b of the generalized van der Waalsequation of state as

$$b - c = \left(\frac{n-1}{n+1}\right) V_c \tag{10}$$

Using Eqs.(8)-(10), the parameters of the generalized van der Waalsequation of state can be determined. Moreover, using the Riedel's parameter along with the critical volume, the values of the parameter b and c can be determined.

4. Spinodal

The knowledge of the spinodal, a characteristic curve on the phase diagram, is essentialin describing the properties of a substance in the critical and in the metastable states. Fig 1 schemtically depicts [21] the vapour-liquid equilibrium curve (bimodal) and the stability boundary curve (spinodal) of substances.



The spinodal defines the thermodynamic Metastable Stateof the phase envelope. The spinodal encloses theregionof

unstable states for which the elasticity is negative.For stable states, the elasticity is positive.In the region between the binodal and the spinodal on the phase diagram,the liquid is in the metastable state.Considering the scientific and technological significance,in recent years,several studies have been made [22-32] on the behavior of the superheated metastable fluids. The spinodal is therefore, defined by the condition:

$$-\left(\frac{\partial P}{\partial V}\right)_T = 0 \tag{11}$$

Applying the condition given by Eq.(11) to Eq.(7), we get the equation of spinodal in T^* , V^* coordinates as

$$T_{s}^{*} = \left[\frac{\left(n^{2} - 1\right)\left(NV^{*} - 1\right)^{2}}{4NV^{*n+1}} \right]$$
(12)

Substituting Eq. (12) into Eq. (7), we get the equation of spinodal in P^* , V^* coordinates as

$$P_{s}^{*} = \frac{N}{V^{*n}} \left[\frac{n \left(V^{*} - \frac{1}{N} \right)}{V^{*}} - 1 \right]$$
(13)

With a decrease in pressure, the superheat of substances increases. The thermodynamic limit of superheat is attained at

Applying the condition given by Eq.(14) to Eq.(7) and using Eq.(12),we get

$$V_{s,0}^* = \frac{n+1}{n}$$
(15)

Where $V_{s,0}^*$ - The reduced volume of the fluid at the thermodynamic limit of superheat

$$T_{s,0}^* = \frac{1}{4} \left(\frac{n+1}{n}\right)^{n+1} (16)$$

Where $T_{s,0}^*$ - The thermodynamic limit of superheat

That is, thermodynamic limit of superheat depends only on the parameter n but not on the parameters a, b and c of the of generalized van der Waalsequation of state.

5. Determination of Equation-Of- State Parameters

The parameters of the generalized van der Waals equation of state can be determined using any characteristic point on the phase diagram. However, the use of the critical-point parameters in determining the equation of state parameters will improve the accuracy of the equation of state in describing the high-temperature properties of substances. The parameter nforsilicon, germanium, tin and leadis determined through the Eq. (8) using experimental data [33] on the critical compressibility factor. The obtainzd values of п are presented in Table1. The parameteraforsilicon, germanium, tin and leadis determined through the Eqs (9)using experimental data on criticalpoint parameters along with the values of n. The parameter

Volume 7 Issue 4, April 2018 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY b-c forsilicon, germanium, tin and leadis determined through the Eq.(10) using experimental data on criticalpoint parameters along with the values of *n*. The obtained values of *a* and *b*-care presented in Table 1.

Tuble If Equation of State parameters			
	$a \ 10^{5}$	b-c	п
Substance	$Pa(m^3.mol^{-1})^n$	$10^{-6}(m^3.mol^{-1})$	
Silicon	4.5476	6.1637	1.7285
Germanium	12.5332	11.7115	1.7195
Tin	13.6081	15.3236	1.7197
Lead	8.7167	16.8070	1.7160

Table 1: Equation-of-state parameters

6. Determination of Spinodal

Considering the values of n (Table1) forsilicon, germanium, tin and lead, the spinodal is determined by Eqs.(12) and (13).The obtained spinodal-parameters are presented in Tables2-5.These spinodal-parametersdefine the Metastable State of silicon, germanium, tin and leadin the phasediagram.

	1	
V_s^*	T_s^*	P_s^*
0.55	0.7619	-1.1641
0.6	0.8320	-0.3679
0.6335	0.8683	0
0.65	0.8846	0.1459
0.7	0.9237	0.4804
0.75	0.9521	0.6969
0.8	0.9722	0.8351
0.85	0.9858	0.9204
0.9	0.9944	0.9701
0.95	0.9988	0.9933
1	1	1.0170



Plot 1.1: Spinodal of Silicon in P-V Coordinates



Plot 1.2: Spinodal of Silicon in T-P Coordinates

Table 3: Spinodal of Germanium		
V_s^*	T_s^*	P_s^*
0.55	0.7644	-1.1372
0.6	0.8345	-0.3522
0.6323	0.8698	0
0.65	0.8863	0.1555
0.7	0.9247	0.4858
0.75	0.9527	0.6942
0.8	0.9724	0.8367
0.85	0.986	0.9211
0.9	0.9942	0.9696
0.95	0.9987	0.9933
1	1	1



Plot 2.1: Spinodal of Germanium in P-V Coordinates



Plot 2.2: Spinodal of Germanium in T-P Coordinates

Volume 7 Issue 4, April 2018 www.ijsr.net

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296

Table 4: Spinodal of Tin			
V_s^*	T_s^*	P_s^*	
0.55	0.7656	-1.1382	
0.6	0.8344	-0.3527	
0.6323	0.8697	0	
0.65	0.8863	0.1553	
0.7	0.9218	0.4857	
0.75	0.9527	0.6999	
0.8	0.9725	0.9031	
0.85	0.9860	0.9213	
0.9	0.9942	0.9697	
0.95	0.9987	0.9935	
1	1	1	







Plot 3.2: Spinodal of Tin in T-P Coordinates

Table 5: Spinodal of Lead			
V_s^*	T_s^*	P_s^*	
0.55	0.7671	-1.7291	
0.6	0.8356	-0.4988	
0.6318	0.8701	0	
0.65	0.8871	0.1593	
0.7	0.9252	0.4879	
0.75	0.9529	0.7012	
0.8	0.9726	0.8374	
0.85	0.9859	0.9215	
0.9	0.9943	0.9697	
0.95	0.9987	0.9934	
1	1	1	



Plot 4.1: Spinodal of Lead in P-V Coordinates



Plot 4.2: Spinodal of Lead in T-P Coordinates

7. Determination of thermodynamic limit of superheat

The volume at the thermodynamic limit of superheat forsilicon, germanium, tin and leadaredetermined through Eq.(15) using the values of the parameters n (Table1). The obtained values are presented in Table 6. The thermodynamic limit of superheat forsilicon, germanium, tin and lead is determined through Eq.(16) using the values of the parameters n (Table1). The obtained values are presented in Table 6.Below the thermodynamic limit of superheat, heterogeneous nucleation prevails. And, above the thermodynamic limit of superheat, homogeneous nucleation will prevail resulting in the explosive boiling of fluids.

 Table 6:
 Thermodynamic limit of superheat

Substance	$T^{*}_{s,0}$	$V_{s,0}^{*}$	T _{s,0} k	$V_{s,0}$ 10 ⁻⁵ m ³ /mol
Silicon	0.8683	0.6335	4481	1.4625
Germanium	0.8697	0.6323	7975	2.7989
Tin	0.8697	0.6323	7161	3.6615
Lead	0.8701	0.6318	4333	4.0280

8. Results and Discussion

The four-parameter generalized van der Waalsequation of state has been employed to calculate the spinodal, and thermodynamic limit of superheat ofsilicon, germanium,

Volume 7 Issue 4, April 2018 www.ijsr.net

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296

tin and lead. The performance characteristics of the generalized van der Waals type equation of state in evaluating the spinodal, and the thermodynamic limit of superheat ofsilicon, germanium, tin and leadhave been studied.The parameters of the generalized van der Waalsequation of state are expressed in terms of the critical-point parameters ofsilicon, germanium, tin and lead. Thus, is has been established that the three characteristic properties of the fluids viz., the critical pressure, the critical volume and the critical temperature characterize the generalized van der Waalsequation of state. It has been established thatsilicon, germanium, tin and leadcan be superheated, under rapid heating, up to temperatures 4481K, 7975K, 7161K and 4333K respectively. That is silicon can be superheated to above 943 K above their normal boiling temperatures. germanium can be superheated to above 4819 K above their normal boiling temperatures. tin, can be superheated to above 4302 K above their normal boiling temperatures and lead can be superheated to above 2310 K above their normal boiling temperatures. This fact is to be taken into account whensilicon, germanium, tin and leadare subjected to rapid heating.

9. Conclusion

A new four-parameter generalized van der Waalsequation of state is proposed for describing the high-temperature properties of silicon, germanium, tin and lead. It is established that silicon, germanium, tin and leadobey the single parameter law of corresponding states. It is established that the newly introduced parameter n is a thermodynamic similarity parameter. The spinodal (Metastable State on the phase diagram) of silicon, germanium, tin and leadhas been determined. The thermodynamic limit of superheat of silicon, germanium, tin and leadhas been determined.

References

- [1] A. Antonopoulos, et al., "Introducing a silicon carbide inverter for hybrid electric vehicles," in *Power Electronics Specialists Conference*, 2008. PESC 2008. IEEE, 2008, pp. 1321-1325.
- [2] J. Hornberger, et al., "Silicon-carbide (SiC) semiconductor power electronics for extreme hightemperature environments," in *Aerospace Conference*. *Proceedings*. 2004 IEEE, 2004, pp. 2538-2555 Vol.4.
- [3] B. J. Baliga, "Power semiconductor device figure of merit for high-frequency applications,"*Electron Device Letters, IEEE*, vol. 10, pp. 455-457, 1989.
- [4] L. Del castillo, Y. Chen, N. Aranki, C. Assad, M. Mazzola, M. Mojarradi, and E. Kolawa, "Reliability Assessment of High Temperature Electronics and Packaging Technologies for NASA Missions," 2008 International Conference on High Temperature Electronics (HiTEC), Albuquerque, NM, 2008, pp. 10-16.
- [5] Radovsky, G.; Popovitz-Biro, R.; Stroppa, D.G.; Houben, L.; Tenne, R. *Nanotubes from chalcogenide misfit compounds:* Sn–S and Nb–Pb–S. Acc. Chem. Res. 2013, 47, 406–416.
- [6] Radovsky, G.; Popovitz-Biro, R.; Staiger, M.; Gartsman, K.; Thomsen, C.; Lorenz, T.; Seifert, G.; Tenne, R. Synthesis of copious amounts of SnS2 and

SnS2/SnS nanotubes with ordered superstructures. *Angew. Chem. Int. Ed.* 2011, 50, 12316–12320.

- [7] C. Nordling, J. "Osterman, Physics Handbook for Science and Engineering, Studentlitteratur, (1999).
- [8] M. M. Martynyuk, "generalized equation of state for liquids and gases," *Zhurnal Fizicheskoi Khimii*, vol. 65, pp. 1716–1717(1991).
- [9] R. J. Sadus, "New Dieterici-Type Equations of State for Fluid Phase Equilibria," *Fluid Phase Equilibria*, vol. 212, pp. 31–39(2003).
- [10] P.A.Tamanga, D.Lissouck, F. Lontisi, and M. Tchoffo, "The Boundary Curve of Thermodynamic Stability of the Liquid Phase on the basis of a Generalised Berthelots equation," *African Journal of Sci. and Tech.*, vol. 5, pp. 1–8(2004).
- [11] A.A.Sobko, "generalized van der Waals Berthelot Equation of state,"*Doklady Physics*,**53**,No.8, 416 – 419 (2008).
- [12] S.B.Kiselev, 'Cubic Crossover Equation of State', *Fluid phase Equilib.*, 147, 7 – 23, (1998).Valderrama J.O, "A generalized Patel-Teja Equation of State for polar and Nonpolar Fluids and Mixtures,"*J.Chem.Eng. Jpn.*,23, 87-91 (1990).
- [13] K.Jibeom and J.Joonhyeon, "A Mathematical Recursive Model For Accurate Description of the Phase Behavior in the Near – critical Region by generalized van der Waals Equation," *J.Physics Conference Series*, **574**, No.1, 12006 – 12009 (2015).
- [14] R.Balasubaramanian, K.Gunavathi, R.Jegan and D.Roobanguru, "A Study on the Generlistion of Equations of State for Liquids and Gases," Open Journal of Modern Physics, 1, 54-60 (2014).
- [15] R.Balasubramanian and R.Kamala, "A New Three-Parameter generalized Berthelot Equation of State for Hydrocarbons: Superheat", *Open Science Journal of Modern Physics*, 3(1): 1-4 (2016)
- [16] R.Balasubramanian and S. Menaka, "A Study on the Thermodynamic Limit of Superheat of Refractory Metals",*International Journal of Science and Research*, Volume **5**, Issue 11,634-636 (2016).
- [17] R.Balasubramanian and C.Arul, "Thermophysical Properties of Metastable Helium-3 and Helium-4", *International Journal of Science and Research*, Volume **6**, Issue 1, 2315-2319 (2017).
- [18] R.Balasubramanian and Sugumar, "A New Three-Parameter generalized generalized van der Waals Equation of State for Water, Heavy Water, Tritium Oxide and Silica," *International Journal of Science and Research (IJSR) ISSN (Online):* 2319-7064 (2017).
- [19] Balasubramanian R and Murugesan R, "A New Four-Parameter Berthelot Equation of State: Stability Boundary of Isomers and Isotopes of Hydrogen",*International Journal of Science and Research*, **6**, Issue 8, 2282-2286, (2017).
- [20] Balasubramanian R, RamyaRayen X and Murugesan R, "Second Virial Coefficients of Noble Gases", *International Journal of Science Research*, 6, Issue 10, 1025-1029,(2017).
- [21] R.Balasubramanian, "Superheating of Liquid Alkali Metals," *Int.J. Thermophys*, **27**, 1494-1500 (2006).

Volume 7 Issue 4, April 2018

<u>www.ijsr.net</u>

- [22] M.M.Martynyuk, "Phase Explosion of a Metastable Fluid," *Combustion, Explosion, and Shock Waves* **13** (2): 178–191 (1977)
- [23] B.J. Garrison, T.E. Itina, L.V. Zhigilei, "Limit of overheating and the Threshold Behaviour in Laser Ablation,"*Phys. Rev. B*, 68, 041501 (2003).
- [24] V.I. Mazhukin, A.V. Shapranov, A.A. Samokhin, A.Yu. Ivochkin, "Mathematical Modeling of Non-Equilibrium Phase Transition in Rapidly Heated Thin Liquid Film,"*Mathematica Montisnigri*, 27, 65-90 (2013).
- [25] V.I. Mazhukin, A.V. Shapranov, A.A. Samokhin, A.V. Mazhukin and O.N. Koroleva, "Vizualization and Analysis of the Results of Molecular Dynamic Modeling of Intensive Evaporation of Liquid in the Near-critical Region," *Scientific Visualization Electronic Journal*, 6 (4), 72-95 (2014).
- [26] V.I. Mazhukin, A.A.Samokhin, A.V.Shapranov and M.M.Demin, "Modeling of Thin Film Explosive
- [27] Boiling-Surface Evaporation and Electron Thermal Conductivity Effect,"*Mater. Res. Express*, 2, 016402 (2015). [27]. V.P. Skripov, Metastable Liquids(Halsted Press, John Wiley & Sons, New York) (1974)
- [28] V.P. Skripov, M.Z. Faizullin, "Crystal–Liquid–Gas Phase Tranzition and Thermodynamic Similarity," (Wiley–VCH Verlag GmbH & Co. KGaA, Weinheim) (2006)
- [29] V.G. Baidakov, "Explosive Boiling of Superheated Cryogenic Liquids," (Wiley–VCH Verlag GmbH & Co. KGaA, Weinheim) (2007)
- [30] P.V. Skripov, A.P.Skripov, The Phenomenon of Superheat of Liquids: Memory of Vladimir P.Skripov, *Int. J. Thermophys.* **31**, 816-830 (2010)
- [31] G.V. Ermakov, "Thermodynamic Properties and Boiling-Up Kinetics of Superheated Liquids,"(UrO RAN, Ekaterinburg) (2002) [in Russian]
- [32] G.V. Ermakov, E.V. Lipnyagov, S.A. Perminov,L. Gurashkin, J. Chem. Phys. **131**, 031102 (2009)
- [33] Thermodynamic Properties of Individual Substances, Vol 3, Edited by L.V.Medveder et al., Moscow, Nauka, 1981.

Author Profile



Dr. R. Balasubramanian is an Assistant Professor of Physics at the Arignar Anna Government Arts College, Namakkal, Tamilnadu, India. He has 25 years of post- doctoral research and teaching experience. He did his Doctorate at the Russian Peoples' Friendship

University, Moscow, Russia under the supervision of renowned Professor M.M.Martynyuk who conceptualized the phenomenon of Explosive Boiling of Liquids. His research interests include Phase transitions, Critical phenomena and Equations of state.



Dr.R.Balasubramanian.

Volume 7 Issue 4, April 2018 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY